

## Versatile ruthenium(II)-catalyzed C–H cyanations of benzamides†

Weiping Liu and Lutz Ackermann\*

Cite this: *Chem. Commun.*, 2014, 50, 1878Received 15th December 2013,  
Accepted 2nd January 2014

DOI: 10.1039/c3cc49502g

www.rsc.org/chemcomm

**Direct cyanations of arenes and heteroarenes bearing only weakly coordinating amides were accomplished using a robust ruthenium(II) catalyst. The user-friendly C(sp<sup>2</sup>)–H activation occurred with the assistance of carboxylate with high site-selectivity, excellent functional group tolerance and ample scope.**

Aromatic nitriles constitute key structural motifs of important pharmaceuticals, dyes, agrochemicals and natural products.<sup>1–3</sup> The cyano group serves as a versatile functional group that can easily be transformed into amines, ketones or aldehydes, among others.<sup>4</sup> The syntheses of aryl nitriles continue to rely on classical approaches, such as the Sandmeyer<sup>5</sup> or the Rosenmund–von Braun reaction,<sup>6,7</sup> which have severe limitations, including the use of stoichiometric or even super-stoichiometric amounts of metal cyanides as well as harsh reaction conditions. An alternative approach is represented by transition metal-catalyzed cyanations of aryl halides or boronic acids under milder reaction conditions.<sup>8</sup> These catalyzed coupling reactions thereby also display a considerably improved functional group tolerance. However, this strategy exploits prefunctionalized substrates, the preparation of which requires numerous functional group interconversions and generates undesired waste.


The catalytic activation of otherwise inert C(sp<sup>2</sup>)–H bonds as latent functional groups has in recent years emerged as an increasingly viable tool for improving the atom- and step-economy of organic synthesis.<sup>9</sup> Thus, Yu developed direct cyanations of 2-arylpyridines.<sup>10,11</sup> The research groups of Jiao<sup>12</sup> and Chang<sup>13</sup> reported on oxidative cyanations of electron-rich substrates with nonmetallic cyano-group sources in the presence of stoichiometric sacrificial oxidants,<sup>14,15</sup> while Hartwig disclosed a two-step iridium-catalyzed borylation/copper-mediated cyanation protocol.<sup>16</sup> Very recently, rhodium-catalyzed C–H cyanations of arenes were accomplished, albeit exploiting rather strongly coordinating directing groups, such

as oxazolines, oximes, pyrazoles or pyridines.<sup>17–19</sup> Herein, we wish to disclose an alternative strategy, which involves the use of relatively inexpensive<sup>20</sup> ruthenium(II) catalysts<sup>21</sup> for the first time for C(sp<sup>2</sup>)–H cyanations. Importantly, the versatile ruthenium catalyst enabled expedient C–H cyanations with easily accessible *N*-cyano-*N*-phenyl-*p*-toluenesulfonamide (NCTS, 2) as the most user-friendly cyanation reagent for the functionalization of only weakly coordinating<sup>22</sup> aromatic and heteroaromatic amides – key structural scaffolds in organic synthesis and medicinal chemistry.

We initiated our studies by probing the effect of different additives and solvents on the envisioned C–H bond cyanation using synthetically useful benzamide 1a (Table 1).

Thus, [RuCl<sub>2</sub>(*p*-cymene)]<sub>2</sub> enabled the desired direct cyanation when using both AgSbF<sub>6</sub> and AgOAc as additives in DCE as the

Table 1 Optimization of C–H cyanation with benzamide 1a<sup>a</sup>

			
Entry	Additive	Solvent	Yield (%)
1	AgOAc	DCE	77 (70)
2	Cu(OAc) <sub>2</sub>	DCE	30
3 <sup>b</sup>	AgOAc	DCE	80 (67)
4 <sup>c</sup>	AgOAc	DCE	< 5
5 <sup>d</sup>	AgOAc	DCE	67 (54)
6	AgOAc	1,4-Dioxane	38
7	AgOAc	<i>o</i> -Xylene	21
8	AgOAc	Toluene	23
9	AgOAc	DMF	< 2
10 <sup>e</sup>	AgOAc	DCE	< 2
11	KOAc	DCE	< 5
12	CsOAc	DCE	17
13	NaOAc	DCE	90 (80)
14 <sup>f</sup>	NaOAc	DCE	95 (84)

<sup>a</sup> Reaction conditions: 1a (0.5 mmol), 2 (1.0 mmol), [RuCl<sub>2</sub>(*p*-cymene)]<sub>2</sub> (5.0 mol%), AgSbF<sub>6</sub> (20 mol%), additive (20 mol%), solvent (2.0 mL), 120 °C, 18 h; GC-conversion, isolated yields are given in parentheses.


<sup>b</sup> AgOAc (30 mol%). <sup>c</sup> Without AgSbF<sub>6</sub>. <sup>d</sup> AgSbF<sub>6</sub> (30 mol%). <sup>e</sup> Without [RuCl<sub>2</sub>(*p*-cymene)]<sub>2</sub>. <sup>f</sup> 24 h.

Institut fuer Organische und Biomolekulare Chemie, Georg-August-Universitaet,  
Tammannstrasse 2, 37077 Goettingen, Germany.

E-mail: Lutz.Ackermann@chemie.uni-goettingen.de

† Electronic supplementary information (ESI) available: Experimental procedures, characterization data, and <sup>1</sup>H and <sup>13</sup>C NMR spectra of products. See DOI: 10.1039/c3cc49502g





Scheme 1 Effect of *N*-substituents on C–H cyanations.

solvent (entries 1–5). DCE proved to be the solvent of choice (entries 1 and 6–9), while a test reaction illustrated that the C–H bond activation did not occur in the absence of the ruthenium catalyst (entry 10). Among a set of representative metal acetate additives,<sup>23,24</sup>  $\text{NaOAc}$  furnished the highest yields of the product **3a** (entries 11–14).

With the optimized catalytic system in hand, we explored the effect exerted by the *N*-substituent at the amide moiety (Scheme 1). A variety of tertiary amides **1a–1f** proved to be suitable substrates with optimal results being accomplished with the sterically hindered substrate **1a**.<sup>25</sup> A comparable catalytic efficacy was obtained when performing the C–H bond cyanation on a larger 5 mmol scale.

Subsequently, we tested the scope of the catalytic system for the cyanation of aromatic amides **1** (Scheme 2). The robust ruthenium(II) complex proved to be highly chemo-selective. Thus, the catalyst tolerated a set of valuable electrophilic functional groups, such as ester, fluoro, chloro, bromo and even iodo groups, which should be invaluable for the further diversification of the thus obtained products **3**.

Scheme 2 Scope of ruthenium-catalyzed C–H cyanation with amides **1**.Scheme 3 C–H cyanation with heteroaromatic amides **1**.


The user-friendly ruthenium(II) catalyst was not limited to the functionalization of arenes. Indeed, challenging heteroaromatic substrates **1p–1u** were efficiently directly functionalized both at positions C-2 (Scheme 3a) as well as C-3 (Scheme 3b). Thereby, cyanated thiophenes, furanes, benzothiophenes, benzofuranes and indoles were obtained in a chemo- and site-selective fashion.

The well-defined ruthenium(II) biscalboxylate complex **4**<sup>26</sup> displayed a catalytic activity comparable to the one observed when using the *in situ* generated system, notably even in the absence of  $\text{NaOAc}$  (Scheme 4).

In consideration of the unique chemo-selectivity and outstanding efficacy of our ruthenium catalyst, we became interested in delineating its mode of action. To this end, intramolecular competition experiments with *meta*-substituted substrates **1v–1x** revealed a considerable secondary directing group effect<sup>27</sup> exerted by a methoxy substituent and, even more pronounced, by a fluoro group present in benzamides **1w** and **1x**, respectively (Scheme 5).

Intermolecular competition experiments between differently decorated amides indicated electron-rich arenes to be preferentially converted (Scheme 6), which can be rationalized in terms of an electrophilic-type activation mode of the cationic ruthenium species.

In support of this hypothesis, we observed a rather low kinetic isotope effect (KIE) of  $k_{\text{H}}/k_{\text{D}} \approx 1.2$  (see the ESI†). Moreover, C–H bond activation conducted in the presence of  $\text{D}_2\text{O}$  as the cosolvent

Scheme 4 Well-defined complex **4** as the catalyst.



- Chem. Sci.*, 2013, **4**, 886; (e) P. B. Arockiam, C. Bruneau and P. H. Dixneuf, *Chem. Rev.*, 2012, **112**, 5879; (f) L. Ackermann, *Pure Appl. Chem.*, 2010, **82**, 1403; (g) F. Kakiuchi and S. Murai, *Acc. Chem. Res.*, 2002, **35**, 826 and cited references.
- 22 K. M. Engle, T.-S. Mei, M. Wasa and J.-Q. Yu, *Acc. Chem. Res.*, 2012, **45**, 788.
- 23 L. Ackermann, *Chem. Rev.*, 2011, **111**, 1315.
- 24 D. Lapointe and K. Fagnou, *Chem. Lett.*, 2010, **39**, 1118.
- 25 The mass balance of the cyanation reactions with substrates **1b–1f** is accounted for by the unreacted starting materials.
- 26 L. Ackermann, P. Novák, R. Vicente and N. Hofmann, *Angew. Chem., Int. Ed.*, 2009, **48**, 6045.
- 27 D. Balcells, E. Clot and O. Eisenstein, *Chem. Rev.*, 2010, **110**, 749.
- 28 Preliminary experiments with anilides or aryl carbamates as the substrates provided thus far less satisfactory results.
- 29 For recent examples of ruthenium(II)-catalyzed C–H oxygenations, see: (a) W. Liu and L. Ackermann, *Org. Lett.*, 2013, **15**, 3484; (b) F. Yang and L. Ackermann, *Org. Lett.*, 2013, **15**, 718; (c) V. S. Thirunavukkarasu and L. Ackermann, *Org. Lett.*, 2012, **14**, 6206; (d) V. S. Thirunavukkarasu, J. Hubrich and L. Ackermann, *Org. Lett.*, 2012, **14**, 4210; (e) Y. Yang, Y. Lin and Y. Rao, *Org. Lett.*, 2012, **14**, 2874.
- 30 A recent ruthenium(II)-catalyzed C–H halogenation: L. Wang and L. Ackermann, *Chem. Commun.*, 2014, **50**, 1083.

