This article was downloaded by: [University of Illinois Chicago] On: 28 November 2014, At: 12:12 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/uopp20

REINVESTIGATION OF THE SYNTHESIS OF 2-BENZYL-3-ARYL-QUINAZOLIN-4-[3H]-ONES. AN IMPROVED MULTI-COMPONENT PROCEDURE USING MICROWAVES

Anshu Dandia^a, Ruby Singh^a & Pritima Sarawgi^a ^a Department of Chemistry, University of Rajasthan, Jaipur, 302004, INDIA E-mail: Published online: 06 Feb 2009.

To cite this article: Anshu Dandia , Ruby Singh & Pritima Sarawgi (2005) REINVESTIGATION OF THE SYNTHESIS OF 2-BENZYL-3-ARYL-QUINAZOLIN-4-[3H]-ONES. AN IMPROVED MULTI-COMPONENT PROCEDURE USING MICROWAVES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 37:4, 397-402, DOI: <u>10.1080/00304940509354972</u>

To link to this article: <u>http://dx.doi.org/10.1080/00304940509354972</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

OPPI BRIEFS

REINVESTIGATION OF THE SYNTHESIS OF 2-BENZYL-3-ARYL-QUINAZOLIN-4-[*3H*]-ONES. AN IMPROVED MULTI-COMPONENT PROCEDURE USING MICROWAVES[†]

Anshu Dandia*, Ruby Singh and Pritima Sarawgi

Submitted by (04/28/04)

Department of Chemistry, University of Rajasthan Jaipur-302004, INDIA e-mail: dranshudandia@eth.net

4[3H]-Quinazolinones have wide ranging biological activities^{1,2} and have been used as anti-convulsant drugs with anti-inflammatory, analgesic and hypnotic properties.³⁻⁵ Some 4[3H]-quinazolinones also display CNS, anti-fungal, antibacterial and anti-microbial activities^{6,7} and are also useful as anti-tumor agents.⁸ 2,3-Disubstituted quinazolones have been associated with antiviral, antibacterial and antifungal activity.^{9,10}

An earlier reported conventional synthesis of 2,3-disubstituted quinazolin-4(3*H*)-ones (4) involves two steps,¹¹ cyclodehydration of 2-benzamidobenzoic acid (1) with excess acetic anhydride under anhydrous conditions to give benzoxazin-4-one (2) followed by reflux of 2 with amines in glacial acetic acid or pyridine.¹² The products were obtained in moderate yields and the reaction requires 10-12 h refluxing. In view of our desire to synthesize a series of 2,3-disubstituted quinazolones (4) with pharmacophoric groups and to establish structure-activity relationship in quinazolones, the condensation of benzoxazin-4-one with substituted anilines was studied. While *m*- and *p*-substituted anilines gave the expected products 4, the *o*-substituted anilines afforded the intermediate *o*-acylaminobenzanilides (3), in contrast to earlier report of Mishra *et al.*¹¹ The synthesis of a compound with the same substituent ($\mathbf{R} = o$ -nitrophenyl) was repeated under the conditions specified by Mishra *et al.*, and the products were identified as 3 and not as quinazolones (4). Hence, the method was found to be unsuitable for the reaction involving *o*-substituted anilines. This observation is in agreement with the report of Zentmyer *et al.*¹³

Microwave irradiation is an alternate energy source whose popularity and synthetic utility in organic chemistry has increased considerably in recent years.¹⁴ Solvent-free conditions are especially suitable for microwave irradiation. A literature survey reveals examples of certain reactions, which do not occur under conventional conditions but become possible under microwave irradiation.¹⁵ Recently multi-component reactions (MCRs) constitute an especially attractive synthetic strategy for the rapid and efficient generation of libraries of compound.¹⁶

© 2005 by Organic Preparations and Procedures Inc.



i) Stirring (room temp.); ii) Ac₂O, Δ ; iii) HOAc, *o*-substituted anilines; iv) HOAc, *m*- and *p*-substituted anilines Scheme 1

The synthesis of 4[3H]-quinazoline-4-one core nucleus using microwaves has been reported from the reaction of anthranilic acid with formamide.¹⁷ Herein we describe an improved procedure using the multi-component reaction of anthranilic acid, acid chlorides and amines under microwave irradiation using neat reaction conditions. The results showed that the reaction occurred in a shorter time with facile work-up and thus provides a general route to the synthesis of 4[3H]-quinazolinones (4) with a variety of amines and acid chlorides.

-		• • •			
Yield	mp	Elemen	Elemental Analysis (Found)		
(%)	(°Ċ)	С	Н	Ν	
65	166/16511	67.19(67.01)	4.56(4.55)	11.19(11.17)	
68	174-176	63.17(63.00)	4.04(4.03)	7.02(7.01)	
62	170-172	76.72(76.52)	5.85(5.82)	8.13(8.11)	
	Yield (%) 65 68 62	Yield mp (%) (°C) 65 166/165 ¹¹ 68 174-176 62 170-172	Yield mp Element (%) (°C) C 65 166/165 ¹¹ 67.19(67.01) 68 174-176 63.17(63.00) 62 170-172 76.72(76.52)	Yield mp Elemental Analysis (I (%) (°C) C H 65 166/165 ¹¹ 67.19(67.01) 4.56(4.55) 68 174-176 63.17(63.00) 4.04(4.03) 62 170-172 76.72(76.52) 5.85(5.82)	

Table 1. Physical and Analytical Data of Compounds (3a-c)

The formation of *o*-acylaminobenzanilide **3a-c** and 4(3H)-quinazoline-4-one **4a-n** was confirmed on the basis of spectral studies. The IR spectra of **3a-c** displayed characteristic bands at 3360-3400 (NH), 1700 and 1680 (both C=O) cm⁻¹. IR spectra of quinazolones **4a-n** showed only one carbonyl band at 1700 and 1605-1610 (C=N) cm⁻¹ confirming the ring closure in all compounds. The detailed spectral data of compound **3a-c** and **4a-n** are given in *Table 3*. The mass



a) $R = PhCH_2$, $R' = 2-NO_2 \bullet C_6H_4$; b) $R = PhCH_2$, $R' = 2,6-Cl_2 \bullet C_6H_3$; c) $R = PhCH_2$, $R' = 2-Me \bullet C_6H_4$; d) $R = PhCH_2$, $R' = 3-Cl \bullet C_6H_4$; e) $R = PhCH_2$, $R' = 4-Me \bullet C_6H_4$; f) R = Ph, $R' = 4-Cl \bullet C_6H_4$; g) R = Ph, $Ar = 2-Cl \bullet C_6H_4$; h) R = Ph, $R' = 4-Br \bullet C_6H_4$; i) R = Ph, $R' = 3-CF_3$, $4-Cl \bullet C_6H_3$; j) $R = CH_3$, $R' = 2-F \bullet C_6H_4$; k) $R = CH_3$, $R' = 4-NO_2 \bullet C_6H_4$; l) R = Ph, $R' = C_2H_5$; m) R = Ph, $R' = 3-HO \bullet C_6H_4$; n) R = Ph, $R' = 4-CH_3O \bullet C_6H_4$

Scheme 2

OPPI BRIEFS

spectrum of representative compounds 3a and 4a exhibited the molecular ion peaks at m/z 375 and 357 (100%) which correspond to the molecular weight of compounds 3a and 4a, respectively.

Cmpd	Yield	Time	mp	Elemental Analysis (Found)			
•	(%)	(min/hs)	(°Č)	С	H	Ν	
4 a	88	5	156-157	70.58(70.39)	4.23(4.22)	11.76(11.73)	
4b	91	6	124-126	66.16(65.96)	3.70(3.71)	7.35(7.33)	
4c	93	5	116-118	80.96(80.74)	5.56(5.54)	8.58(8.60)	
4d	90ª/61 ^b	4ª/12 ^b	164-166	72.73(72.52)	4.36(4.37)	8.08(8.06)	
4e	92ª/60 ^b	5ª/12 ^b	110-112	80.96(80.74)	5.56(5.55)	8.58(8.56)	
4f	92	5	183-185	72.18(72.00)	3.94(3.93)	8.42(8.40)	
4g	90	4	140-142	72.18(72.38)	3.94(3.93)	8.42(8.44)	
4h	91	5	220-222	63.68(63.48)	3.47 (3.46)	7.43(7.41)	
4i	89	4	170-172	62.93(62.75)	3.02(3.01)	6.99(6.97)	
4j	90	4	225-227	70.86(70.65)	4.36(4.35)	11.02(10.98)	
4k	90	5	112-114 ¹²	64.05(64.20)	3.94(3.93)	14.94(14.90)	
41	92	4	230-231	76.78(76.58)	5.64(5.63)	11.19(11.16)	
4m	89	5	235-237	76.42(76.20)	4.49(4.50)	8.91(8.93)	
4n	90	5	303-305	76.81(76.60)	4.91(4.92)	8.53(8.51)	

Table 2.	Physical	and Analy	tical Data o	of Com	pounds ((4a-n))
----------	----------	-----------	--------------	--------	----------	--------	---

a) Yield and time (min) correspond to microwave method. b) Yield and time (hs) correspond to conventional method.

EXPERIMENTAL SECTION

Mps were determined in open glass capillary and are uncorrected. IR spectra were recorded on a Perkin-Elmer (model-577) (KBr). ¹H NMR and ¹³C NMR were recorded on Jeol model FX 90Q and Bruker-DRX-300 using CDCl₃ as solvent and TMS as an internal reference at 89.55 and 75.47 MHz, respectively. Mass spectra of representative compounds were obtained on Kratos 50 mass spectrometer at 70 eV. The purity of all compounds was checked by TLC using silica gel 'G' coated glass plates and benzene-ethylacetate (8;2) as eluent. The microwave-induced reactions were carried out in BMO-700T modified multimode oven fitted with a condenser and a magnetic stirrer.

Synthesis of 2-Phenylmethyl-3-(2-nitrophenyl)quinazolin-4(3H)-one(4a).- To an equimolar mixture of anthranilic acid (2 mmol) and phenylacetyl chloride (2 mmol) contained in an Erlenmeyer flask fitted with a condenser, 2-nitroaniline (2 mmol) was added slowly. The flask was then placed in the microwave oven and irradiated for 5 min (TLC) at 640 watts. The reaction mixture was cooled at room temperature to give a solid, which was recrystallized from ethanol to give 4a. All compounds 4b-n listed in *Table 1* were synthesized similarly in high yields and

reduced time. In case of reaction of m-amino- phenol, the sticky product obtained after irradiation of reaction mixture was triturated with pet-ether and recrystallized from ethanol to give crystals of 4m.

Table 3	8. ¹ H and ¹³ C NMR Spectra of Compounds	(3a-c) and (4a-n)
Cmpd	¹ H NMR (δ)	¹³ C NMR (δ)
3a	4.21 (s, 2H, $C\underline{H}_2Ph$), 6.90-8.21 (m, 13H, Ar-H), 8.99 & 10.12 (two bs, two NH, D_2O exchangeable)	170.2, 169.2 (two C=O), 142.8-119.2 (aromatic carbons), 45.9 (CO- <u>C</u> H ₂)
3b	4.24(s, 2H, $C\underline{H}_2$ Ph), 6.98-8.18(m, 12H, Ar-H), 9.01 & 10.15(two bs, two NH, D_2O exchangeable)	171.1, 168.2 (two C=O), 145.2-118.9 (aromatic carbons), 44.8 (CO- <u>C</u> H ₂)
3c	2.24(s, 3H, CH ₃), 4.22(s, 2H, C $\underline{\text{H}}_2$ Ph), 6.92-8.16(m, 13H, Ar-H), 8.96 & 10.12 (two bs, two NH, D ₂ O exchangeable)	169.8, 168.0 (two C=O), 141.2-117.7 (aromatic carbons), 44.1 (CO- <u>C</u> H ₂), 19.2 (CH ₃)
4 a	3.93 (s, 2H, C <u>H</u> ₂ Ph), 6.85-7.15 (m, 5H CH ₂ <u>Ph</u>), 7.20-8.15 (m, 7H, Ar-H), 8.36 (dd, 1H, 5-H)	169.5 (C=O), 157.1 (C=N), 142.8-119.2 (aromatic carbons), 23.4 (<u>C</u> H ₂ Ph)
4 b	3.93 (s, 2H, C <u>H</u> ₂ Ph), 6.89-7.16 (m, 5H CH ₂ <u>Ph</u>), 7.22-8.10 (m, 6H, Ar-H), 8.35 (dd, 1H, 5-H).	168.9 (C=O), 156.2 (C=N) 144.5-118.4 (aromatic carbons), 24.4 (<u>C</u> H ₂ Ph)
4 c	2.21 (s, 3H, CH ₃), 3.90 (s, 2H, C <u>H₂Ph)</u> , 6.86-7.18 (m, 5H CH ₂ <u>Ph</u>), 7.20-8.14 (m, 7H, Ar-H), 8.34 (dd, 1H, 5-H)	170.1 (C=O), 157.3 (C=N) 141.5-116.4 (aromatic carbons), 24.6 ($\underline{C}H_2Ph$), 18.2 (CH ₃)
4 d	3.93 (s, 2H, $C\underline{H}_2Ph$), 6.85-7.20 (m, 5H $CH_2\underline{Ph}$), 7.23-8.16 (m, 7H, Ar-H), 8.30 (dd, 1H, 5-H)	169.5 (C=O), 156.8 (C=N), 143.5-117.4 (aromatic carbons), 25.1 (<u>C</u> H ₂ Ph)
4e	2.18 (s, 3H, CH ₃), 3.92 (s, 2H, C \underline{H}_2 Ph), 6.88-7.16 (m, 5H CH ₂ <u>Ph</u>), 7.20-8.10 (m, 7H, Ar-H), 8.31 (dd, 1H, 5-H)	168.8 (C=O), 156.9 (C=N) 142.5-118.2 (aromatic carbons), 24.8 (<u>C</u> H ₂ Ph), 18.8 (CH ₃)
4f	6.86-7.41 (m, 5H, Ph), 7.50-8.01 (m, 7H, Ar-H), 8.12 (dd, 1H, 5-H)	170.6 (C=O), 157.2 (C=N), 143.2-118.2 (aromatic carbons)
4g	6.89-7.44 (m, 5H, Ar-H), 7.48-8.09 (m, 7H, Ar-H), 8.17 (dd, 1H, 5-H)	171.2 (C=O), 156.8 (C=N) 144.5-119.2 (aromatic carbons)
4h	6.91-7.48 (m, 5H, Ar-H), 7.51-8.12 (m, 7H, Ar-H) 8.20 (dd, 1H, 5-H)	172.4 (C=O), 158.1 (C=N), 143.6-120.3 (aromatic carbons)
4 i	6.84-7.38(m, 5H, Ar-H) and 7.45-7.01 (m, 6H, Ar-H), 8.10(dd, 1H, 5-H)	
4j	.96 (s, 3H, CH ₃), 6.51-7.88 (m, 7H, Ar-H), 7.99 (dd, 1H, 5-H)	171.8 (C=O), 159.2 (C=N), 142.8-119.8 (aromatic carbons), 17.0 (CH ₃)
4k	.98 (s, 3H, CH ₃), 6.54-7.92 (m, 7H, Ar-H), 8.05 (dd,1H, 5-H)	

Table 3. Continued...

Cmpd ¹H NMR (δ)

- **4l** 1.22 (t, 3H, CH₃), 3.24 (q, 2H, CH₂), 7.29-7.90 (m, 8H, Ar-H), 8.14 (dd, 1H, 5-H)
- **4m** 8.78 (s, 1H, OH, D₂O exchangeable), 6.71-7.47 (m, 5H, Ar-H) and 7.52-7.99 (m, 7H, Ar-H), 8.19 (dd, 1H, 5-H)
- **4n** 4.24 (s, 1H, OCH₃), 6.88-7.38 (m, 5H, Ar-H), 7.48-8.09 (m, 7H, Ar-H), 8.17 (dd, 1H, 5-H)

¹³C NMR (δ) 170.8 (C=O), 158.3 (C=N), 142.8-124.2 (aromatic carbons), 33.4 (CH₂), 12.8 (CH₃)

172.1 (C=O), 160.2 (C=N), 145.8-122.1 (aromatic carbons)

171.8 (C=O), 161.8 (C=N), 143.8-120.8 (aromatic carbons), 52.4 (OCH₃)

Acknowledgement.- Financial assistance from UGC and CSIR (No 01(1907)/03/EMR-II), New Delhi is acknowledged and one of author (R. S.) is thankful for the award of a Senior Research Fellowship.

REFERENCES

- [†]Presented at the Seventh International Electronic Conference on Microwave Chemistry (ECSOC-7), 1-30 Nov. 2003.
- 1. M. Hori, R. Iemura, H. Hara and A. Ozaki, Chem. Pharm. Bull., 38, 1286 (1990).
- J. B. Jiang, D. P. Hesson, B. A. Dusak, D. L. Dexter, G. J. Kang and E. Hamel, J. Med. Chem., 33, 1721(1990).
- 3. N. A. Santagati, E. Bousquet, A. Spadaro and G. Ronsisvalle, Il Farmaco, 54, 780 (1999).
- A. Hitkari, M. Bhalla, A. K. Saxena, M. Verma, M. P. Gupta and K. Shanker, *Bull. Chim. Farm.*, **134**, 604 (1995).
- 5. A. A. Bekhit and A. Khalil, *Pharmazie*, **53**, 539 (1998).
- 6. G. Capan, N. Ergenc, S. Bueyuektimkin and N. Yulug, Sci. Pharm., 61, 243 (1993).
- 7. R. Lakhan and J. Rai Babban, J. Chem. Eng. Data, 32, 384 (1987).
- M. J. Deetz, J. P. Malerich, A. M. Beatty and B. D. Smith, *Tetrahedron Lett.*, 42, 1851 (2001).
- 9. V. K. Pandey and H. S. Negi, Biol. Mem., 25, 29 (1999).
- 10. E. Feky and H. Said, Pharmazie, 48, 894 (1993).
- 11. P. Mishra, S. Jain and S. Jain, J. Indian Chem. Soc., 74, 816 (1997).

- 12. P. Mishra, P. N. Gupta and A. K. Shakya, *J. Indian Chem. Soc.*, **68**, 618 (1991); P. Mishra, P. Paneerselvam and S. Jain, *J. Indian Chem. Soc.*, **72**, 559 (1995).
- 13. D. T. Zentmyer and E. C. Wagner, J. Org. Chem., 14, 967 (1949).
- S. Caddick, *Tetrahedron*, **51**, 10403 (1995); R. S. Varma, *Green Chemistry*, 43(1999); P. Lidstron, J. Tierney, B. Wathey and J. Westman, *Tetrahedron Lett.*, **57**, 9225 (2001).
- B. Boruah, J. Boruah, D. Prajapati, J. C. Sandhu and A. C. Gosh, *Tetrahedron Lett.*, 37, 4203 (1996); B. Garringnes, C. Laporte, R. Laurnte, A. Laporterie and J. Dubac, *Ann.*, 739 (1996).
- K. Groebke, L. Weber and F. Mehlin, *Synlett.*, 661, (1998); R. W. Armstrong, A. P. Combs, P. A. Tempest, S. D. Brown and T. A. Keating, *Acc. Chem. Res.*, 26, 123 (1996); R. S. Varma, *Green Chemistry*, 93 (2003); C. O. Kappe, D. Kumar, R. S. Varma, *Synthesis*, 1799 (1999); R. S. Varma and D. Kumar, *Tetrahedron Lett.*, 40, 7665 (1999).
- 17. F. R. Alexandre, A. Berecibar and T. Besson, Tetrahedron Lett., 43, 3911 (2002).
- A. Dandia, R. Singh and K. Arya, Org. Prep. Proced. Int., **35**, 387 (2003); A. Dandia, M. Sati, K. Arya and A. Loupy, *Heterocycles*, **60**, 563 (2003); A. Dandia, M. Sati and A. Loupy, Green Chemistry, **4**, 599 (2002); A. Dandia, H. Taneja and R. Singh, J. Chem. Res. (S), 272 (**2000**); A. Dandia, H. Sachdeva and R. Singh, Synth. Commun., **31**, 1879 (2001).
