


# Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: F. Zhang, W. Xu, Q. Li, C. Cao and H. zheng, *Org. Biomol. Chem.*, 2015, DOI: 10.1039/C5OB00577A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

communication

# Intramolecular Oxidative Coupling: I<sub>2</sub>/TBHP/NaN<sub>3</sub>-mediated Synthesis of Benzofuran Derivatives

Wengang Xu, Qingcui Li, Chuanpeng Cao, Fanglin Zhang\* and Hua Zheng\*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000

A novel intramolecular oxidative coupling reaction is established to prepare benzofuran derivatives via direct C(sp<sup>2</sup>)-H functionalization for the formation of C-O bond. This transformation is mediated by I<sub>2</sub>/TBHP/NaN<sub>3</sub> under metal-free conditions and catalytic amount of NaN<sub>3</sub> plays a crucial role in the reaction. Further, the reaction shows a broad substrate scope with middle to excellent yields.

Oxidative coupling reaction is a novel strategy to construct C-C bond<sup>1</sup> and C-heteroatom bond<sup>2</sup>, which avoids pre-functionalization of easily available chemicals while improving the atom economy and step economy and has received significant interest over the last ten years. Especially, the efficient construction of useful heterocyclic compounds by using oxidative coupling reaction is an important research area in organic synthesis.<sup>3</sup> Numerous methods have been developed for carbon-heteroatom bond formation in last decades.<sup>4</sup> It is well-known that traditional oxidative coupling reactions are most dependent on transition metal catalyst. As a result, residual transition metal contaminant seriously limits their applications. Consequently, an economical, environmentally friendly and metal-free strategy is drawing many chemists' attention.<sup>5</sup>


Benzofuran and their derivatives are the core structure of many clinical drugs and natural products, which have attracted the exploration of their synthesis methods. Molecules containing these structural units have been shown to exhibit activities against cancer, plasmodium falciparum and so on.<sup>6</sup> Series of methods for the synthesis of benzofuran derivatives have been reported.<sup>7</sup> However, investigation on direct C-H functionalization and C-O bond formation of benzofuran derivatives remains rare.

To date, a lot of novel strategies of C-H functionalization and C-O bond formation via oxidative coupling make a big breakthrough.<sup>8</sup> However, there are extreme conditions in oxidative coupling reaction with phenolic hydroxyl due to phenols are ready to undergo oxidative dearomatization to yield quinones.<sup>9</sup> To our best knowledge, oxidative C(sp<sup>2</sup>)-H functionalization and C-O bond formation with phenolic hydroxyl generally falls into one of the following types. Type I, early in 2000, Boudet and his co-workers<sup>10</sup> revealed that 4-hydroxycinnamyl aldehydes incorporate into lignins by examining transgenic plants via intermolecular radical coupling reactions (Scheme 1). And authors think that this process appears to reflect simple chemical coupling propensities. Type II,


Alessandra Lattanzi<sup>11</sup> and Duan<sup>12</sup> separately developed a new strategy for the synthesis of benzofuran derivatives via C(sp<sup>2</sup>)-O coupling reaction (Type II, a). However, their methods are only efficient for electron-rich systems. Later, liu<sup>13</sup>, Yoshikai<sup>14</sup> and Zhu<sup>15</sup> reported Pd or Cu catalyzed C(sp<sup>2</sup>)-H functionalization of 2-arylphenols via oxidative cyclization respectively (Type II, b). At present, there is no report describing intramolecular oxidative functionalization of electron-deficient systems for preparing benzofuran derivatives under metal-free conditions. Accordingly, exploration of this novel oxidative coupling methods is still of great significance.

## Previous work

Type I Intermolecular C(sp<sup>2</sup>)-H bond functionalization and C-O bond formation




Type II Intramolecular C(sp<sup>2</sup>)-H bond functionalization and C-O bond formation



## This work

Intramolecular C(sp<sup>2</sup>)-H bond functionalization and C-O bond formation




**Scheme 1** Different kinds of C-H Functionalization

Recently, I<sub>2</sub> (iodine) system<sup>16</sup> has received considerable attention as a mild, non-toxic and selective reagent in organic synthesis. As a continuous study on the direct C-H functionalization and C-O bond formation, we herein describe an efficient I<sub>2</sub>-catalyzed intramolecular oxidative coupling reaction of 2-hydroxychalcones under metal-free conditions. In addition, catalytic amount of NaN<sub>3</sub> plays a crucial role in the oxidative

coupling reaction and a new plausible mechanism which may be virtually through C(sp<sup>3</sup>)-H functionalization has been proposed.

**Table 1** Optimization of the Reaction Conditions<sup>a</sup>

|  |                     |                                    |                                     |                      |                    |
|---|---------------------|------------------------------------|-------------------------------------|----------------------|--------------------|
| Entry   | Oxidant             | Additive                           | Solvent                             | Catalyst             | Yield <sup>b</sup> |
| 1   | TBHP <sup>c</sup>   | NaN <sub>3</sub>                   | CH <sub>3</sub> CN                  | I <sub>2</sub>       | 40                 |
| 2   | /                   | NaN <sub>3</sub>                   | CH <sub>3</sub> CN                  | I <sub>2</sub>       | NO                 |
| 3   | TBHP                | /                                  | CH <sub>3</sub> CN                  | I <sub>2</sub>       | NO                 |
| 4   | TBHP                | NaN <sub>3</sub>                   | CH <sub>3</sub> CN                  | /                    | NO                 |
| 5   | Others <sup>d</sup> | NaN <sub>3</sub>                   | CH <sub>3</sub> CN                  | I <sub>2</sub>       | NO                 |
| 6   | TBHP                | Others <sup>e</sup>                | CH <sub>3</sub> CN                  | I <sub>2</sub>       | NO                 |
| 7   | TBHP                | NaN <sub>3</sub>                   | CH <sub>3</sub> NO <sub>2</sub>     | I <sub>2</sub>       | NO                 |
| 8   | TBHP                | NaN <sub>3</sub>                   | DCE                                 | I <sub>2</sub>       | NO                 |
| 9   | TBHP                | NaN <sub>3</sub>                   | dioxane                             | I <sub>2</sub>       | NO                 |
| 10  | TBHP                | NaN <sub>3</sub>                   | CH <sub>3</sub> CN <sup>f</sup>     | I <sub>2</sub>       | 60                 |
| 11  | TBHP                | NaN <sub>3</sub>                   | toluene                             | I <sub>2</sub>       | 20                 |
| 12  | TBHP                | NaN <sub>3</sub>                   | C <sub>2</sub> H <sub>5</sub> OH    | I <sub>2</sub>       | 87                 |
| 13  | TBHP                | NaN <sub>3</sub> <sup>g</sup>      | C <sub>2</sub> H <sub>5</sub> OH    | I <sub>2</sub>       | 87                 |
| 14  | <b>TBHP</b>         | <b>NaN<sub>3</sub><sup>h</sup></b> | <b>C<sub>2</sub>H<sub>5</sub>OH</b> | <b>I<sub>2</sub></b> | <b>88</b>          |
| 15  | TBHP                | NaN <sub>3</sub> <sup>h</sup>      | C <sub>2</sub> H <sub>5</sub> OH    | Others <sup>i</sup>  | NO                 |
| 16  | TBHP                | NaN <sub>3</sub> <sup>h</sup>      | C <sub>2</sub> H <sub>5</sub> OH    | KI                   | 78                 |
| 17  | TBHP                | NaN <sub>3</sub> <sup>h</sup>      | C <sub>2</sub> H <sub>5</sub> OH    | TBAI                 | 83                 |

<sup>a</sup>Reaction conditions: **1a** (0.25 mmol), oxidant (0.5 mmol), additive (1 mmol) and catalyst (0.025 mmol) in solvent (4.0 mL) at 80 °C, unless otherwise stated.

<sup>b</sup>Isolated yields. NO means that no product was observed. <sup>c</sup>All TBHP (70% in water). <sup>d</sup>Other oxidants include PIDA, mCPBA, K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>, DTBP. <sup>e</sup>Other additives include K<sub>2</sub>CO<sub>3</sub>, Na<sub>2</sub>CO<sub>3</sub>, NaHCO<sub>3</sub>, NaOH, NaH<sub>2</sub>PO<sub>4</sub>·2H<sub>2</sub>O, KH<sub>2</sub>PO<sub>4</sub>, K<sub>3</sub>PO<sub>4</sub>·3H<sub>2</sub>O, KSCN, (NH<sub>4</sub>)<sub>3</sub>PO<sub>3</sub>·3H<sub>2</sub>O. <sup>f</sup>CH<sub>3</sub>CN:H<sub>2</sub>O=1:1. <sup>g</sup>2.0 equivalent of NaN<sub>3</sub> was used. <sup>h</sup>0.2 equivalent of NaN<sub>3</sub> was used. <sup>i</sup>Other acids include p-TSA, BF<sub>3</sub>·Et<sub>2</sub>O, TfOH, FeCl<sub>3</sub>, CuCl<sub>2</sub>.

Initially, trans-2-hydroxychalcone (**1a**, 0.25 mmol) was used as the model substrate for the synthesis of desired product (**2a**). As shown in table 1, **2a** was obtained in 40% yield in the presence of catalytic amount of I<sub>2</sub> (iodine), 0.5 mmol TBHP (tert-butylhydroperoxide, 70% in water) and 1 mmol NaN<sub>3</sub> (Table 1, entry 1). In the absence of I<sub>2</sub>, TBHP or NaN<sub>3</sub>, no product was obtained (Table 1, entries 2-4). To further optimize the conditions, different oxidants, additives and catalysts were tested. Employing other oxidants: PIDA (iodobenzene diacetate), mCPBA (m-chloroperoxybenzoic acid), K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> or DTBP (tert-butyl peroxide), unfortunately, yielded no product (Table 1, entry 5). Using other additives instead of NaN<sub>3</sub>, we also did not observe desired product (Table 1, entry 6). Solvent screening showed that EtOH, MeCN/H<sub>2</sub>O, and toluene were also effective for the reaction, while DCE and 1,4-dioxane were ineffective for the reaction (Table 1, entries 7-12). Decreasing NaN<sub>3</sub> to 0.5 mmol made no difference to the yield (Table 1, entry 13). Catalytic amount of NaN<sub>3</sub> (0.05 mmol) was employed out of concern that NaN<sub>3</sub> could be recycled in reaction process and, as expected, target product was obtained in good yield (Table 1, entry 14). In

addition, when I<sub>2</sub> was replaced by other acids, including CuCl<sub>2</sub>, FeCl<sub>3</sub>, BF<sub>3</sub>·Et<sub>2</sub>O, p-TSA and TfOH, no product was observed (Table 1, entry 15). What's more, when KI or TBAI (tetrabutylammonium iodide) was used, the desired coupling product **2a** was also obtained in the yield lesser than I<sub>2</sub> as the catalyst (Table 1, entries 16 and 17). On the basis of these results, the optimized reaction conditions were concluded to be 0.1 equivalent of I<sub>2</sub>, 0.2 equivalent of NaN<sub>3</sub> and 2.0 equivalent of TBHP in C<sub>2</sub>H<sub>5</sub>OH at 80 °C.

**Table 2** Exploring Generality and Scope of the Novel Reaction<sup>a,b</sup>

$$\text{1} \xrightarrow[\text{C}_2\text{H}_5\text{OH}]{\begin{matrix} 10\% \text{ I}_2 \\ 20\% \text{ NaN}_3 \\ 2\text{equiv. t-BuOOH} \end{matrix}} \text{2}$$

2a(88%)

2b(94%)

2c(82%)

2d(92%)

2e(84%)

2f(73%)

2g(49%)

2h(82%)

2i(85%)

2j(51%)

2k(86%)

2l(63%)

2m(98%)

2n(85%)

2o(61%)

2p(48%)

2q(87%)

2r(86%)


2s(75%)

<sup>a</sup>All the reactions were carried out using **1** (0.25 mmol), I<sub>2</sub> (0.025 mmol), TBHP (0.5 mmol) and NaN<sub>3</sub> (0.05 mmol) in C<sub>2</sub>H<sub>5</sub>OH (4.0 mL) at 80 °C.

<sup>b</sup>Isolated yields.


With the optimal reaction conditions in hand, we then examined the transformation of trans-2-hydroxychalcones equipped with a variety of substituents to explore generality and

scope. It was observed that substituents on the R<sup>1</sup> ring almost led to good yields (Table 2, **2b-2f**, **2h**, **2i**, **2k** and **2l**). O-methyl and p-methyl conducted almost complete transformation (Table 2, **2b** and **2d**). M-methyl, p-methoxyl, m-bromo, o-bromo and o-chloro gave target products in >80% yields (Table 2, **2c**, **2e**, **2h** and **2k**), while p-Br, p-Cl led to the corresponding products in moderate yields (Table 2, **2g** and **2j**). When changing the substituents on R<sup>2</sup> Ring, different results appeared (Table 2, **2m-2p**). Substrates equipped with 5-Cl or 5-Br gone through great transformation. Similarly, substrate equipped with methoxyl got middle yield. When R<sup>2</sup> was a strong electron-withdrawing group (5-NO<sub>2</sub>), trace desired product was observed under the optimized reaction conditions. Remarkably, this strategy was further successfully applied to heterocyclic substrates to synthesize corresponding products in good yields (Table 2, **2q-2s**).



**Scheme 2** Further Application of the Novel Reaction

We further applied this practical procedure to the trans-2-hydroxychalcone substituted by alkyl groups (Scheme 3). An interesting result occurred. When alkyl groups linked to the carbonyl group with primary carbon or secondary carbon, no desired products was isolated. However, trans-2-hydroxychalcone bearing tertiary carbon was successfully transformed to 1-(benzofuran-2-yl)-2-methylpropan-1-one in 48% yield (Scheme 3, **2t**).



**Scheme 3** Plausible Mechanism

According to our experimental data, when other additives were employed instead of NaN<sub>3</sub>, no desired product was obtained (Table 1, entry 6). Besides, target product was got while catalytic amount of NaN<sub>3</sub> was used (Table 1, entry 14). On the basis of these results, a plausible mechanism was shown in Scheme 4. First, NaN<sub>3</sub> as a nucleophile reacts with **1a** via Michael-type addition reaction to deliver unstable intermediate **A**. This transformation efficiency could be improved in the presence of catalytic amount of iodine. Subsequently, **A** could easily be oxidized by iodine to give intermediate **B**, which is converted to the product **C** after loss of NaI. Then, **C** losses HN<sub>3</sub> catalyzed by I<sub>2</sub>, which is similar to loss H<sub>2</sub>O. At last, NaI and HN<sub>3</sub> exchanged and oxidized by TBHP to I<sub>2</sub> and NaN<sub>3</sub>. Obviously, formation of **2a** was more favorable and fast process.

In summary, this paper described a novel and efficient method for the synthesis of benzofuran derivatives under metal-free conditions. In this transformation, a broad substrate scope has been demonstrated and catalytic amount of NaN<sub>3</sub> plays a crucial role in the oxidative coupling reaction. The possible domino Michael addition and intramolecular oxidative coupling reaction mechanism is also proposed. Studies on novel oxidative functionalization are being actively pursued in this laboratory.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (51273156).

## Notes and references

- <sup>a</sup>School of Chemistry, Chemical Engineering and Life Sciences, Wuhan University of Technology, Wuhan, 430070, Hubei, China. Fax: (+86)027 8774 9300; Tel: (+86)027 8774 9300; E-mail: fanglinzhang0210@gmail.com or zhenghua.whut@126.com
- †Electronic Supplementary Information (ESI) available: See DOI: 10.1039/b000000x/
- (a) J. A. Ashenhurst, *Chem. Soc. Rev.*, 2010, **39**, 540-548. (b) W. Shi, C. Liu and A. Lei, *Chem. Soc. Rev.*, 2011, **40**, 2761-2776.
- (a) A. Corma, A. Leyva-Perez and M. J. Sabater, *Chem. Rev.*, 2011, **111**, 1657-1712. (b) I. P. Beletskaya and V. P. Ananikov, *Chem. Rev.*, 2011, **111**, 1596-1636.
- (a) W. Gong, L. Xu, T. Ji, P. Xie, X. Qi, C. U. Pittman and A. Zhou, *RSC Adv.*, 2014, **4**, 6854-6857. (b) P. Li, J. Zhao, C. Xia and F. Li, *Org. Lett.*, 2014, **16**, 5992-5995.
- (a) J. Yuan, C. Liu and A. Lei, *Chem. Commun.*, 2015, **51**, 1394-1409. (b) Q. Wang, H. Zheng, W. Chai, D. Chen, X. Zeng, R. Fu and R. Yuan, *Org. Biomol. Chem.*, 2014, **12**, 6549-6553. (c) X. Li, B. Li, J. You and J. Lan, *Org. Biomol. Chem.*, 2013, **11**, 1925-1928.
- (a) J. Zhao, P. Li, C. Xia and F. Li, *Chem. Commun.*, 2014, **50**, 4751-4754. (b) R. Samanta, K. Matcha and A. P. Antonchick, *Eur. J. Org. Chem.*, 2013, **2013**, 5769-5804.
- (a) Z. Yu, J. A. Brannigan, D. K. Moss, A. M. Brzozowski, A. J. Wilkinson, A. A. Holder, E. W. Tate and R. J. Leatherbarrow, *J. Med. Chem.*, 2012, **55**, 8879-8890. (b) R. E. Ziegert, J. Torang, K. Knepper and S. Brase, *J. Comb. Chem.*, 2005, **7**, 147-169. (c) B. L. Flynn, E. Hamel and M. K. Jung, *J. Med. Chem.*, 2002, **45**, 2670-2673. (d) B. Carlsson, B. N. Singh, M. Temciuc, S. Nilsson, Y. L. Li, C. Mellin and J. Malm, *J. Med. Chem.*, 2002, **45**, 623-630.
- (a) J. Chen, J. Li and W. Su, *Org. Biomol. Chem.*, 2014, **12**, 4078-4083. (b) Y. Ji, P. Li, X. Zhang and L. Wang, *Org. Biomol. Chem.*, 2013, **11**, 4095-4101. (c) C. Martinez, R. Alvarez and J. M. Aurrecoechea, *Org. Lett.*, 2009, **11**, 1083-1086. (d) Z. Liang, W. Hou, Y. Du, Y. Zhang, Y. Pan, D. Mao and K. Zhao, *Org. Lett.*, 2009, **11**, 4978-4981. (e) X. Guo, R. Yu, H. Li and Z. Li, *J. Am. Chem. Soc.*, 2009, **131**, 17387-17393. (f) B. Lu, B. Wang, Y. Zhang and D. Ma, *J. Org. Chem.*, 2007, **72**, 5337-5341.

- 8 (a) X. Q. Hao, L. J. Chen, B. Ren, L. Y. Li, X. Y. Yang, J. F. Gong, J. L. Niu and M. P. Song, *Org. Lett.*, 2014, **16**, 1104-1107. (b) D. Liang, M. Wang, Y. Dong, Y. Guo and Q. Liu, *RSC Adv.*, 2014, **4**, 6564-6567. (c) X. F. Cheng, Y. Li, Y. M. Su, F. Yin, J. Y. Wang, J. Sheng, H. U. Vora, X. S. Wang and J. Q. Yu, *J. Am. Chem. Soc.*, 2013, **135**, 1236-1239.
- 9 (a) M. Ciufolini, N. Braun, S. Canesi, M. Ousmer, J. Chang and D. Chai, *Synthesis*, 2007, **2007**, 3759-3772. (b) S. Quideau, L. Pouysegu and D. Deffieux, *Curr. Org. Chem.*, 2004, **8**, 113-148. (c) P. Wipf and S. Rodríguez, *Synthesis*, 2004, 2767-2783.
- 10 11 H. Kim, J. Ralph, N. Yahiaoui, M. Pean and A. M. Boudet, *Org. Lett.*, 2000, **2**, 2197-2200.
- 12 A. Lattanzi, A. Senatore, A. Massa and A. Scettri, *J. Org. Chem.*, 2003, **68**, 3691-3694.
- 15 13 X. F. Duan, J. Zeng, Z. B. Zhang and G. F. Zi, *J. Org. Chem.*, 2007, **72**, 10283-10286.
- 14 B. Xiao, T. J. Gong, Z. J. Liu, J. H. Liu, D. F. Luo, J. Xu and L. Liu, *J. Am. Chem. Soc.*, 2011, **133**, 9250-9253.
- 15 Y. Wei and N. Yoshikai, *Org. Lett.*, 2011, **13**, 5504-5507.
- 20 16 (a) J. Zhao, Y. Wang, Y. He, L. Liu and Q. Zhu, *Org. Lett.*, 2012, **14**, 1078-1081. (b) J. Zhao, Q. Zhang, L. Liu, Y. He, J. Li, J. Li and Q. Zhu, *Org. Lett.*, 2012, **14**, 5362-5365.
- 25 17 (a) M. Chennapuram, N. R. Emmadi, C. Bingi and K. Atmakur, *RSC Adv.*, 2015, **5**, 19418-19425. (b) K. K. D. R. Viswanadham, M. Prathap Reddy, P. Sathyanarayana, O. Ravi, R. Kant and S. R. Bathula, *Chem. Commun.*, 2014, **50**, 13517-13520. (c) B. A. Dar, N. A. Dangroo, A. Gupta, A. Wali, M. A. Khuroo, R. A. Vishwakarma and B. Singh, *Tetrahedron Lett.*, 2014, **55**, 1544-1548. (d) N. M. Mishra, D. D. Vachhani, S. G. Modha and E. V. Van der Eycken, *Eur. J. Org. Chem.*, 2013, **2013**, 693-700. (e) J. Dhineshkumar, M. Lamani, K. Alagiri and K. R. Prabhu, *Org. Lett.*, 2013, **15**, 1092-1095.
- 30