



# Chemical Science

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. Eisen and R. J. Batrice, *Chem. Sci.*, 2015, DOI: 10.1039/C5SC02746B.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

*Accepted Manuscripts* are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



## Chemical Science

## EDGE ARTICLE

## Catalytic insertion of E-H bonds (E=C, N, P, S) into heterocumulenes by amido-actinide complexes†

Received 00th January 20xx,  
Accepted 00th January 20xx

Rami J. Batrice and Moris S. Eisen\*

DOI: 10.1039/x0xx00000x

www.rsc.org/

We report herein the actinide-mediated insertion of E-H bonds (E=C, N, P, S) into various heterocumulenes including carbodiimides, isocyanates, and isothiocyanates. The precatalysts are prepared by a simple, one-pot procedure using readily available starting materials, and challenging insertions are achieved with excellent selectivity in short reaction times. Spectroscopic data are utilised to propose the active catalytic mechanism and derive thermodynamic activation parameters, which are described in this study. The only example of an actinide-mediated C-H bond insertion into a carbon-heteroatom bond is presented within this study.

## Introduction

The structural and catalytic behaviour of the actinides has attracted considerable attention in recent years owing to their unique coordination geometries,<sup>1–10</sup> as well as their impressive reactivity profiles in numerous stoichiometric and catalytic processes. Hydroaminations,<sup>11–18</sup> hydroalkoxylations,<sup>19</sup> hydrosilylations,<sup>20–22</sup> polymerisations,<sup>23–29</sup> alkyne oligomerisations,<sup>30–32</sup> and numerous small molecule activations<sup>28, 33–40</sup> represent a few of the advancements made in this field. However, the insertion of heterocumulenes into E-H bonds (E=N, P, S) has only recently appeared in the scope of accessible reactions in *f*-block chemistry.<sup>41–47</sup> The guanidine, phosphaguanidine, and thiourea products obtained have found considerable utility as ligands in coordination compounds,<sup>48</sup> medicinal applications,<sup>49</sup> and synthons for challenging organic transformations.<sup>50</sup>

Carbodiimides in particular have been shown to insert stoichiometrically into the actinide-carbon bond of Cp\*<sub>2</sub>AnMe<sub>2</sub> (An=U, Th), giving rise to the corresponding pentamethylcyclopentadienyl-amidinate complexes of the early actinides,<sup>51</sup> however only in this year was the process made catalytic, mediated by a mono(imidazolin-2-iminato) thorium(IV) complex and often accompanied by extended reaction times and moderate conversions.<sup>52</sup> Moreover, the

low-yielding multi-step ligand synthesis provides a further challenge accompanying the production and application of this complex. Recently, we have shown that a series of simple amido-actinide complexes of the formula U[N(SiMe<sub>3</sub>)<sub>2</sub>]<sub>3</sub> (**1**) and [(Me<sub>3</sub>Si)<sub>2</sub>N]<sub>2</sub>An[κ<sup>2</sup>-(N,C)-CH<sub>2</sub>Si(CH<sub>3</sub>)<sub>2</sub>N(SiMe<sub>3</sub>)] (An=U (**2**), Th (**3**)) (Fig. 1) are capable of effecting the efficient oligomerisation or cyclotrimerisation of terminal alkynes,<sup>53</sup> the sole example of a catalytic transformation employing this class of complexes prior to the work described in this study. These complexes are generated in a one-pot process using readily available starting materials and have already shown impressive stoichiometric reactivity,<sup>54, 55</sup> making them exceptionally attractive in their use as catalysts. In our attempt to uncover new reactivity with this class of complexes, we have applied them toward the catalytic insertion of E-H bonds into heterocumulenes (Y=C=X), wherein E= C, N, P, or S, and Y=RN, and X=RN, O, or S. Most interestingly was the catalytic insertion of C-H bonds into carbodiimides, the only example of actinide-mediated C-H insertion into a C-X unsaturated bond to date.

Previously, the effects of opening the coordination sphere of an actinide catalyst was achieved in cyclopentadienyl-actinide systems by the use of an *ansa*-bridged tetramethylcyclopentadienyl ligand, resulting in markedly increased activity and selectivity.<sup>56</sup> In the catalytic study of the amido-actinide complexes (**1–3**), the coordinative unsaturation is considerably increased as compared to the analogous actinide(imidazolin-2-iminato) complex, similarly providing superior reactivity over the previously reported system. These findings open the scope of possible catalytic transformations toward new, challenging reactions.




Fig. 1 Precatalysts employed for the catalytic insertion of E-H bonds into heterocumulenes.


Schulich Faculty of Chemistry, Technion – Israel Institute of Technology, Technion City, 32000, Haifa, Israel. E-mail: chmoris@tx.technion.ac.il


† Electronic Supplementary Information (ESI) available: Detailed experimental conditions, product characterisation, kinetic, and thermodynamic data are provided. See DOI: 10.1039/x0xx00000x



## Results and Discussion

The catalytic reactions of *N,N'*-diisopropylcarbodiimide (DIC), di-*p*-tolylcarbodiimide (DTC), phenylisocyanate, and phenylisothiocyanate with a series REH moieties are presented below. To establish the potential reactivity of these complexes, the insertion of aniline into diisopropylcarbodiimide was monitored using  $^1\text{H}$  NMR spectroscopy (Fig. 2), showing the reaction progress as a function of DIC consumption and guanidine formation, (Fig. 2, (a) and (b), respectively). Further investigation of the mode of activation of the actinide centre was performed by independent addition of aniline and DIC. In each of the uranium and thorium complexes, it is seen that complete displacement of the hexamethyldisilazane occurs after the addition of aniline, and that conservation of the precatalyst oxidation state is maintained throughout the reaction. Conversely, the reaction with DIC necessitated elevated temperatures to activate the metal centre. These findings inform us that activation of the precatalyst is achieved by the amine moiety to generate the catalytically active species. Attempts to isolate the intermediates generated and characterisation by crystallisation were unsuccessful, however previous studies have shown that activation of the thorium complex **3** with terminal alkyne similarly results in complete displacement of the amido moieties.<sup>53</sup> It is worth note that the order of addition of the reagents to the precatalyst solution did not markedly affect the reaction rate or product distribution. The nature of the REH substrate additionally affects the outcome of the insertion product (Fig. 3); where E is a monoprotic nucleophile, product **4** is obtained as the sole insertion product. However, when E is a diprotic nucleophile (such as a primary amine), isomerisation of compound **4** is possible, yielding product **5** as the major product. The isomerisation with diprotic nucleophiles is dependent on the resulting guanidine generated, wherein the formation of the C=N bond is directed toward the more electron withdrawing arm of the structure. The insertion of a series of REH moieties into diisopropylcarbodiimide, mediated by the three actinide complexes presented (**1-3**), was investigated and the results summarised in Table 1. All experiments presented herein were



**Table 1** Catalytic reactions of diisopropylcarbodiimide with E-H species mediated by complexes **1-3**<sup>a</sup>


Entry	Precatalyst	R = (R <sub>2</sub> EH/REH)	Reaction Time (h)	Conv <sup>b</sup> (%)	Product
1	<b>1</b>		6	>99	
2	<b>2</b>	PhNH <sub>2</sub>	9	>99	<b>5ae</b>
3	<b>3</b>		4	94	
4	<b>1</b>		3	>99	
5	<b>2</b>	<i>o</i> MeO-PhNH <sub>2</sub>	12	93	<b>5af</b>
6	<b>3</b>		24	96	
7	<b>1</b>		6	>99	
8	<b>2</b>	<i>o</i> Me-PhNH <sub>2</sub>	12	93	<b>5ag</b>
9	<b>3</b>		6	95	
10	<b>1</b>		9	>99	
11	<b>2</b>	<i>p</i> Cl-PhNH <sub>2</sub>	12	>99	<b>5ah</b>
12	<b>3</b>		4	>99	
13	<b>1</b>	<i>i</i> PrNH <sub>2</sub>	24	5	<b>4ai</b>
14	<b>3</b>		24	15	
15	<b>1</b>	Et <sub>2</sub> NH	24	23	<b>4aj</b>
16	<b>3</b>		24	31	
17	<b>1</b>		24	58	
18	<b>2</b>	Ph <sub>2</sub> PH	24	53	<b>4ak</b>
19	<b>3</b>		2	>99	
20	<b>3</b>	BnSH	2	>99	<b>6al</b>
21	<b>1</b>	<i>t</i> BuCCH	24	51	<b>6am</b>
22	<b>2</b>		24	81	
23	<b>1</b>		9	96	
24	<b>2</b>	Me <sub>3</sub> SiCCH	24	96	<b>6an</b>
25	<b>3</b>		12	>99	
26	<b>1</b>		9	97	
27	<b>2</b>	PhCCH	24	97	<b>6ao</b>
28	<b>3</b>		18	>99	


<sup>a</sup>Reaction conditions: ~3.5 μmol precatalyst (1 mol%), 600 μL C<sub>6</sub>D<sub>6</sub>, 75 °C.<sup>b</sup>Determined by <sup>1</sup>H NMR of the crude reaction mixture.

whereas the thorium complex (**3**) (lacking 5f electrons) is unable to perform this stabilising interaction and allows for rapid protonolysis and insertion. The insertion of benzylthiol was attempted with each of the uranium precatalysts, however upon addition of the thiol, immediate formation of a green precipitate was observed, which reacted no further after heating. The thorium complex **3**, conversely, reacted within two hours to near-quantitative conversion toward product **6al**


(Table 1, entry 20). It is interesting to note the lower activity of the thorium complex (**3**) as compared to the uranium complexes employed in the reaction with *ortho*-anisidine (Table 1, entries 4-6). It is proposed that the highly electrophilic thorium catalyst forms a dative coordination to the *o*-methoxy functionality of the aniline, thus inhibiting the catalytic turnover.

Encouraged by the diversity of E-H moieties demonstrating favourable reactivity in the actinide-mediated insertion, attention was turned to the possibility of insertion of a C-H bond, a process hitherto unseen in actinide chemistry for heteroatom-containing multiple bonds. Performing this reaction with trimethylsilyl- or phenyl-acetylene generated the desired insertion product with all complexes in excellent yield and short reaction times (Table 1, entries 23-28), providing the substituted amidine. The insertion of 3,3-dimethylbutyne was observed in 51 and 81% conversion after 24 hours after heating in the presence of complexes **1** and **2**, respectively (Table 1, entries 21 and 22), generating *N,N'*-diisopropyl-4,4-dimethylpent-2-ynimidamide. The analogous reaction performed with complex **3** produced alkyne dimerization product, however no detectable insertion into the carbodiimide was observed. This finding illustrates the competitive process between the alkyne oligomerisation and insertion into the heterocumulene, revealing a lower barrier energy for insertion into the C≡C bond, favouring the former process (Fig. 4). These reactions were additionally attempted using 1-hexyne, however oligomerisation forming organic enynes was dominant, generating a mixture of products from competitive insertions.

Similar reactions were performed using di-*p*-tolylcarbodiimide (DTC) as the heterocumulene source, and the results are presented in Table 2. Initial analysis of the reactions performed with DTC revealed superb substrate conversions in moderate reaction times. The reaction of aromatic amines with DTC generated the corresponding guanidines in excellent yield after heating at 75 °C (Table 2, entries 1-12). The most apparent trend of these reactions is the shorter reaction times when utilising the thorium metallacycle (**3**), and the observed acceleration of the reaction rate as a more electron deficient aniline is used (Table 2, entries 3, 6, 9, and 12). This trend in reactivity is an expected corollary of the decreased electron donation from the aniline derivatives to the metal centre, resulting in a weaker, more labile bond, and allowing for a more rapid insertion of the carbodiimide into the Th-N bond. Similar short reaction times are observed for the insertion of Ph<sub>2</sub>PH using the thorium

**Fig. 4** Competitive alkyne and carbodiimide insertion into actinide-acetylide bond.

## Edge Article

**Table 2** Catalytic reactions of di-*p*-tolylcarbodiimide with E-H species mediated by complexes **1–3**<sup>a</sup>

Entry	Precatalyst	R= (R <sub>2</sub> EH/REH)	Reaction Time (h)	Conv <sup>b</sup> (%)	Product
1	<b>1</b>		24	89	
2	<b>2</b>	PhNH <sub>2</sub>	24	93	<b>5be</b>
3	<b>3</b>		6	>99	
4	<b>1</b>		24	91	
5	<b>2</b>	<i>o</i> MeO-PhNH <sub>2</sub>	24	>99	<b>4bf</b>
6	<b>3</b>		24	90	
7	<b>1</b>		24	96	
8	<b>2</b>	<i>o</i> Me-PhNH <sub>2</sub>	24	>99	<b>4bg</b>
9	<b>3</b>		9	>99	
10	<b>1</b>		24	98	
11	<b>2</b>	<i>p</i> Cl-PhNH <sub>2</sub>	24	94	<b>5bh</b>
12	<b>3</b>		6	>99	
13	<b>1</b>		12	98	
14	<b>2</b>	Ph <sub>2</sub> PH	24	85	<b>4bk</b>
15	<b>3</b>		24	83	
16	<b>1</b>	BnSH	6	>99	<b>6bl</b>
17	<b>3</b>		3	>99	

<sup>a</sup>Reaction conditions: ~3.5 μmol precatalyst (1 mol%), 600 μL C<sub>6</sub>D<sub>6</sub>, 75 °C.<sup>b</sup>Determined by <sup>1</sup>H NMR of the crude reaction mixture.

precatalyst (**3**) as compared to the uranium precatalysts (**1** and **2**) (Table 2, entries 13–15).

Performing these reactions with benzylthiol resulted in rapid insertion into DTC for complexes **1** and **3** (Table 2, entries 16 and 17), however the reaction with precatalyst **2** again resulted in the formation of an intractable green precipitate, which formed immediately upon addition of the thiol. This is most likely due to the formation of insoluble U(IV) dimers bridged by the thiol moiety, which has been previously shown to occur, especially for U(IV) metallocene systems.<sup>57</sup> Similar reactions were performed with aliphatic amines and terminal alkynes, however no catalytic turnover was observed in any of these reactions, showing that the insertion of the electron deficient carbodiimide is unable to surpass the bonding energy of the actinide-nitrogen and actinide-carbon bonds described.

The scope of these reactions illustrating the robust nature of these complexes was expanded further yet in the insertion into isocyanates and isothiocyanates. Using phenylisocyanate and phenylisothiocyanate as the heterocumulene source, the insertion of diphenylphosphine, benzylthiol, and terminal

alkynes was studied, and the results presented in Table 3. It is most notable that the catalytic insertion into these substrates was only achieved using the uranium complexes; this observation is justified by the greater bond-disruption enthalpy of the Th-O and Th-S bond as compared to the uranium analogues (Th-O = 208.0, U-O = 181.0, Th-S = 145.2, and U-S = 121.9 kcal mol<sup>-1</sup>).<sup>58</sup> These bonding energies indicate a higher barrier of insertion into the Th-S or Th-O species, which is not exceeded under the given reaction conditions, additionally supporting the observed reactivity of the thorium complex with *ortho*-anisidine. The insertion of diphenylphosphine gives only moderate conversions using either uranium complex with both heterocumulenes (Table 3, entries 1, 2, 5, and 6), and a similar trend is observed using benzylthiol, however the reaction of this E-H moiety with phenylisocyanate mediated by the uranium(III) complex (**1**) shows an enhanced reactivity as compared to the uranium(IV) complex (**2**), providing a 99% conversion to product **8cl** after 24 hours at 75 °C (Table 3, entry 3).

Analysis of the rate order reveals first order kinetics in precatalyst, heterocumulene, and REH, giving rise to the rate equation 1:

$$\frac{\partial p}{\partial t} = k_{obs} \cdot [catalyst]^1 \cdot [DIC]^1 \cdot [aniline]^1 \quad (1);$$

(a more thorough presentation of the kinetic rate law is presented within the Supporting Information). The mechanism of activation was studied using DIC and aniline. The first step is a rapid protolytic cleavage by the aniline which displaces all hexamethyldisilazide moieties at room temperature (Fig. 5, step (a)), followed by rapid reversible migratory insertion of

**Table 3** Catalytic reactions of phenylisocyanate or phenylisothiocyanate with E-H species mediated by complexes **1** and **2**<sup>a</sup>

Entry	Precatalyst	Ph-N=C=X	R= (R <sub>2</sub> EH/REH)	Conv <sup>b</sup> (%)	Product
1	<b>1</b>			52	
2	<b>2</b>		Ph <sub>2</sub> PH	64	<b>7ck</b>
3	<b>1</b>	O	BnSH	99	<b>8cl</b>
4	<b>2</b>			68	
5	<b>1</b>		Ph <sub>2</sub> PH	49	<b>7ck</b>
6	<b>2</b>			55	
7	<b>1</b>	S	BnSH	47	<b>8dk</b>
8	<b>2</b>			52	

<sup>a</sup>Reaction conditions: ~3.5 μmol precatalyst (1 mol%), 600 μL C<sub>6</sub>D<sub>6</sub>, 75 °C, 24 h.<sup>b</sup>Determined by <sup>1</sup>H NMR of the crude reaction mixture.






Fig. 5 Plausible mechanism of the actinide-mediated catalytic insertion of E-H bond into carbodiimide.

DIC into the An-N bond (step (b)). A slow protolytic cleavage by an additional equivalent of amine (step (c)) is operative (turnover limiting step) to complete the catalytic cycle with concomitant liberation of the organic product and regeneration of the active catalyst. Deuterium labelling studies were performed in this reaction using aniline-*N,N*-*d*<sub>2</sub>, revealing a primary kinetic isotope effect ( $k_H/k_D = 1.22$ ), further supporting the protonolysis of the guanidinate is the turnover limiting step of the catalytic cycle (Figure 5, step (c)).

Thermodynamic parameters were experimentally calculated from <sup>1</sup>H NMR data of aniline insertion into DIC using Eyring plots (see Supporting Information), and the enthalpy of activation ( $\Delta H^\ddagger$ ) for complexes **1-3** were found to be 9.9(1), 10.5(1), and 9.7(1) kcal mol<sup>-1</sup>, respectively, showing a low energetic barrier to the actinide-mediated insertion. The proposed  $\sigma$ -bond metathesis by aniline via a four-centred transition state is additionally supported by the remarkably low entropies of activation ( $\Delta S^\ddagger$ ) of -49.5(3), -48.0(4), and -52.8(3) e.u., respectively, for complexes **1-3**. Calculating the Gibbs energy of activation ( $\Delta G^\ddagger$ ) for these reactions at 75 °C showed extremely similar values for each of the precatalysts employed (27.1(1), 27.2(1), and 28.1(1) kcal mol<sup>-1</sup> for **1-3**, respectively).

## Conclusions

This work presents a series of simple coordinative unsaturated amido-actinide complexes capable of performing the catalytic insertion of E-H bonds into heterocumulenes, where E=C, N, P, or S. The insertion into carbodiimides showed excellent reaction rates and functional group tolerances. The insertion of terminal alkynes represents a new facet of actinide-mediated catalysis. The insertion into phenylisocyanate and phenylisothiocyanate showed a more limited reactivity, however the reaction performed with phosphines and thiols yielded the desired phosphanecarboxamide and carbamothioate.

Deuterium-labelling studies in the reaction of PhND<sub>2</sub> with DIC revealed a KIE of 1.22, indicating protolytic cleavage of the guanidinate to be the rate-determining step of the catalytic cycle. Preliminary studies in our group show the possibility of using group 4 metal complexes to effect similar chemical transformations. The investigation of further catalytic transformation using these actinide precatalysts is currently underway.

## Notes and references

- P. L. Arnold, A. F. Pecharman and J. B. Love, *Angew Chem Int Ed Engl*, 2011, **50**, 9456-9458.
- H. S. La Pierre, A. Scheurer, F. W. Heinemann, W. Hieringer and K. Meyer, *Angew Chem Int Ed Engl*, 2014, **53**, 7158-7162.
- P. L. Arnold, C. J. Stevens, J. H. Farnaby, M. G. Gardiner, G. S. Nichol and J. B. Love, *J. Am. Chem. Soc.*, 2014, **136**, 10218-10221.
- P. L. Diaconescu and C. C. Cummins, *Inorg. Chem.*, 2012, **51**, 2902-2916.
- B. Vidjayacoumar, S. Ilango, M. J. Ray, T. Chu, K. B. Kolpin, N. R. Andreychuk, C. A. Cruz, D. J. H. Emslie, H. A. Jenkins and J. F. Britten, *Dalton Trans.*, 2012, **41**, 8175-8189.
- R. E. Jilek, L. P. Spencer, D. L. Kuiper, B. L. Scott, U. J. Williams, J. M. Kikkawa, E. J. Schelter and J. M. Boncella, *Inorg. Chem.*, 2011, **50**, 4235-4237.
- M. J. Monreal, S. I. Khan, J. L. Kiplinger and P. L. Diaconescu, *Chem. Commun.*, 2011, **47**, 9119-9121.
- O. J. Cooper, D. P. Mills, J. McMaster, F. Tuna, E. J. L. McInnes, W. Lewis, A. J. Blake and S. T. Liddle, *Chem. Eur. J.*, 2013, **19**, 7071-7083.
- D. Patel, F. Tuna, E. J. L. McInnes, J. McMaster, W. Lewis, A. J. Blake and S. T. Liddle, *Dalton Trans.*, 2013, **42**, 5224-5227.
- W. J. Evans, N. A. Siladke and J. W. Ziller, *Chem. Eur. J.*, 2010, **16**, 796-800.
- W. Ren, G. Zi, D.-C. Fang and M. D. Walter, *Chem. Eur. J.*, 2011, **17**, 12669-12682.
- B. D. Stubbart and T. J. Marks, *J. Am. Chem. Soc.*, 2007, **129**, 6149-6167.
- B. D. Stubbart and T. J. Marks, *J. Am. Chem. Soc.*, 2007, **129**, 4253-4271.
- M. S. Eisen, T. Straub and A. Haskel, *J. Alloy Compd.*, 1998, **271-273**, 116-122.
- A. Haskel, T. Straub and M. S. Eisen, *Organometallics*, 1996, **15**, 3773-3775.
- C. E. Hayes, R. H. Platel, L. L. Schafer and D. B. Leznoff, *Organometallics*, 2012, **31**, 6732-6740.
- T. Straub, A. Haskel, T. G. Neyroud, M. Kapon, M. Botoshansky and M. S. Eisen, *Organometallics*, 2001, **20**, 5017-5035.
- B. D. Stubbart, C. L. Stern and T. J. Marks, *Organometallics*, 2003, **22**, 4836-4838.
- S. D. Wobser and T. J. Marks, *Organometallics*, 2013, **32**, 2517-2528.
- A. K. Dash, J. Q. Wang and M. S. Eisen, *Organometallics*, 1999, **18**, 4724-4741.
- A. K. Dash, J. Q. Wang, J. Wang, I. Gourevich and M. S. Eisen, *J. Nucl. Sci. Technol.*, 2002, **39**, 386-392.



## Edge Article

## Chemical Science

22. A. K. Dash, J. X. Wang, J. C. Berthet, M. Ephritikhine and M. S. Eisen, *J. Organomet. Chem.*, 2000, **604**, 83-98.
23. E. Barnea, D. Moradove, J.-C. Berthet, M. Ephritikhine and M. S. Eisen, *Organometallics*, 2006, **25**, 320-322.
24. J. Fang, A. Walshe, L. Maron and R. J. Baker, *Inorg. Chem.*, 2012, **51**, 9132-9140.
25. C. E. Hayes, Y. Sarazin, M. J. Katz, J.-F. Carpentier and D. B. Leznoff, *Organometallics*, 2013, **32**, 1183-1192.
26. I. S. R. Karmel, M. Botoshansky, M. Tamm and M. S. Eisen, *Inorg. Chem.*, 2014, **53**, 694-696.
27. I. S. R. Karmel, T. Elkin, N. Fridman and M. S. Eisen, *Dalton Trans.*, 2014, **43**, 11376-11387.
28. S. M. Mansell, J. H. Farnaby, A. I. Germeroth and P. L. Arnold, *Organometallics*, 2013, **32**, 4214-4222.
29. A. Walshe, J. Fang, L. Maron and R. J. Baker, *Inorg. Chem.*, 2013, **52**, 9077-9086.
30. A. Haskel, T. Straub, A. K. Dash and M. S. Eisen, *J. Am. Chem. Soc.*, 1999, **121**, 3014-3024.
31. T. Straub, A. Haskel and M. S. Eisen, *J. Am. Chem. Soc.*, 1995, **117**, 6364-6365.
32. J. Q. Wang, A. K. Dash, J. C. Berthet, M. Ephritikhine and M. S. Eisen, *Organometallics*, 1999, **18**, 2407-2409.
33. P. L. Arnold, S. M. Mansell, L. Maron and D. McKay, *Nat. Chem.*, 2012, **4**, 668-674.
34. P. L. Arnold, Z. R. Turner, R. M. Bellabarba and R. P. Tooze, *Chem. Sci.*, 2011, **2**, 77-79.
35. O. P. Lam, S. M. Franke, F. W. Heinemann and K. Meyer, *J. Am. Chem. Soc.*, 2012, **134**, 16877-16881.
36. V. Mougél, C. Camp, J. Pécaut, C. Copéret, L. Maron, C. E. Kefalidis and M. Mazzanti, *Angew. Chem. Int. Ed.*, 2012, **51**, 12280-12284.
37. A.-C. Schmidt, A. V. Nizovtsev, A. Scheurer, F. W. Heinemann and K. Meyer, *Chem. Commun.*, 2012, **48**, 8634-8636.
38. C. L. Webster, J. W. Ziller and W. J. Evans, *Organometallics*, 2012, **31**, 7191-7197.
39. E. Zhou, W. Ren, G. Hou, G. Zi, D.-C. Fang and M. D. Walter, *Organometallics*, 2015, DOI: 10.1021/acs.organomet.5b00454.
40. A. R. Fox, S. C. Bart, K. Meyer and C. C. Cummins, *Nature*, 2008, **455**, 341-349.
41. I. J. Casely, J. W. Ziller and W. J. Evans, *Organometallics*, 2011, **30**, 4873-4881.
42. A. C. Behrle and J. A. R. Schmidt, *Organometallics*, 2013, **32**, 1141-1149.
43. W. Yi, J. Zhang, L. Hong, Z. Chen and X. Zhou, *Organometallics*, 2011, **30**, 5809-5814.
44. J. Tu, W. Li, M. Xue, Y. Zhang and Q. Shen, *Dalton Trans.*, 2013, **42**, 5890-5901.
45. Z. Li, M. Xue, H. Yao, H. Sun, Y. Zhang and Q. Shen, *J. Organomet. Chem.*, 2012, **713**, 27-34.
46. Y. Cao, Z. Du, W. Li, J. Li, Y. Zhang, F. Xu and Q. Shen, *Inorg. Chem.*, 2011, **50**, 3729-3737.
47. Z. Du, H. Zhou, H. Yao, Y. Zhang, Y. Yao and Q. Shen, *Chem. Commun.*, 2011, **47**, 3595-3597.
48. W.-X. Zhang, L. Xu and Z. Xi, *Chem. Commun.*, 2015, **51**, 254-265.
49. S. G. Zarate, A. G. Santana, A. Bastida and J. Revuelta, *Curr. Org. Chem.*, 2014, **18**, 2711-2749.
50. T. Ishikawa and T. Kumamoto, *Synthesis*, 2006, **2006**, 737-752.
51. W. J. Evans, J. R. Walensky, J. W. Ziller and A. L. Rheingold, *Organometallics*, 2009, **28**, 3350-3357.
52. I. S. R. Karmel, M. Tamm and M. S. Eisen, *Angew. Chem. Int. Ed.*, 2015, DOI: 10.1002/anie.201502041.
53. R. J. Batrice, J. McKinven, P. L. Arnold and M. S. Eisen, *Organometallics*, 2015, **34**, 4039-4050.
54. S. J. Simpson and R. A. Andersen, *J. Am. Chem. Soc.*, 1981, **103**, 4063-4066.
55. S. J. Simpson, H. W. Turner and R. A. Andersen, *J. Am. Chem. Soc.*, 1979, **101**, 7728-7729.
56. A. K. Dash, I. Gourevich, J. Q. Wang, J. Wang, M. Kapon and M. S. Eisen, *Organometallics*, 2001, **20**, 5084-5104.
57. S. D. Stults, R. A. Andersen and A. Zalkin, *Organometallics*, 1990, **9**, 1623-1629.
58. *CRC Handbook of Chemistry and Physics*, CRC Press, Boca Raton, FL, 96th ed. edn., 2015-2016.

