


# Tandem Reduction / Intramolecular Hydroxyalkylation of (3-Hydroxyphenyl)alkanoates: a New Regioselective Approach to 1,8-Dihydroxytetralins

Giuseppe Guanti,\* Luca Banfi, Enrica Narisano, Renata Riva, and Sergio Thea

Istituto di Chimica Organica e C.N.R., C.d.S. sui Diariloidi, corso Europa 26, I-16132 GENOVA (Italy)


**Abstract:** 4-(3-hydroxyphenyl)-butanoates **4**, on treatment with DIBALH, followed by hydrolytic work-up, undergo a novel completely regioselective intramolecular hydroxyalkylation reaction to give 1,8-dihydroxytetralins. The yield of the reaction depends heavily on the structure of starting material, best results being achieved with 3,3-disubstituted butanoates.

The discovery that 11-deoxyanthracyclinones **1** are present as aglycone portions in some antibiotics possessing significant anti-cancer activity and reduced cardiotoxicity has attracted the interest of several research groups, stimulating extensive synthetic work in this field.<sup>1</sup> A possible strategy for the preparation of these linearly fused units involves first the assemblage of the AB ring system followed by its coupling with the remaining CD rings to complete the tetracyclic arrangement. Various different disconnections can be envisaged for the AB fragment, and some elegant syntheses had already been published.<sup>2</sup> In the course of a research program directed to the development of new synthetic routes to this class of antibiotics, we planned to prepare the 1,8-dihydroxytetralin fragment **2** through a completely new approach, based on an intramolecular C-C bond formation between the positions 1 and 8a, starting from a monocyclic precursor **3**.



In order to examine the feasibility of this project, we decided to synthesize some suitably substituted 4-(3-hydroxyphenyl)butanals, and to study in details the factors controlling the *ortho*-*para* ratio in the intramolecular hydroxyalkylation. For this purpose the model "meta" ester **4a**<sup>3</sup> was first prepared, planning to accomplish its conversion into the corresponding phenolic aldehyde **7** either *via* a protection-reduction-deprotection sequence, or through direct reduction (Scheme 1).<sup>4</sup> Thus **4a** was first protected as the dimethyl-*t*-butylsilyl ether and then converted into silylated aldehyde **5** either through direct chemoselective reduction with DIBALH, or *via* a two-step sequence (reduction to the primary alcohol followed by Swern oxidation). Further treatment with ethanolic NaOH removed the protecting group<sup>5</sup> and the resulting phenolic aldehyde **7** immediately cyclized, under these basic conditions, to give in good yield exclusively the 1,6-dihydroxytetralin **6**, deriving from attack of the carbonyl group at the *para* position of the aromatic ring. On the other hand, when the reduction with DIBALH was carried out directly on the unprotected ester **4a**, the main product turned out to be, instead than

Scheme 1



a)  $\text{Me}_2\text{t-BuSiCl}$ , imidazole, DMF, r.t.; b) DIBALH,  $\text{CH}_2\text{Cl}_2$ ,  $-78^\circ\text{C}$ ; c) 1.  $\text{LiBH}_4$ ,  $\text{Et}_2\text{O}$ ; 2.  $(\text{COCl})_2$ , DMSO,  $\text{Et}_3\text{N}$ ,  $-78^\circ \rightarrow -40^\circ\text{C}$ ; d)  $\text{NaOH}$ ,  $\text{MeOH}$ , r.t.; e) see typical procedure in the Table.

the expected aldehyde 7, an alcohol whose structure, determined by  $^1\text{H}$  and  $^{13}\text{C}$  NMR, corresponded to the 1,8-dihydroxytetralin 8a, as confirmed by its conversion into iso-propylidene derivative 11a (see Table). This result was particularly interesting, since one-pot transformation of the starting phenolic ester 4a into 8a, through a tandem reduction-cyclization reaction, was realized. Moreover the reaction proceeded with complete *ortho* regioselectivity (no *para* compound 6 was detected) and in satisfactory yield (49%), which was further improved by *in situ* derivatization of 8a to give the diacetate 10a (75% yield from non-recovered 4a).<sup>7</sup> This outcome demonstrated the possibility of performing C-C bond formation between position 1 and 8a, thus opening a new synthetic route to fragment 2a.

In order to define the scope and limitations of this unprecedented reaction, we synthesized a series 4-(3-hydroxyphenyl)butanoates 4b-f, variously substituted at position 3 of the aliphatic chain,<sup>3</sup> as well as ethyl 3-(3-hydroxyphenyl)propanoate 12 and 5-(3-hydroxyphenyl)pentanoate 13,<sup>3</sup> and subjected them to the same reaction conditions already employed for 4a. To our surprise we noted big differences in their behaviour. Actually the yield and the product distribution appeared to depend dramatically upon the length of carbon chain as well as on the presence of substituents (at the moment we examined only the effect of modifications at position 3). First of all we observed that compounds 12 and 13 did not furnish at all respectively the five and seven-membered cyclization compounds, leading instead in good yield to the corresponding aldehydes deriving from simple reduction. On the other hand, compounds 4c and 4d, which bear no substituents at position 3, afforded 8c and 8d only in low yields, due to the formation of several by-products.<sup>7</sup> It should be noted that in this case, like for 4a, very little or no aldehyde was detected. On the contrary, good yields were obtained starting from dithiane 4b and 3,3-dimethyl derivative 4f, which are both disubstituted at position 3, while the behaviour of monosubstituted 4e falls in between. Thus, although further data may be desirable in order to draw generalization, it seems that the presence of a double substitution at position 3 is crucial for the achievement of satisfactory yields.<sup>8</sup>

Finally, about the mechanism and the factors responsible for the observed highly regioselective cyclization, at the moment we are only able to advance a tentative explanation. Addition at  $-78^\circ\text{C}$  of 2 eq. DIBALH to the phenolic esters 4 may generate initially a diisobutylaluminium phenolate and then effects ester reduction to give an O-(diisobutylaluminium) acetal, which does not probably cyclize under these conditions.<sup>9</sup> Aqueous quenching (saturated  $\text{NH}_4\text{Cl}$ ) then frees the aldehyde group, which adds to the aromatic ring under the intramolecular catalysis given by the aluminium phenolate. The intervention of chelated transition state 9 (Scheme 1) can explain the preferential formation of *ortho* cyclization products 8.<sup>10</sup> On the contrary, when the intramolecular hydroxyalky-





Table: Intramolecular Cyclization of Esters 4a-h<sup>a</sup>

a) 1) DIBALH, CH<sub>2</sub>Cl<sub>2</sub>; 2) NH<sub>4</sub>Cl b) Ac<sub>2</sub>O, CH<sub>2</sub>Cl<sub>2</sub>, DMAP, r.t., 2h c) 2-methoxypropene, TsOH, CH<sub>2</sub>Cl<sub>2</sub>, 0°C, 15 min

| Entry | Starting ester | X                                   | Yield of 8 <sup>b</sup> | Yield of 10 <sup>c</sup><br>(from 4) | Yield of 11<br>(from 4) |
|-------|----------------|-------------------------------------|-------------------------|--------------------------------------|-------------------------|
| 1     | 4a             |                                     | 42% (49%)               | 54% (75%)                            | 40%                     |
| 2     | 4b             |                                     | 32%                     | 40% (54%)                            | 40%                     |
| 3     | 4c             | -CH <sub>2</sub> -                  | 10% (13%)               | 10% (18%)                            | 2%                      |
| 4     | 4d             | -O-                                 | 8%                      | not det.                             | not det.                |
| 5     | 4e             | -CH(CH <sub>3</sub> )-              | not det.                | 38% (52%) <sup>d</sup>               | 8%                      |
| 6     | 4f             | -C(CH <sub>3</sub> ) <sub>2</sub> - | 65% (69%)               | 60% (64%)                            | 55%                     |

a) **Typical procedure:** A solution of 4 (1.0 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (10 ml) was cooled to -78°C and treated with DIBALH (1N in CH<sub>2</sub>Cl<sub>2</sub>) (2.1 mmol). After 15 min, the mixture was quenched with saturated aqueous NH<sub>4</sub>Cl (15 ml), warmed to r.t. and stirred for 5h. After dilution with saturated aqueous Na,K tartrate (25 ml), and Et<sub>2</sub>O (30 ml), the mixture was stirred for 30 min. Separation of the phases (in some cases filtration through a celite pad was necessary to remove emulsions) and evaporation afforded crude 8, which was purified through chromatography or *in situ* derivatized to give 10 or 11. b) Yield from non-recovered 4 are in brackets. c) Yield from non-recovered O-acetylated 4 in brackets. d) 10e was obtained as an approximately 1:1 diastereomeric mixture.

lation is carried out under basic conditions and in the absence of a metal atom capable to promote chelation, the *para* position seems to be favoured, most likely for steric reasons, affording 1,6-dihydroxytetralin 6.

Since the dithiolane and dithiane rings are readily removable protecting groups, which can furnish the corresponding carbonyl compounds, the 1,8-dihydroxytetralins 8a and 8b represent versatile intermediates that may find useful applications in the chemistry of 11-deoxyanthracyclinones. Work in this field is under development in our laboratory.

We thank Mr. S. Brusco for his appreciate collaboration, and M.U.R.S.T. and C.N.R. (Progetto Finalizzato Chimica Fine) for financial assistance.

## REFERENCES AND NOTES

- Krohn, K. *Angew. Chem. Int. Ed. Engl.*, **1986**, *25*, 790-807. *Tetrahedron*, **1990**, *46*, 291-318. Thomas, G.J. Synthesis of Anthracyclines related to Daunomycin. In *Recent Progress in the Chemical Synthesis of Antibiotics*, Lukacs, G.; Ohno, M. Eds., Springer, Berlin, **1990**, p. 467-496.
- Li, T.; Wu, Y.L.; Walsgrove, T.C. *Tetrahedron*, **1984**, *40*, 4701-4710. Rizzi, J.P.; Kende, A.S. *Tetrahedron*, **1984**, *40*, 4693-4700. Confalone, P.N.; Pizzolato, G. *J. Am. Chem. Soc.*, **1981**, *103*, 4251-4254. Meyers, A.I.; Higashiyama, K. *J. Org. Chem.*, **1987**, *52*, 4592-4597. Hauser, F.M.; Mal, D. *J. Am. Chem. Soc.*, **1983**, *105*, 5688-5690. Hauser, F.M.; Prasanna, S.; Combs, D.W. *J. Org.*

- Chem.*, 1983, 48, 1328-1333. Hauser, F.M.; Tommasi, R.A. *J. Org. Chem.*, 1991, 56, 5758-5759. Vedejs, E.V.; Pribish, J.R. *J. Org. Chem.*, 1988, 53, 1593-1599. Davis, A.F.; Kumar, A. *Tetrahedron Lett.*, 1991, 32, 7671-7674.
3. Esters **4a-f**, **12** and **13** are all new compounds and their preparation was in some cases not trivial. Synthesis of the esters **4a-c** (starting from commercially available 3-hydroxyphenylacetic acid) : **4a** and **4b**: a) MOMCl, Et<sub>3</sub>N, CH<sub>3</sub>CN, reflux; b) KOH/MeOH 9:1, r.t., 30'; c) 1) CDI, CH<sub>3</sub>CN, 30'; 2) magnesium enolate of EtO<sub>2</sub>C-CH<sub>2</sub>CO<sub>2</sub>H, THF, 60°C, overnight, 54% (3 steps); d) HS(CH<sub>2</sub>)<sub>n</sub>SH, BF<sub>3</sub>·Et<sub>2</sub>O, CH<sub>2</sub>Cl<sub>2</sub>, r.t., 80% (**4a**), 76% (**4b**); **4c**: a) H<sub>2</sub>SO<sub>4</sub>, abs. EtOH, reflux, 24 h; b) BzIbR, K<sub>2</sub>CO<sub>3</sub>, DMF, 88% (2 steps); c) 1 eq DIBALH, CH<sub>2</sub>Cl<sub>2</sub>, -78°C; d) Ph<sub>3</sub>P=CH-CO<sub>2</sub>Et, CH<sub>2</sub>Cl<sub>2</sub>, r.t., powdered 4 Å molecular sieves, 3 h, 84%; e) H<sub>2</sub>, PtO<sub>2</sub>, Pd/C, EtOH, r.t., 24 h, 96%. **4e**: a) see a) and b) in preparation of **4c**; b) KOH, MeOH/H<sub>2</sub>O 9:1, r.t., 6 h, 91%; c) MeONHMe·HCl, WSC, H<sub>2</sub>O/THF 4:1, r.t., 2.5 h, 94%; d) MeLi, THF, -60°C, 1.25 h, 85%; e) Me<sub>3</sub>SiCH<sub>2</sub>CO<sub>2</sub>Et, lithium dicyclohexylamide, THF, -78° → -50°C, 2 h, 87%; f) H<sub>2</sub>, PtO<sub>2</sub>, Pd/C, EtOH, r.t., 5.5 h, 88%. Synthesis of esters **4f**, **12**, and **13** starting from commercially available 3-hydroxybenzaldehyde: **12**: a) malonic acid, pyridine, piperidine, reflux, 8 h, 93%; b) abs. EtOH, H<sub>2</sub>SO<sub>4</sub>, reflux, 7 h, 97%; c) H<sub>2</sub>, PtO<sub>2</sub>, EtOH, r.t., 20 h, 98%. **13**: (starting from **12**): a) BzIbR, K<sub>2</sub>CO<sub>3</sub>, DMF, 80°C, 24 h, 91%; b) 1) 1 eq DIBALH, CH<sub>2</sub>Cl<sub>2</sub>, -78°C, 30'; 2) Ph<sub>3</sub>P=CH-CO<sub>2</sub>Et, CH<sub>2</sub>Cl<sub>2</sub>, powdered 4 Å mol. sieves, 2.5 h, r.t., 61% (2 steps); c) H<sub>2</sub>, PtO<sub>2</sub>, Pd/C, EtOH, r.t., 24 h, 96%. **4f**: a) see a) and b) in preparation of **12**; b) 1) LDA, HMPA, MeI, THF, -78°C, 45'; 2) LDA, HMPA, MeI, -78°C, 15', 47%; c) DIBALH (2 equiv.), CH<sub>2</sub>Cl<sub>2</sub>, -65° → -40°C, 5.5 h, 86% or LiBH<sub>4</sub>, Et<sub>2</sub>O, t. amb., 24 h, 93%; d) (COCl)<sub>2</sub>, DMSO, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, -75° → -30°C; e) 2-lithium-2-trimethylsilyl-1,3-dithiane, THF, 0° → 25°C, 81% (2 steps); f) HgCl<sub>2</sub>, abs. EtOH, r.t., 1 h, 79%; g) BF<sub>3</sub>·Et<sub>2</sub>O, EtSH, t. amb., 2 h, 88%. Synthesis of ester **4e** starting from commercially available 3-hydroxybenzyl alcohol: a) BzIbR, K<sub>2</sub>CO<sub>3</sub>, DMF, 70°C, 6 h, 75%; b) BrCH<sub>2</sub>CO<sub>2</sub>t-Bu, NaOH 50%, benzene, Bu<sub>4</sub>N<sup>+</sup>HSO<sub>4</sub><sup>-</sup>, r.t., overnight, 73%; c) abs. EtOH, H<sub>2</sub>SO<sub>4</sub>, reflux, 2 h; d) H<sub>2</sub>, Pd/C, EtOH, r.t., 1.5 h, 83% (2 steps).
  4. Preliminary attempts to effect the cyclization at the ester stage failed.
  5. Desilylation of **5** with n-Bu<sub>4</sub>NF·3 H<sub>2</sub>O in THF afforded aldehyde **7** in only poor yield due to extensive decomposition. Formation of **6** or **8a** was not observed under these conditions.
  6. The substrate selectivity of DIBALH in the reaction herein studied is never total, and so the cyclization products are always accompanied by variable amounts of unreacted **4** as well as of the primary alcohol deriving from further reduction of the aldehyde.
  7. Compounds **5c**, **5d**, and, although to a lesser extent, **5e** were found to be somewhat unstable under the reaction conditions as well as during purification or derivatization, especially in the presence of acids. See for example the very low yield of iso-propylidene derivatives **7c** and **7e**. On the contrary **4a**, **4b**, and **4f** seem to be definitely more stable. Thence the low yields of entries 3-4 may be due to relatively fast decomposition of **5c** and **5d** in the reaction conditions.
  8. A tentative explanation of this phenomenon will be reported in a forthcoming full paper.
  9. By quenching the reaction mixture with methanolic NaBH<sub>4</sub>, instead of NH<sub>4</sub>Cl, only the primary alcohol deriving from further aldehyde reduction was obtained, together with some unreacted **4**, while no cyclization product **5** was observed. On the other hand, by using more acidic (i.e. AcOH) aqueous quenching conditions, the yield dropped significantly. Finally lower yields were obtained when the aluminium salts were removed soon after quenching with aqueous NH<sub>4</sub>Cl. These results strongly suggest that cyclization occurs only after aqueous quenching.
  10. *Ortho*-directed intermolecular acylations and hydroxyalkylations of phenols, which take advantage of the formation of a six-membered cyclic transition state deriving from covalent bonding of the Lewis acidic catalyst with the phenolic hydroxyl, have been extensively studied in recent years.<sup>11</sup>
  11. Casnati, G.; Casiraghi, G.; Pochini, A.; Sartori, G.; Ungaro, R. *Pure Appl. Chem.*, 1983, 55, 1677; Sartori, G.; Casnati, G.; Bigi, F.; Predieri, G. *J. Org. Chem.*, 1990, 55, 4371-4377 and references therein.