Angewandte

[(NHC)₂Cu]X Complexes as Efficient Catalysts for Azide–Alkyne Click Chemistry at Low Catalyst Loadings**

Silvia Díez-González* and Steven P. Nolan*

Dedicated to Professor Carl D. Hoff on the occasion of his 60th birthday

In 2001, Sharpless and co-workers defined the concept of "click chemistry" and the criteria for a transformation to be considered a "click" reaction.^[1] Since then, the coppercatalyzed reaction of azides with alkynes to produce 1,2,3triazoles regioselectively^[2] (the 1,3-dipolar Huisgen cycloaddition^[3]) has become the most widely used click reaction. As a result of its mild conditions and high efficiency, this reaction has found a myriad of applications in biology and materials science.^[4] Less attention has been focused on the development of novel copper(I)-based well-defined catalytic systems^[5–7] and on decreasing the amount of copper used.^[5b,8] This last point is extremely important for future industrial applications and might be one of the last challenges to overcome for this transformation.

We recently reported the remarkable activity of [(NHC)CuX] complexes (NHC=N-heterocyclic carbene; X = Cl, Br) in the Huisgen cycloaddition.^[9] Such catalytic systems have already been applied to the preparation of triazole-containing carbanucleosides,^[10] porphyrins,^[11] and platinum-based anticancer drugs,^[12] as well as to the development of a latent catalyst for this transformation.^[13]

We also studied a family of cationic NHC-containing complexes of the general formula $[(NHC)_2Cu]X$ (X = PF₆, BF₄).^[14] During the examination of their activity in the hydrosilylation of ketones, we observed enhanced reactivity of these complexes relative to that of their neutral analogues [(NHC)CuCl].^[15] This improved activity was rationalized in terms of a more efficient pathway for the activation of the cationic precatalyst. Specifically, the second NHC ligand was postulated to have an active role in the catalytic cycle. Under hydrosilylation conditions, one NHC ligand is displaced by the base in the reaction mixture. We wondered if an alkyne

[*] Dr. S. Díez-González, Prof. Dr. S. P. Nolan
Institute of Chemical Research of Catalonia (ICIQ)
Av. Països Catalans 16, 43007 Tarragona (Spain)
Fax: (+34) 977-920-244
E-mail: sdiez@iciq.es
snolan@iciq.es
Homepage: http://www.iciq.es/english/research/grups_eng nolan/entrada.htm

^[**] The ICIQ Foundation and the Spanish Ministerio de Educación y Ciencia are gratefully acknowledged for financial support. S.D.G. thanks the MEC for financial support through the Torres Quevedo program. S.P.N. is an ICREA Research Professor. We thank Dr. Nicolas "La Click, c'est chic" Marion for insightful reading of this manuscript. NHC = N-heterocyclic carbene.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.200803289.



Scheme 1. Catalyst design.

could play a similar role to produce an active copper acetylide from $[(NHC)_2Cu]X$ species (Scheme 1).

We investigated the activity of [(NHC)₂Cu]X complexes 1-7 in the Huisgen cycloaddition with benzyl azide and phenylacetylene as model substrates (Table 1). The reactions were conducted on water with 2 mol% of the copper catalyst. The SIMes-containing complexes 4a and 4b displayed poor activity under these conditions, in contrast to the results of our previous studies with [(NHC)CuX] complexes.^[9] This lack of reactivity can not simply be correlated with the medium steric bulk of the SIMes ligand, as complexes 7a and 7b with very bulky ItBu ligands were also ineffective.^[16] [(ICy)₂Cu]PF₆ (5a) provided the best result: The cycloaddition reached completion within 90 min. No general trend in reactivity was found for this reaction with respect to the counterion of the metal complex. Further optimization with complex 5a showed that acetonitrile, instead of water, was the most suitable solvent for this transformation.[17] Moreover, the reaction was found to proceed smoothly under neat conditions with a significantly decreased catalytic loading.

The scope of the reaction was examined with 5a (0.5 mol%) under neat conditions at room temperature (Scheme 2). All reactions proceeded smoothly to completion

© 2008 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim



Communications

Table 1: Catalyst screening.

	Ph	^ _{N3 +}	[(NH	C)₂Cu]X 1 – 7 (2 mol %) water, RT Ph	N ^N N N 8a Ph		
[(NHC) ₂ Cu]PFe	5	<i>t</i> [h]	Conv. [%] ^[a]	[(NHC) ₂ Cu]BF	<i>t</i> [h]	Conv. [%] ^[a]	
[(IPr) ₂ Cu]PF ₆	la	18	71	[(IPr) ₂ Cu]BF ₄	1 b	8	100
[(SIPr) ₂ Cu]PF ₆	2a	5	100	[(SIPr) ₂ Cu]BF ₄	2 b	5	100
[(IMes) ₂ Cu]PF ₆	3 a	6	100	[(IMes) ₂ Cu]BF ₄	3 b	6	100
[(SIMes) ₂ Cu]PF ₆	4 a	18	5	[(SIMes) ₂ Cu]BF ₄	4 b	18	13
[(ICy) ₂ Cu]PF ₆	5 a	1.5	99	[(ICy) ₂ Cu]BF ₄	5 b	5	95
[(IAd) ₂ Cu]PF ₆	6 a	5	100	[(IAd)₂Cu]BF₄	6 b	3	100
[(ItBu) ₂ Cu]PF ₆	7 a	18	76	[(ItBu) ₂ Cu]BF ₄	7 b	18	35

[a] The conversion (determined by ¹H NMR spectroscopy) is an average of the values for at least two independent experiments.

in short reaction times (from minutes to 9 h), and triazoles 8 were isolated in excellent yields and high purity after simple filtration or evaporation. According to the "click laws", no precautions to exclude oxygen or moisture were taken in any of these reactions. Neither copper disproportionation with the precipitation of metallic copper nor copper oxidation was observed under the optimized reaction conditions, which shows, once more, the high ability of NHCs to stabilize copper(I) species.^[18]

Electron-rich, electron-poor, and hindered alkynes were found to be suitable cycloaddition partners, as well as enynes. Benzyl, alkyl, and aryl azides were successfully employed. A number of functional groups were tolerated, such as alcohol, ketone, ester, amine, pyridine, nitrile, and halogen functionalities. No direct correlation can be drawn between the



Scheme 2. $[(ICy)_2Cu]PF_6$ -catalyzed formation of triazoles.

electronic nature of the reactants and the outcome of the reaction. As we had observed previously,^[9] the physical state of the reaction product is critical: Longer reaction times were required for the formation of oily or low-melting-point triazoles. In these cases, the results were improved simply by increasing the reaction temperature to 40 °C.^[19]

The high activity of this catalytic system naturally led us to examine the possibility of further decreasing the amount of metal used in this transformation. Again with benzyl azide and phenylacetylene as model

substrates, we tested different catalyst loadings at various temperatures under neat conditions. In this particular case, reactions at 40 or 50°C were faster than those at room temperature and could be carried out with the lowest catalyst loadings.^[20] However, minor formation of the 1,5-disubstituted triazole regioisomeric to 8a was observed consistently by GC and ¹H NMR spectroscopy after prolonged reaction times (8 h at 40 °C and 5 h at 50 °C). Thus, extreme caution is required to avoid undesired thermal processes. Results for the formation of a number of triazoles are presented in Table 2.

In all cases, the loading of 5a could be lowered at least to 300 ppm at room temperature or 100 ppm at 40 °C. Under these conditions, good conversion was still attained after short to reasonable reaction times. The best results were observed for the synthesis of triazole 8a: A copper loading as low as 40 ppm led to a remarkable turnover number (TON) above 20000 and a turnover frequency above $5000 h^{-1}$.^[21] It is important to note that, under these reaction conditions, the

Table 2: [(ICy)2Cu]PF6-catalyzed synthesis of triazoles at low catalyst loadings.

R ¹ —N ₃ +	$\equiv -R^2$	[(ICy)₂Cu]PF ₆ neat		R ¹ _N, N, N 8 R ²	
8	Т [°С]	[Cu] [ppm]	<i>t</i> [h]	Conv. [%] ^[a]	TON
	RT	50	48	80	16000
Ph' '	40	50	8	89	17800
8a Ph	50	40	4	81	20250
Ph N ^N N 8d N	RT	75	6	91	12133
Hept-N ^N N 8f Ph	RT	200	20	72	3600
Ph~N ^N N	RT	300	43	85	2833
8i \}_3Cl	40	100	18	70	7000
Ph ^(h) 2N ^N N 8j NMe ₂	RT 40	300 100	40 18	45 71	1500 7100

[a] The conversion (determined by GC) is an average of the values for at least two independent experiments.

use of our previous catalyst [(SIMes)CuBr] led to no conversion after 24 h.

To gain some insight into the activation pathway of the precatalyst, we next carried out a number of stoichiometric experiments. Whereas no interaction was observed between benzyl azide and 5a in CD₃CN, the same copper complex was completely consumed in minutes in the presence of phenyl-acetylene. The equimolar formation of two new species was evidenced by ¹H NMR spectroscopy (Figure 1). One of them



Figure 1. ¹H NMR spectrum for the reaction of 5a with phenylacetylene. Cy = cyclohexyl.

was identified unequivocally as $ICy \cdot HPF_6$ (**B**) by comparison with a pure sample.^[22] The second species was also isolated cleanly and identified as the mononuclear copper(I) acetylide A by NMR spectroscopy. To determine the identity of A unambiguously, we carried out the same reaction with the IPrcontaining complex 1a. In this case, no reaction was observed at room temperature. However, when the reaction mixture was heated overnight at 80°C, the formation of two new species was observed, as well as remaining starting material. The new products were separated and identified as the known complex $[(IPr)Cu-C \equiv CPh]^{[23]}$ and $IPr \cdot HPF_6$. The lower reactivity of 1a relative to that of 5a can be correlated directly to its moderate catalytic activity (Table 1). Thus, it appears that one of the NHC ligands on the copper center can act as a base, deprotonating the alkyne, to initiate the catalytic cycle.

A proposed mechanism for this transformation is depicted in Scheme 3. The bis(NHC) complex **5a** would be transformed into an acetylide **A** and an azolium salt **B** by reaction with the alkyne. Upon the interaction of **A** with the azide, the reaction would follow the pathway commonly accepted for this transformation^[24] to form a triazolide intermediate **C**, the reaction of which with the azolium salt **B** would lead to the formation of the expected triazole **8a** and the regeneration of the catalyst. However, a binuclear mechanism can not be ruled out at this time.^[25] We are currently carrying out theoretical calculations to clarify this point.

In an attempt to isolate intermediate C, we treated isolated complex **A** with benzyl azide in CD₃CN. However, we observed the formation of a complex mixture of unknown products as well as considerable precipitation, presumably of copper salts. We attribute this result to the extreme instability



Scheme **3.** Postulated mechanism. Bn = benzyl.

of **C** in the absence of a proton source. In contrast, when a mixture of **A** and **B** was treated with the azide, clean formation of triazole 8a was observed, along with the regeneration of [(ICy)₂Cu]PF₆.

Further experiments to isolate the triazolide intermediate C met with limited success. The treatment of 5a with an equimolar amount of phenylacetylene and a slight excess of benzyl azide in acetonitrile led either to the decomposition of the reaction mixture or to the direct formation of triazole 8a. Nevertheless, we were able to observe by ¹H NMR spectroscopy the consumption of the starting materials and the formation of a new compound. The spectrum of the new species was attributed to \mathbf{C} ;^[20] however, the formation of this intermediate as a minor component of the mixture and its high instability precluded its full characterization. Moreover, upon acidic hydrolysis this crude mixture rapidly evolved to triazole 8a and ICy·HPF₆. These findings are in line with a report by Straub and co-workers, who recently isolated a NHC triazolide complex in which the environment around the copper center was highly encumbered, probably stabilizing the complex.^[26]

In conclusion, we have developed a new catalytic system, $[(ICy)_2Cu]PF_6$ (**5a**) without a solvent, for the [3+2] cycloaddition of azides and alkynes under click conditions. This system has been shown to be broad in scope and highly efficient (TOFs up to 5000 h⁻¹) even at very low catalyst loadings (down to 40 ppm). Furthermore, preliminary mechanistic studies have shown the ability of the NHC ligand on the copper center to act as a base and deprotonate the starting alkyne to initiate the catalytic cycle. Further applications of this catalyst system, especially with a low catalyst loading, are currently under investigation in our laboratory.

Received: July 7, 2008 Revised: August 14, 2008 Published online: October 10, 2008

Communications

Keywords: alkynes · azides · click chemistry · copper · N-heterocyclic carbenes

- H. C. Kolb, M. G. Finn, K. B. Sharpless, Angew. Chem. 2001, 113, 2056–2075; Angew. Chem. Int. Ed. 2001, 40, 2004–2021.
- [2] a) C. W. Tornøe, C. Christensen, M. Meldal, J. Org. Chem. 2002, 67, 3057–3064; b) V. V. Rostovtsev, L. G. Green, V. V. Fokin, K. B. Sharpless, Angew. Chem. 2002, 114, 2708–2711; Angew. Chem. Int. Ed. 2002, 41, 2596–2599.
- [3] a) R. Huisgen, Pure Appl. Chem. 1989, 61, 613–628; b) A.
 Padwa in Comprehensive Organic Synthesis, Vol. 4 (Ed.: B. M. Trost), Pergamon, Oxford, 1991, pp. 1069–1109.
- [4] For reviews, see: a) J. E. Moses, A. D. Moorhouse, *Chem. Soc. Rev.* 2007, *36*, 526–539; b) A. Dondoni, *Chem. Asian J.* 2007, *2*, 700–708; c) J.-F. Lutz, *Angew. Chem.* 2007, *119*, 1036–1043; *Angew. Chem. Int. Ed.* 2007, *46*, 1018–1025; d) Special issue: Click Chemistry, *QSAR Comb. Sci.* 2007, *26*, 1110–1323.
- [5] a) T. R. Chan, R. Hilgraf, K. B. Sharpless, V. V. Fokin, Org. Lett.
 2004, 6, 2853–2855; b) V. O. Rodionov, S. I. Presolsky, S. Gardinier, Y.-H. Lim, M. G. Finn, J. Am. Chem. Soc. 2007, 129, 12696–12704; c) S. S. Gupta, J. Kuzelka, P. Singh, W. G. Lewis, M. Manchester, M. G. Finn, Bioconjugate Chem. 2005, 16, 1572–1579.
- [6] F. Pérez-Balderas, M. Ortega-Muñoz, J. Morales-Sanfrutos, F. Hernández-Mateo, F. G. Calvo-Flores, J. A. Calvo-Asín, J. Isac-García, F. Santoyo-González, Org. Lett. 2003, 5, 1951–1954.
- [7] B. Gerard, J. Ryan, A. B. Beeler, J. A. Porco, Jr., *Tetrahedron* 2006, 62, 6405-6411.
- [8] N. Candelon, D. Lastécouères, A. K. Diallo, J. Ruiz Aranzaes, D. Astruc, J.-M. Vincent, *Chem. Commun.* 2008, 741–743.
- [9] S. Díez-González, A. Correa, L. Cavallo, S. P. Nolan, *Chem. Eur. J.* 2006, *12*, 7558–7564.
- [10] J. Broggi, S. Díez-González, J. L. Petersen, S. Berteina-Raboin, S. P. Nolan, L. A. Agrofoglio, *Synthesis* 2008, 141–148.
- [11] M. Séverac, L. Le Pleux, A. Scarpaci, E. Blart, F. Odobel, *Tetrahedron Lett.* 2007, 48, 6518–6522.
- [12] A. Maisonial, P. Serafin, M. Traïkia, E. Debiton, V. Théry, D. J. Aitken, P. Lemoine, B. Voissat, A. Gautier, *Eur. J. Inorg. Chem.* 2008, 298–305.
- [13] a) S. Díez-González, E. D. Stevens, S. P. Nolan, *Chem. Commun.* 2008, 4747–4749.
- [14] a) S. Díez-González, N. M. Scott, S. P. Nolan, *Organometallics* 2006, 25, 2355-2358; b) S. Díez-González, E. D. Stevens, N. M. Scott, J. L. Petersen, S. P. Nolan, *Chem. Eur. J.* 2008, 14, 158-168.

- [15] a) H. Kaur, F. K. Zinn, E. D. Stevens, S. P. Nolan, Organometallics 2004, 23, 1157–1160; b) S. Díez-González, H. Kaur, F. K. Zinn, E. D. Stevens, S. P. Nolan, J. Org. Chem. 2005, 70, 4784–4796.
- [16] For discussion on the stereoelectronic parameters of NHC ligands, see: a) T. Strassner, *Top. Organomet. Chem.* 2004, *13*, 1–20; b) L. Cavallo, A. Correa, C. Costabile, H. Jacobsen, *J. Organomet. Chem.* 2005, *690*, 5407–5413; c) S. Díez-González, S. P. Nolan, *Coord. Chem. Rev.* 2007, *251*, 874–883.
- [17] During the optimization studies, complete conversion was observed in dimethyl sulfoxide (2 h), water (1.5 h), *N*,*N*-dimethylformamide (1.5 h), THF (1 h), acetone (0.5 h), and MeCN (0.2 h) in the presence of **5a** (2 mol%). Sluggish reactions were observed when alcohols were used as solvents.
- [18] For reviews, see: a) S. Díez-González, S. P. Nolan, *Synlett* 2007, 2158–2167; b) S. Díez-González, S. P. Nolan, *Aldrichimica Acta* 2008, 41, 43–51.
- [19] For example, triazole 8g was produced quantitatively at 40 °C in 3 h instead of 9 h. For triazoles 8n and 8o, a reaction time of only 1.5 h was required at 40 °C.
- [20] For further details, see the Supporting Information.
- [21] Vincent and co-workers recently reported a copper(I)-tren catalyst (tren = tris(2-aminoethylamine)) that promoted the Huisgen cycloaddition with high turnover numbers (TONs). TON values of up to 54000 were observed for the preparation of **8a** at 60 °C in 24 h in the absence of a solvent.^[8] However, during our studies, even for the formation of the same triazole, we observed the non-negligible formation of the 1,5-regioisomer at 50 °C after only 5 h.
- [22] F. Liu, W. Chen, X. You, J. Chem. Crystallogr. 2002, 32, 27-31.
- [23] L. A. Goj, E. D. Blue, C. Munro-Leighton, T. B. Gunnoe, J. L.
- Petersen, *Inorg. Chem.* 2005, 44, 8647–8649.
 [24] F. Himo, T. Lovell, R. Hilgraf, V. V. Rostovtsev, L. Noodleman, K. B. Sharpless, V. V. Fokin, *J. Am. Chem. Soc.* 2004, 127, 210– 216.
- [25] For a review, see: V. D. Bock, H. Hiemstra, J. H. van Maarseveen, *Eur. J. Org. Chem.* **2006**, 51–68; see also: a) B. F. Straub, *Chem. Commun.* **2007**, 3868–3870; b) M. Ahlquist, V. V. Fokin, *Organometallics* **2007**, 26, 4389–4391.
- [26] C. Nolte, P. Mayer, B. F. Straub, *Angew. Chem.* 2007, *119*, 2147–2149; *Angew. Chem. Int. Ed.* 2007, *46*, 2101–2103; for a stable gold-triazolide complex, see: D. V. Partyka, J. B. Updegraff III, M. Zeller, A. D. Hunter, T. G. Gray, *Organometallics* 2007, *26*, 183–186.