Ruthenium-Catalyzed Oxidative Heck Reactions**

Edward J. Farrington, John M. Brown,* C. F. J. Barnard, and E. Rowsell

The coupling of organic electrophiles with alkenes is one of the cornerstones of Pd complex catalysis. The basis of the chemistry involved lies in early contributions by Mizoroki^[1] and Heck.^[2] Many of Heck's examples involve stoichiometric reactions of organomercurials and other potential organic nucleophiles in addition to Pd salts. Since then, but with a notable recent exception (see below),^[3] palladium complexes have been used essentially exclusively in Heck catalysis.

Ruthenium complexes are readily accessible, but less widely utilized in the catalysis of C–C coupling reactions.^[4] Faller and Chase demonstrated that complex **1b**, prepared from the dichloride **1a** and PhMgBr, reacted with ethene and AgSbF₆ to form styrene and the corresponding Ru–H complex **2**.^[5] An intramolecular variant through insertion of ethene into the Ru–C bond of complex **3** was reported by Pfeffer and co-workers.^[6] These stoichiometric reactions



Blount's Court, Sonning Common, Reading RG4 9NH (UK)

- [**] We thank the EPSRC for CASE support (to EJF) with Johnson Matthey, who also provided the Ru salts. Dr. R. T. Aplin was very helpful in acquiring electrospray mass spectra.
- Supporting information for this article is available on the WWW under http://www.angewandte.com or from the author.

provided the key to catalysis, since reversion from a Ru-H complex to the halide together with a compatible method for the formation of a Ru-C bond would, in principle, complete the cycle.

In preliminary experiments it was established that the reaction of **1a** with PhSnBu₃ or PhB(OH)₂ occurred readily to give **1c** as a stable product.^[7] This observation formed the basis of a successful stoichiometric reaction between PhB(OH)₂ and methyl propenoate, promoted by complex **1a** in the presence of NEt₃ under mild conditions. The use of more than 2 equivalents of Cu(OAc)₂ to reoxidize the expected Ru-H complex led to catalytic turnover; the process was more efficient when PPh₃ was omitted from the initial Ru catalyst (Scheme 1).^[8] Optimization experiments



21% D 49% isolated

169



Scheme 1. Ruthenium-catalyzed Heck reactions: A) optimum conditions; B) with stereospecifically labeled butyl propenoate.

demonstrated that other oxidizing agents (e.g. Me₃NO, Fremy's salt) also promoted the reaction, but less effectively than Cu^{II}. Significant improvements in the base component were achieved by this approach: 3-quinuclidone gave the highest turnover (Scheme 1 A). The conditions were successfully applied with *n*-butyl propenoate and other ArB(OH)₂ species (Ar = *p*-CHO, OMe, OCF₃).

To compare the pathway with the more familiar Pd case, the addition of PhB(OH)₂ to butyl (*Z*)-3-D-propenoate^[9] was effected (Scheme 1B). The label was largely, but not completely, absent in the product. This is in accord with partial stereoselectivity in the catalytic process, most probably through stereospecific *cis* addition of Ru–Ph^[10] to the alkene followed by some randomization (Ru keto-enol equilibria?)^[11] prior to stereospecific Ru-H/D *cis* elimination. Electrospray mass spectrometry revealed intermediates in the catalytic cycle, best observed with 4-BrC₆H₄B(OH)₂ in a CH₃CN matrix (Scheme 2). Before the addition of the alkene, three monomeric complexes **5a**–**c** and one dimeric cationic complex **5d** can be characterized on the basis of the typical

Angew. Chem. Int. Ed. 2002, 41, No. 1 © WILEY-VCH Verlag GmbH, 69451 Weinheim, Germany, 2002 1433-7851/02/4101-0169 \$ 17.50+.50/0

COMMUNICATIONS



Scheme 2. Cationic species observed by electrospray mass spectrometry recorded on a Micromass BioQ II-ZS at a cone voltage of +20 V, Ru-102, Br-81, Cl-35 ion specified. Reagents and conditions: a) [Ru(p-cymene)Cl₂]₂ (3.26×10^{-5} mol), *p*-bromobenzeneboronic acid (3.26×10^{-5} mol), NEt₃ (3.26×10^{-5} mol), MeCN (20 mL), 30 min, room temperature; b) same conditions as (a), but allowed to stand with a tenfold excess of methyl propenoate for 15 h at -20 °C before assay.

isotope distributions in the parent molecular ion. In the presence of methyl propenoate, the same set of monomeric ions are observed together with two distinct new halide-free species, corresponding to monoalkene complexes with one or two coordinated acetonitrile ligands. The simplest representation in which these preserve an 18e configuration at the ruthenium center is as the aryl and alkyl complexes **6a** before, and **6b** after the expected migration step; ring slippage is an alternative possibility.^[12] The nature of propenoate binding



was explored through the use of methylene malonate 7, which gave rise to a strong acetonitrile-free molecular ion 8 for the coordinated alkene complex, under conditions in which the parent malonate 9 remains passive.

The inertness of the aryl bromide residue towards ruthenium (i.e. dominance of nucleophilic attack by the boronic acid) indicated valuable synthetic potential. Several haloaryl boronic acids were subjected to our best conditions of catalytic turnover, with *n*-butyl propenoate as the alkene component. The results are recorded in Scheme 3, and



Scheme 3. Ru-catalyzed reactions of haloaryl boronic acids. Reagents and conditions: a) boronic acid (0.4 mmol), [Ru(p-cymene)Cl₂]₂ (6.2 mg, 0.02 mmol), Cu(OAc)₂ (201 mg, 1 mmol), quinuclidin-3-one (126 mg, 1 mmol), and butyl propenoate (155 mg, 1.2 mmol), toluene, 24 h, room temperature, X = m-Cl (77%), X = p-Cl (93%), X = p-Br (63%),^[13] X = m-Br (96%), X = m-I (76%), X = p-I (77%); b) butyl *p*-iodocinnamate (50 mg, 0.15 mmol), tris(*o*-tolyl)phosphane (4.56 mg, 0.015 mmol), reiethylamine (76 mg, 0.76 mmol), DMF, 24 h, room temperature, **10** (90%); dba = *trans*, *trans*-dibenzylideneacetone, DMF = *N*,*N*-dimethyl-formamide. For product characterization, see Supporting Information.

demonstrate the generality of the selective C–B activation and transfer. Good yields are maintained over a range of boronic acids substituted with Cl, Br, or I. In one case, a tandem Pd- and Ru-catalyzed coupling reaction in which C–B is activated first, followed by C–X, gave the unsymmetrical phenylene-bis-propenoate **10**.^[14]

In related work, it was demonstrated that aryl stannanes,^[15] aryl silanols,^[16]and aryl boronic acids^[17] can be coupled with electrophilic alkenes under Pd(OAc)₂ catalysis (5–10 mol %). In the first two cases Cu^{II} is present as a reoxidant. In the last case, the authors favor, but do not prove, an oxidative addition of ArB(OH)₂ as the initial step; the reaction works only in AcOH. Very recently, Lautens and co-workers observed the coupling of styrenes with an excess of functionally diverse boronic acids in aqueous solution, to form *E* stilbenes in the absence of a formal oxidant.^[3] The relationship of these procedures to ours is intriguing, but as yet unclear.

The synthetic potential of the ruthenium Heck reaction is being actively investigated further.^[18]

Experimental Section

Butyl (E)-3-bromocinnamate: 3-bromobenzeneboronic acid (81 mg, 0.4 mmol), [Ru(p-cymene)Cl₂]₂ (6.2 mg, 0.02 mmol), Cu(OAc)₂ (201 mg, 1 mmol), and 3-quinuclidinone (126 mg, 1 mmol) were combined in a dry Schlenk tube, which was sealed and placed under vacuum for 30 min. The Schlenk tube was flushed with argon, and butyl propenoate (155 mg, 1.2 mmol) and freshly distilled toluene (10 mL) were added through a syringe. The reaction mixture was stirred for 24 h at ambient temperature. The volatile components were removed in vacuo, and the product was purified by flash column chromatography on silica gel (diethyl ether) to give butyl (E)-3-bromocinnamate (112 mg, 96%). M.p. 36 $^{\circ}\mathrm{C};\ ^{1}\mathrm{H}\,\mathrm{NMR}$ (500 MHz, CDCl₃): δ = 0.96 (3H, t, J = 7.3 Hz; CH₃), 1.44 (2H, m; CH₂), 1.70 (2 H, m; CH₂), 4.22 (2 H, t, J = 6.7 Hz; OCH₂), 6.46 (1 H, d, J = 16 Hz; H1), 7.26 (1 H, dd, J = 8 Hz and 7.95 Hz; H5'), 7.46 (1 H, d, J = 8 Hz; H4'), 7.52 (1 H, d, J = 7.95 Hz; H6'), 7.61 (1 H, d, J = 16 Hz; H3), 7.68 (1 H, s; H2'); ^{13}C NMR (125.8 MHz, CDCl₃): δ = 13.6 (CH₃), 19.1 (CH₂), 30.6 (CH₂), 64.5 (OCH2), 119.7 (C2), 122.9, 126.5, 130.3, 130.6, 132.9, 136.4, (C1', C2', C3', C4', C5', C6'), 142.7 (C3) 166.5 (C1); IR (Nujol): $\tilde{v}_{max} = 2920.9$, 1727.3 (C=O), 1641.7 cm⁻¹ (C=C); HR-MS: calcd for C₁₃H₁₆O₂Br: 283.0335, found: 283.0335; MS (atmospheric pressure chemical ionisation; APCI +): m/z (%): 284 [MH +], 226 [M - Bu].

Coupling of butyl (*Z*)-3-D-propenoate: Benzeneboronic acid (10 mg, 8.2 × 10⁻⁵ mol), [Ru(*p*-cymene)Cl₂]₂ (1.25 mg, 4.1×10^{-7} mol), and Cu(OAc)₂ (40.9 mg, 0.2 1mmol) were combined in a vial, and THF (0.9 mL) and

1433-7851/02/4101-0170 \$ 17.50+.50/0 Angew. Chem. Int. Ed. 2002, 41, No. 1

triethylamine (0.1 mL) were added. Butyl (*Z*)-3-D-propenoate was added through a syringe. The vial was sealed, and the reaction mixture was stirred at ambient temperature for 17 h. The solvent was removed in vacuo, and the product was purified by filtration through a short plug of silica gel (diethyl ether) to afford butyl (*E*)-cinnamate (8.2 mg, 49%). B.p. 280–284 °C; ¹H NMR (500 MHz, CDCl₃): $\delta = 0.99$ (3H, t, J = 7.3 Hz; CH₃), 1.43 (2H, m; CH₂), 1.72 (2H, m; CH₂), 4.23 (2H, t, J = 6.7 Hz; OCH₂), 6.48 (1H, d, J = 16 Hz; H2), 7.40 (3H, m; ArH), 7.53 (2H, m; ArH), 7.77 (1H, d, J = 16 Hz; H3); ¹³C NMR (125.8 MHz, CDCl₃): $\delta = 13.6$ (CH₃), 19.1 (CH₂), 30.6 (CH₂), 64.3 (OCH₂), 118.2 (C2), 127.9, 128.7, 130.1, 134.3, (Ar-C), 144.4 (C3), 167.0 (C1); UV/Vis: \tilde{v}_{max} (MeOH) = 276 nm (ε = 22000); MS (CI +): m/z (%): 205 [*M*H⁺].

Received: August 10, 2001 [Z17712]

- [1] T. Mizoroki, K. Mori, A. Ozaki, Bull. Chem. Soc. Jpn. 1971, 44, 581.
- [2] R. F. Heck, J. Am. Chem. Soc. 1968, 90, 5518-5526; R. F. Heck, Acc. Chem. Res. 1979, 12, 146-151; R. F. Heck, Org. React. 1982, 27, 345-391, and references therein.
- [3] M. Lautens, A. Roy, K. Fukuoka, K. Fagnou, B. Martin-Matute, J. Am. Chem. Soc. 2001, 123, 5358-5359.
- [4] Notably, the C-H activation discovered by Murai: S. Murai, F. Kakiuchi, S. Sekine, Y. Tanaka, A. Kamatani, M. Sonoda, N. Chatani, *Nature* 1993, 366, 529-531; F. Kakiuchi, S. Sekine, Y. Tanaka, A. Kamatani, M. Sonoda, N. Chatani, S. Murai, *Bull. Chem. Soc. Jpn.* 1995, 68, 62-83; S. Busch, W. Leitner, *Adv. Synth. Catal.* 2001, 343, 192-195; for an intramolecular variant, see: H. Weissman, X. P. Song, D. Milstein, *J. Am. Chem. Soc.* 2001, 123, 337-338; for Ru cyclo-isomerizations, see: Y. Yamamoto, Y.-i. Nakagai, N. Ohkoshi, K. Itoh, *J. Am. Chem. Soc.* 2001, 123, 6372-6380; for a general review, see: B. M. Trost, F. D. Toste, A. B. Pinkerton, *Chem. Rev.* 2001, 101, 2067-2096.
- [5] J. W. Faller, K. J. Chase, Organometallics 1995, 14, 1592-1600.
- [6] V. Ritleng, J. P. Sutter, M. Pfeffer, C. Sirlin, *Chem. Commun.* 2000,129-130.
- [7] The X-ray structure of complex 1c has been determined: A. R. Cowley, E. J. Farrington, *Acta Crystallogr. Sect. E*, submitted.
- [8] A referee has suggested that the Cu(OAc)₂ may activate the aryl boronic acid directly, on the basis of reported Cu-catalyzed nucleo-philic substitutions: P. S. Herradura, K. A. Pendola, R. K. Guy, Org. Lett. 2000, 2, 2019–2022; we have observed a reaction between PhB(OH)₂ and complex 4 at a rate commensurate with the observed turnover, thus trapping the initially formed intermediate by addition of PPh₃ to form 1 c. In the presence of the acetate analogue of complex 4 (5 mol%), the reaction between PhB(OH)₂ and Cu(OAc)₂ yields PhOAc as the only characterized organic product in 61% yield; see: P. Y. S. Lam, G. Vincent, C. G. Clark, S. Deudon, P. K. Jadhav, *Tetrahedron Lett.* 2001, 42, 3415–3418.
- [9] R. K. Hill, G. R. Newkome, J. Org. Chem., 1969, 34, 740-741.
- [10] In the case of Pd, this is assumed on the basis of the overall stereochemical course of catalysis to be a syn addition of Pd–R and then a syn elimination of Pd–H; R. F. Heck, J. Am. Chem. Soc. 1969, 91, 6707–6715; H. A. Dieck, R. F. Heck, J. Am. Chem. Soc. 1974, 96, 1133–1136.
- [11] M. Ikeda, S. El-Bialy, T. Yakura, Heterocycles 1999, 51, 1957-1970.
- [12] M. A. Bennett, Z. Lu, X. Wang, M. Bown, D. C. R. Hockless, J. Am. Chem. Soc. 1998, 120, 10409-10415.
- [13] M. Miura, H. Hashimoto, K. Itoh, M. Nomura J. Chem. Soc. Perkin Trans. 1 1990, 2207–11.
- [14] For tandem Heck reactions of dihalobenzenes, see, for example: J. E. Plevyak, J. E. Dickerson, R. F. Heck, J. Org. Chem. 1979, 44, 4078–4083.
- [15] K. Hirabayashi, J. Ando, Y. Nishihara, A. Mori, T. Hiyama, Synlett 1999, 99–101; K. Hirabayashi, J. Ando, J. Kawashima, Y. Nishihara, A. Mori, T. Hiyama, Bull. Chem. Soc. Jpn. 2000, 73, 1409–1417.
- [16] K. Hirabayashi, Y. Nishihara, A. Mori, T. Hiyama, *Tetrahedron Lett.* 1998, 39, 7893–7896.
- [17] C. S. Cho, S. Uemura, J. Organomet. Chem. 1994, 465, 85-92.
- [18] An oxidative palladium-catalyzed Heck reaction of boronates has been reported; X. L. Du, M. Suguro, K. Hirabayashi, A. Mori, T. Nishikata, N. Hagiwara, K. Kawata, T. Okeda, H. F. Wang, K. Fugami, M. Kosugi, Org. Lett. 2001, 3, 3313–3316.

Enantiopure Double-Helical Alkynyl Cyclophanes

De Lie An, Takehiko Nakano, Akihiro Orita, and Junzo Otera*

Chiral π -conjugated molecules have been the subject of extensive investigation from the standpoints of structural chemistry and material science.^[1] Double-helical molecules, in particular, are of great interest on account of their unique structural features as well as potential applications in optics and electronics. Several twisted alkynyl cyclophanes have been reported, but, unfortunately, they were obtained as racemates.^[2, 3] To the best of our knowledge, only one nonracemic double-helical molecule has been prepared. Namely, a cyclophane was synthesized by connecting a (+)-2,15-diethynyl[6]helicene auxiliary with ortho-phenylene bridges.^[4] However, this synthesis was rather lengthy, and only one enantiomer was obtained. Moreover, in the key coupling of the helicene with an o-diiodobenzene unit, the target molecule was obtained in less than 3 % yield. We report here on a rational synthesis and full characterization of double-helical alkynyl cyclophanes 1 of both enantiopure forms.^[5]

The synthetic route is shown in Scheme 1. Separately (R)and (S)-2,2'-diformyl-1,1'-binaphthyl (2)^[6] underwent carbon-carbon coupling, and the resulting diethynyl compound 3 was converted to the monosilylethynyl derivative 4. Exposure of this compound to an aryl iodide with the diethyltriazene function 5 afforded 6. The triazene derivatives 6 were transformed into iodides 7 or desilylated to give 8.^[7] Sonogashira coupling^[8] of 7 with 8 furnished 9. After conversion of 9 to 11 via 10 through successive functional group transformations, intramolecular Sonogashira coupling provided the desired cyclophanes 1 in enantiopure form (Table 1). These compounds formed white needlelike crystals upon recrystallization, but none of them were suitable for X-ray analyses. Then, we prepared racemic 1b by mixing equimolar amounts of (R,P)- and (S,M)-1b. Recrystallization of this mixture from CH2Cl2/hexane furnished crystals conducive to X-ray crystallographic analysis.^[9]

As is evident from the ORTEP view depicted in Figure 1, the cyclophane skeleton is twisted, resulting in the doublehelical motif. The C=C bonds are slightly deformed from linearity. Notably, the two binaphthyl groups differ significantly in the dihedral angle defined by the naphthalene planes $(68^{\circ} \text{ and } 78^{\circ}, \text{ respectively})$. The space-filling model (Figure 2) indicates that the symmetrical structure places the inside hydrogen atoms of the phenylene rings very close to each other. The resulting ring strain is passed on unsymmetrically into the binaphthyl termini in the crystal.

In contrast to the solid-state molecular structure, ${}^{1}H$ and ${}^{13}C$ NMR spectra of **1** (Table 1) are compatible with a single

1433-7851/02/4101-0171 \$ 17.50+.50/0

Angew. Chem. Int. Ed. 2002, 41, No. 1 © WILEY-VCH Verlag GmbH, 69451 Weinheim, Germany, 2002

^[*] Prof. Dr. J. Otera, Dr. D. L. An, T. Nakano, Dr. A. Orita Department of Applied Chemistry, Okayama University of Science Ridai-cho, Okayama 700-0005 (Japan) Fax: (+81)-86-256-4292 E-mail: otera@high.ous.ac.jp