## Studies toward the Total Synthesis of Gambieric Acids, Potent Antifungal Polycyclic Ethers: Convergent Synthesis of the CDEFG-Ring System

2005 Vol. 7, No. 12 2441–2444

ORGANIC LETTERS

## Kazushi Sato and Makoto Sasaki\*

Laboratory of Biostructural Chemistry, Graduate School of Life Sciences, Tohoku University, Tsutsumidori-Amamiya, Aoba-ku, Sendai 981-8555, Japan masasaki@bios.tohoku.ac.jp

Received April 8, 2005

ABSTRACT



A convergent synthetic route to the CDEFG-ring system of gambieric acids, potent antifungal polycyclic ether marine natural products, has been developed. The present synthesis features convergent union of the CD- and G-rings through esterification, formation of the E-ring as a lactone form, stereoselective allylation to set the  $C_{26}$  stereocenter, and ring-closing metathesis reaction to construct the nine-membered F-ring.

A number of polycyclic ether natural products with potent and diverse biological activities have been isolated from marine sources.<sup>1</sup> Gambieric acids A–D (1–4, Figure 1) were isolated by the Yasumoto group from the culture media of the marine dinoflagellate, *Gambierdiscus toxicus*, which is the causative organism responsible for ciguatera seafood poisoning.<sup>2</sup> These polycyclic ethers have attracted considerable interest because of their potent antifungal activity against a variety of filamentous fungi.<sup>2c</sup> Their antifungal activity against *Aspergillus niger* by the paper disk method was 2000 times greater than that of amphotericin B, and they exhibited only moderate toxicity toward mammalian cells. These findings make them potential lead compounds for the discovery of antifungal agents.<sup>2</sup> In this letter, we describe a



Figure 1. Structures of gambieric acids A–D.

convergent synthesis of the CDEFG-ring system 5 of gambieric acids.<sup>3</sup>

During our studies toward the total synthesis of ciguatoxins, we have previously reported a convergent route to the

<sup>(1)</sup> For reviews on marine polycyclic ethers, see: (a) Yasumoto, T.; Murata, M. *Chem. Rev.* **1993**, *93*, 1897–1909. (b) Murata, M.; Yasumoto, T. *Nat. Prod. Rep.* **2000**, 293–314. (c) Yasumoto, T. *Chem. Rec.* **2001**, *3*, 228–242.

<sup>(2) (</sup>a) Nagai, H.; Torigoe, K.; Satake, M.; Murata, M.; Yasumoto, T.; Hirota, H. J. Am. Chem. Soc. **1992**, 114, 1102–1103. (b) Nagai, H.; Murata, M.; Torigoe, K.; Satake, M.; Yasumoto, T. J. Org. Chem. **1992**, 57, 5448– 5453. (c) Nagai, H.; Mikami, Y.; Yazawa, K.; Gonoi, T.; Yasumoto, T. J. Antibiot. **1993**, 46, 520–522. (d) Morohashi, A.; Satake, M.; Nagai, H.; Oshima, Y.; Yasumoto, T. Tetrahedron **2000**, 56, 8995–9001.

FGH-ring system.<sup>4</sup> The synthesis features an intramolecular radical cyclization to form the seven-membered G-ring with complete stereocontrol and a ring-closing metathesis (RCM) to construct the nine-membered F-ring. Subsequently, this approach was further modified and refined by the Hirama group, culminating in the first total synthesis of ciguatoxin CTX3C.<sup>5</sup> Initially, we planned to synthesize the CDEFG-ring system **5** of gambieric acids by applying this strategy (Scheme 1). The nine-membered F-ring of **5** could be



constructed by RCM from a precursor diene. In turn, the seven-membered E-ring was envisioned to be formed by an intramolecular radical cyclization of 8, which would be derived from alcohol 6 and carboxylic acid 7.

The synthesis of alcohols **6a** and **6b** started with compound **9**<sup>6</sup> (Scheme 2). Protection as the TBS ether and oxidative cleavage of the double bond, followed by Wittig reaction with methyl (triphenylphosphoranilidene)acetate, afforded enoate **10** in 81% yield for the three steps. Treatment of **10** with methylmagnesium bromide in the presence of TMSCI and *i*-propylsalicylaldimine copper(II) complex **11**<sup>7</sup> (THF, -45 °C) gave the desired adduct **12** as the sole product in high yield.<sup>8</sup> The stereochemistry of the newly generated stereocenter was confirmed by conversion to lactone **13** and its NMR analysis as shown. DIBALH reduction of **12** 

(4) (a) Sasaki, M.; Noguchi, T.; Tachibana, J. Org. Chem. 2002, 67, 3301–3310. (b) Sasaki, M.; Noguchi, T.; Tachibana, K. Tetrahedron Lett. 1999, 40, 1337–1340. (c) Sasaki, M.; Inoue, M.; Noguchi, T.; Takeichi, A.; Tachibana, K. Tetrahedron Lett. 1998, 39, 2783–2786.

(5) (a) Hirama, M.; Oishi, T.; Uehara, H.; Inoue, M.; Maruyama, M.; Oguri, H.; Satake, M. *Science* **2001**, *294*, 1904–1907. (b) Inoue, M.; Uehara, H.; Maruyama, M.; Hirama, M. Org. Lett. **2002**, *4*, 4551–4554. (c) Inoue, M.; Hirama, M. Synlett **2004**, *4*, 577–595. (d) Inoue, M.; Miyazaki, K.; Uehara, H.; Maruyama, M.; Hirama, M. Proc. Natl. Acad. Sci. U.S.A. **2004**, *101*, 12013–12018. (e) Inoue, M.; Hirama, M. Acc. Chem. Res. **2004**, *37*, 961–968.



(7) Sakata, H.; Aoki, Y.; Kuwajima, I. *Tetrahedron Lett.* **1990**, *31*, 1161–1164.

(8) Matsuo, G.; Kawamura, K.; Hori, N.; Matsukura, H.; Nakata, T. J. Am. Chem. Soc. **2004**, 126, 14374–14376.



produced primary alcohol **14** (99%), which upon benzylation followed by desilylation yielded alcohol **6a** in 88% yield for the two steps. Alcohol **14** was converted to a terminal olefin via *o*-nitrophenyl selenide by the method of Grieco and Nishizawa.<sup>9</sup> Subsequent desilylation afforded alcohol **6b** in 85% yield for the three steps.

The synthesis of carboxylic acid 7 commenced with alcohol 15,<sup>10</sup> which was converted to methyl ketone 16 by a four-step sequence in 82% overall yield (Scheme 3).



Following desilylation, treatment of the derived alcohol with ethyl propiolate in the presence of *N*-methylmorpholine (NMM) produced  $\beta$ -alkoxyacrylate **17** in 81% yield over the two steps. Treatment of **17** with samarium(II) iodide in the

<sup>(3)</sup> For synthetic studies of gambieric acids, see: (a) Kadota, I.; Oguro, N.; Yamamoto, Y. *Tetrahedron Lett.* **2001**, *42*, 3645–3647. (b) Kadota, I.; Takamura, H.; Yamamoto, Y. *Tetrahedron Lett.* **2001**, *42*, 3649–3651. (c) Clark, J. S.; Fessard, T. C.; Wilson, C. *Org. Lett.* **2004**, *6*, 1773–1776.



presence of methanol effected reductive cyclization<sup>11</sup> to deliver bicyclic ester **18** as the sole product in high yield. After protection as the *tert*-butyldimethylsilyl (TBS) ether, DIBALH reduction of the ester moiety to the aldehyde followed by Wittig reaction gave olefin **19** in 81% yield for the three steps. Hydroboration using 9-BBN followed by oxidative workup gave an alcohol (91%), which was then oxidized by a two-step procedure (SO<sub>3</sub>•pyr/DMSO then NaClO<sub>2</sub>) to provide acid **7** in 96% yield for the two steps.

With the desired fragments **6** and **7** in hand, we next turned our attention to their coupling. Yamaguchi esterification of **6a** with **7** proceeded smoothly to afford ester **20** in high yield (Scheme 4). Conversion to the  $\alpha$ -acetoxy ether **21** was carried out according to the procedure of Rychnovsky (DIBALH, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C; then Ac<sub>2</sub>O, DMAP, pyridine, -78 °C  $\rightarrow$  room temperature).<sup>12</sup> Subsequent treatment with TMSSPh in the presence of TMSOTf and 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP) generated, after desilylation, mixed thioacetal **22** in 85% overall yield from **20**. The  $\beta$ -alkoxyacrylate unit was attached to the tertiary alcohol by treatment with methyl propiolate and trimethylphosphine to provide **23** in modest yield (47%; 84% yield based on recovered **21**). However, radical cyclization of **23** (Bu<sub>3</sub>SnH, AIBN toluene, 80 °C) was sluggish and yielded a complex

(9) Grieco, P. A.; Gilman, S.; Nishizawa, M. J. Org. Chem. 1976, 41, 1485–1486.

mixture of the desired O-linked oxepane **24**,<sup>13</sup> its diastereomer **25**, and reduction product **26** in a ratio of ca. 1:4:3, along with recovered **23**. The outcome of this radical cyclization can be explained as follows (Figure 2). Severe



Figure 2. Possible rationale for radical cyclization.

steric repulsion between the  $C_{21}^{14}$  tertiary methyl group and the  $\beta$ -hydrogen on the acrylate unit in the transition state structure **A** resulted in a preference for the structure **B**, which leads to **25**. A similar result has been reported by Nakata and co-workers in the SmI<sub>2</sub>-mediated reductive cyclization.<sup>15</sup>

Accordingly, we next investigated an alternative route as summarized in Scheme 5. Alcohol **6b** and carboxylic acid **7** 

<sup>(13)</sup> Stereostructure of compound **24** was assigned by NOE between 21-Me and 26-H and the coupling constant,  $J_{25,26} = 3.5$  Hz. In a related O-linked oxepane ring system, a small coupling constant, J = 0 Hz, between H<sub>a</sub> and H<sub>b</sub> was observed. See: Sasaki, M.; Noguchi, T.; Tachibana, K. J. Org. Chem. **2002**, 67, 3301–3310.



(14) Numbering of carbon atoms of all compounds in this paper corresponds to that of gambieric acid A.

(15) Suzuki, K.; Matsukura, H.; Matsuo, G.; Koshino, H.; Nakata, T. Tetrahedron Lett. 2002, 43, 8653–8655.

<sup>(10)</sup> Nicolaou, K. C.; Hwang, C. K.; Marron, B. E.; DeFrees, S. A.; Couladouros, E. A.; Abe, Y.; Carroll, P. J.; Snyder, J. P. *J. Am. Chem. Soc.* **1990**, *112*, 3040–3054.

<sup>(11) (</sup>a) Hori, N.; Matsukura, H.; Matsuo, G.; Nakata, T. *Tetrahedron Lett.* **1999**, *40*, 2811–2814. (b) Hori, N.; Matsukura, H.; Nakata, T. *Org. Lett.* **1999**, *1*, 1099–1101. (c) Matsuo, G.; Hori, N.; Nakata, T. *Tetrahedron Lett.* **1999**, *40*, 8859–8862. (d) Hori, N.; Matsukura, H.; Matsuo, G.; Nakata, T. *Tetrahedron* **2002**, *58*, 1853–1864.

<sup>(12) (</sup>a) Dahanukar, V. H.; Rychnovsky, S. D. J. Org. Chem. **1996**, 61, 8317–8320. (b) Kopecky, D. J.; Rychnovsky, S. D. J. Org. Chem. **2000**, 65, 191–198. (c) Kopecky, D. J.; Rychnovsky, S. D. Org. Synth. **2003**, 80, 177–183.

Scheme 5. Synthesis of the CDEFG-Ring System 5



were coupled through esterification to afford 27 (92%), which was subjected to reductive acetylation to give an  $\alpha$ -acetoxy ether as a 3:1 mixture of diastereomers. Subsequent treatment with TMSCN in the presence of TMSOTf and DTBMP generated nitrile 28,<sup>16</sup> which was desilvlated to produce alcohol 29 in 68% yield over the three steps. The nitrile group was hydrolyzed under alkaline conditions to give carboxylic acid 30 as a 1:1 mixture of isomers. Yamaguchi lactonization produced a mixture of seven-membered lactones, which was separated by flash column chromatography to give 31a and 31b in 38 and 40% yield, respectively.<sup>17</sup> Reduction of lactone 31a with DIBALH followed by in situ acetylation produced acetate 32 in 93% yield as the sole product. Upon treatment of **32** with allyltrimethylsilane in the presence of  $BF_3 \cdot OEt_2$ (MeCN,  $-40 \rightarrow 0$  °C), stereoselective allylation occurred from the less hindered  $\alpha$ -face of the molecule to generate diene 33 in 67% yield. Stereochemistry at the C<sub>26</sub> position was determined by NOE between 21-Me and 26-H. Finally, construction of the nine-membered F-ring was achieved by

the action of the second-generation Grubbs' catalyst  $34^{18,19}$  to furnish the targeted CDEFG-ring system 5 in 98% yield. The stereochemistry of 5 was unambiguously established by NOE experiments and the coupling constant.

In summary, we have developed a synthetic route to the CDEFG-ring system **5** of gambieric acids through convergent union of the CD- and G-rings. Further studies toward the total synthesis of gambieric acids along these lines are in progress and will be reported in due course.

Acknowledgment. This work was financially supported by the Sumitomo Foundation and a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, Culture and Technology, Japan.

**Supporting Information Available:** Experimental procedures and characterization data for compounds **10**, **12**–**14**, **6b**, **16–19**, **7**, **27–33**, and **5**. This material is available free of charge via the Internet at http://pubs.acs.org.

## OL050760K

<sup>(16)</sup> Synthesis of  $\alpha$ -cyano ether in a related system; see: Oishi, T.; Watanabe, K.; Murata, M. *Tetrahedron Lett.* **2003**, *44*, 7315–7319.

<sup>(17)</sup> Attempted isomerization of the undesired **31b** to **31a** under various basic conditions was unsuccessful.

<sup>(18)</sup> Scholl, M.; Ding, S.; Lee, C. W.; Grubbs, R. H. Org. Lett. 1999, 1, 953–956.

<sup>(19)</sup> For a recent review, see: Deiters, A.; Martin, S. F. Chem. Rev. 2004, 104, 2199-2238.