## ORGANIC LETTERS

2004 Vol. 6, No. 13 2193–2196

## Cu- and Pd-Catalyzed Asymmetric One-Pot Tandem Addition—Cyclization Reaction of 2-(2',3'-Alkadienyl)- $\beta$ -keto Esters, Organic Halides, and Dibenzyl Azodicarboxylate: An Effective Protocol for the Enantioselective Synthesis of Pyrazolidine Derivatives

Shengming Ma,<sup>\*,†,‡</sup> Ning Jiao,<sup>†</sup> Zilong Zheng,<sup>†</sup> Zhichao Ma,<sup>‡</sup> Zhan Lu,<sup>‡</sup> Longwu Ye,<sup>‡</sup> Youqian Deng,<sup>‡</sup> and Guofei Chen<sup>‡</sup>

State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R. China, and Department of Chemistry, Zhejiang University, Hangzhou 310027, P. R. China

masm@mail.sioc.ac.cn

Received April 9, 2004





The development of new chemical processes for producing elaborate and important hetereocyclic structures in a rapid, atom-economic,<sup>1</sup> and efficient manner has become an important area of research in organic chemistry. One of the major challenges in these areas is the multicomponent process in which a given target molecule would be assembled from three or more simple and flexible building blocks in a practical one-pot operation.<sup>2,3</sup>

Pyrazolidines are an intensively studied family of heterocycles, which are important subunits in biologically active naturally occurring compounds.<sup>4</sup> This pyrazolidine structural

<sup>\*</sup> Fax: (+86)21-64167510.

<sup>&</sup>lt;sup>†</sup> Chinese Academy of Sciences.

<sup>&</sup>lt;sup>‡</sup> Zhejiang University.

<sup>(1) (</sup>a) Trost, B. M. Science **1991**, 254, 1471. (b) Trost, B. M. Angew. Chem., Int. Ed. Engl. **1995**, 34, 259.

<sup>(2) (</sup>a) Tietze, L. F. *Chem. Rev.* **1996**, *96*, 115. (b) Parsons, P. J.; Penkett, C. S.; Shell, A. J. *Chem. Rev.* **1996**, *96*, 195. (c) Bunce, R. A. *Tetrahedron* **1995**, *51*, 13103.

<sup>(3)</sup> For recent reviews, see: (a) Hulme, C.; Gore, V. Curr. Med. Chem. 2003, 10, 51. (b) Orru, R. V. A.; de Greef, M. Synthesis 2003, 1471. (c) Zhu, J. Eur. J. Org. Chem. 2003, 1133. (d) Balme, G.; Bossharth, E.; Monteiro, N. Eur. J. Org. Chem. 2003, 4101. (e) Bienaymé, H.; Hulme, C.; Oddon, G.; Schmitt, P. Chem. Eur. J. 2000, 6, 3321. (f) Dömling, A.; Ugi, I. Angew. Chem., Int. Ed. 2000, 39, 3168. (g) Tietze, L. F.; Modi, A. Med. Res. Rev. 2000, 20, 304. (h) Weber, L.; Illgen, K.; Almstetter, M. Synlett 1999, 366. (i) Dax, S. L.; Mcnally, J. J.; Youngman, M. A. Curr. Med. Chem. 1999, 6, 255.

unit is usually constructed by the [3 + 2] cycloaddition<sup>5,6</sup> or hetero-Diels–Alder reactions.<sup>7</sup> Although these methods are available, there are still some obvious limitations: the starting material is complex, and the methods for the synthesis of optically active pyrazolidines are still very limited.

Recently, the direct enantioselective amination of 2-ketoesters catalyzed by chiral copper(II)-bisoxazoline complexes was reported.<sup>8</sup> On the basis of our past work,<sup>9,10</sup> we reasoned that pyrazolidines **1** with different substitution patterns may be generally constructed from  $\pi$ -allypalladium intermediate<sup>11</sup> **2**, which in turn, could be generated easily via the carbopalladation of intermediate **3** with allene **4**<sup>12,13</sup> (Scheme 1).



The optically active allene **4** may be produced from enantioselective copper (II)-catalyzed  $\alpha$ -amination of  $\alpha$ -allenicsubstituted  $\beta$ -keto esters **5** with azodicarboxylates **6** (Scheme 1). Herein, we describe the Cu- and Pd-catalyzed one-pot

(4) (a) Rahman, M. T.; Nishino, H.; Qian, C.-Y. *Tetrahedron Lett.* **2003**, 44, 5225. (b) Chauveau, A.; Martens, T.; Bonin, M.; Micouin, L.; Husson, H.-P. *Synthesis* **2002**, 1885. (c) Hanessian, S.; Mcnaughton-Smish, G.; Lombart, H.-G. *Tetrahedron* **1997**, *53*, 12798. (d) Kim, H.-O.; Lum, C.; Lee, M. S. *Tetrahedron Lett.* **1997**, *38*, 4935.

(5) (a) For some reviews, see: (a) Huisgen, R.; Grashey, R.; Sauer, J. In *The Chemistry of Alkenes: Cycloaddition Reactions of Alkenes*; Patai, S., Ed.; Interscience: New York, 1964; pp 739–953. (b) Huisgen, R. *1,3-Dipolar Cycloaddition Chemistry: 1,3-Dipolar Cycloaddition-Introduction, Survey, Mechanism*; Padwa, A., Ed.; Wiley-Interscience: New York, 1984; Vol. 1, pp 1–176. (c) Huisgen, R. *Advance in Cycloaddition: Steric Course and Mechanism of 1,3-Dipolar Cycloaddition*; Curran, D. P., Ed.; JAI Press: Greenwich, CT, 1988; Vol. 1, pp 1–31.

(6) (a) Kobayashi, S.; Hirabayashi, R.; Shimizu, H.; Ishitani, H.; Yamashita, Y. *Tetrahedron Lett.* 2003, 44, 3351. (b) Kobayashi, S.; Shimizu, H.; Yamashita, Y.; Ishitani, H.; Kobayashi, J. J. Am. Chem. Soc. 2002, 124, 13678. (c) Jäger, V.; Bierer, L.; Dong, H.-Q.; Palmer, A. M.; Shaw, D.; Frey, W. J. Heterocycl. Chem. 2000, 37, 455. (d) Mish, M. R.; Guerra, F. M.; Carreira, E. M. J. Am. Chem. Soc. 1997, 119, 8379. (e) Gallos, J. K.; Koumbis, A. E.; Apostolakis, N. E. J. Chem. Soc., Perkin Trans. 1 1997, 2457. (f) Khau, V. V.; Martinelli, M. J. Tetrahedron Lett. 1996, 25, 4323.

three-component asymmetric tandem addition-cyclization leading to optically active pyrazolidine derivatives with high selectivity.

Our first approach was based on the reaction of 2-(2',3'dienyl)- $\beta$ -ketoesters **5a** and dibenzyl azodicarboxylate (DBAD) **6** with PhI (**7a**). First, the enantioselective amination of **5a** with DBAD **6** in CH<sub>2</sub>Cl<sub>2</sub> provided the corresponding optically active allenic amide **4a** (E<sup>1</sup> = COCH<sub>3</sub>; E<sup>2</sup> = COOEt; R<sup>1</sup> = R<sup>2</sup> = COOBn) in quantitative yield. Then, the solvent was removed in vacuo and THF was added to the reaction mixture followed by treatment with PhI, Pd(PPh<sub>3</sub>)<sub>4</sub> (5% mol), and K<sub>2</sub>CO<sub>3</sub> (2.0 equiv) under reflux (Scheme 2). The carbopalladation reaction<sup>14</sup> of the allenes afforded a **2**-type  $\pi$ -allyl palladium intermediate,<sup>11</sup> which can be further trapped by the intramolecular nitrogen giving the corresponding pyrazolidine derivative **1aa** in good overall yields with very high enantioselectivity (Scheme 2). On the



basis of these results, it is quite obvious that the enantioselectivity for the formation of **4a** is very high with the absolute configuration depending on the structure of the ligand.

Although the diastereoselectivity is unsatisfactory, the diastereoisomers differ greatly in terms of molecular polarity  $(R_{f(R,R)-1aa} = 0.48, R_{f(R,S)-1aa} = 0.35; \text{eluent, petroleum ether/} \text{ether} = 1:1)$  and therefore can be easily separated and purified by flash chromatography on silica gel. Then, the four pure isomers (S,S)-, (S,R)-, (R,R)-, and (R,S)-isomer of **1aa** can be conveniently obtained in high yield with very high ee when both enantiomers of Ph-Box were applied as the ligand.

<sup>(7)</sup> Ellis, J. M.; King, S. B. Tetrahedron Lett. 2002, 43, 5833.

<sup>(8) (</sup>a) Marigo, M.; Juhl, K.; Jørgensen, K. A. Angew. Chem., Int. Ed.
2003, 42, 1367. (b) Juhl, K.; Jørgensen, K. A. J. Am. Chem. Soc. 2002, 124, 2420. (c) Evans, D. A.; Nelson, S. G. J. Am. Chem. Soc. 1997, 119, 6452.

<sup>(9)</sup> Ma, S.; Jiao, N. Angew. Chem., Int. Ed. 2002, 41, 4737.

<sup>(10)</sup> For some of our most recent results, see: (a) Ma, S.; Duan, D.;
Shi, Z. Org. Lett. 2000, 2, 1419. (b) Ma, S.; Zhang, J. Chem. Commun.
2000, 117. (c) Ma, S.; Li, L. Org. Lett. 2000, 2, 941. (d) Ma, S.; Xie, H.
Org. Lett. 2000, 2, 3801. (e) Ma, S.; Zhao, S. J. Am. Chem. Soc. 1999, 121, 7943. (f) Ma, S.; Yu, F.; Gao, W. J. Org. Chem. 2003, 68, 5943.

<sup>(11)</sup> Shimizu, I.; Tsuji, J. *Chem. Lett.* **1984**, 233. Ahmar, M.; Cazes, B.; Gore, J. *Tetrahedron Lett.* **1984**, 25, 4505.

<sup>(12)</sup> Schuster, H. F.; Coppola, G. M. Allenes in Organic Synthesis; John Wiley & Sons: New York, 1984. Patai, S., Ed. The Chemistry of Ketenes, Allenes, and Related Compounds; John Wiley & Sons: New York, 1980; Part 1.

<sup>(13) (</sup>a) Ma, S.; Shi, Z. J. Org. Chem. 1998, 63, 6387. (b) Ma, S.; Zhao,
S. Org. Lett. 2000, 2, 2495. (c) Ma, S.; Jiao, N. Zhao, S.; Hou, H. J. Org.
Chem. 2002, 67, 2837. (d) Ma, S.; Gao, W. Org. Lett. 2002, 4, 2989.

<sup>(14)</sup> For intermolecular carbopalladation of allenes, see: (a) Cazes, B. *Pure Appl. Chem.* **1990**, *62*, 1867 and references therein. (b) Larock, R. C.; Berrios-Pena, N. G.; Fried, C. A. *J. Org. Chem.* **1991**, *56*, 2615.

**Table 1.** Cu and Pd-Catalyzed Three-Component Asymmetric Tandem Addition–Cyclization Reactions of 2-(2',3'-Dienyl)- $\beta$ -keto-esters **5a** and Dibenzyl Azodicarboxylate (DBAD) **6** with PhI **7a** in Different Solvents<sup>*a*</sup>



entry	Solvent (1)	<i>t</i> (1), C	time(1), n	Solvent (2)	$l(\omega), C$	time (2), n	yleiu (70)	ui (/0 ee) ( <i>I</i> ( <i>,I</i> ( <i>)</i> /( <i>I</i> ( <i>,S</i> ))
1	CH <sub>2</sub> Cl <sub>2</sub>	0	3	THF	reflux	4	92	36:64 (98;98)
2	THF	0-rt	57	THF	reflux	4	66	48:52 (99;96)
3	$CH_2Cl_2$	0	3	$CH_2Cl_2/A^b$	reflux	4	92	40:60 (98;98)
4	$CH_2Cl_2$	0	3	$CH_2Cl_2/B^b$	100	4	85	33:67 (98;98)
5	$CH_2Cl_2/B^b$	rt	3	$CH_2Cl_2/B^b$	100	4	95	32:68 (95;95)
6	CH <sub>2</sub> Cl <sub>2</sub>	rt	1.5	$CH_2Cl_2/B^b$	100	4	100	37:63 (97;97)
7	$CH_2Cl_2/B^b$	0	10	$CH_2Cl_2/B^b$	100	4	90	36:64 (97;98)
8	$CH_2Cl_2/C^b$	0	20	$CH_2Cl_2/C^b$	100	4	19	76:24 (13;8)
9	$CH_2Cl_2/D^b$	0	20	$CH_2Cl_2/D^b$	100	4	0	0

<sup>*a*</sup> DBAD (1.2 equiv), Ph-Box (1.1 equiv of Cu), PhI (1.2 equiv), and  $K_2CO_3$  (2.0 equiv) were applied. <sup>*b*</sup> A = THF; B = 1,4-dioxane; C = DMSO; D = DMF. The solvent ratio was 1:2.

To identify a more convenient operation for the reaction, we tested the effect of the solvents of the two steps on the reaction. The results are summarized in Table 1. When both steps were carried out in THF, high enantioselectivity was achieved; however, the yield and diastereoselectivity were unsatisfactory (entry 2, Table 1). When a mixture of CH<sub>2</sub>Cl<sub>2</sub>/ 1,4-dioxane (1:2) was used as the solvent in both steps, the results were better (entry 7, Table 1), but the first step was slow (compare entry 7 with entry 4, Table 1). The reaction gave poor results in the mixed solvent of DMSO or DMF and CH<sub>2</sub>Cl<sub>2</sub> (entries 8 and 9, Table 1). The temperature in the first step is important to the enantioselectivity (compare entry 4 with entry 6 and entry 5 with entry 7, Table 1). On the basis of these results, CH<sub>2</sub>Cl<sub>2</sub> was chosen as the solvent at 0 °C in the first step followed by the direct addition of 1,4-dioxane (double volume of CH<sub>2</sub>Cl<sub>2</sub>) to the reaction mixture to conduct the second step at 100 °C.

The results of the Cu- and Pd-catalyzed three-component asymmetric tandem addition—cyclization reactions of 2-(2',3'dienyl)- $\beta$ -ketoesters **5a** and dibenzyl azodicarboxylate (DBAD) **6** with different organohalides are summarized in Table 2. From these results, it should be noted that (1) the yield and the enantioselectivity of these reactions are high and (2) the configuration of the C=C bond in 1-alkenyl iodide remained unchanged during the reaction (entries 7–9, Table 2). To determine the absolute structure of the pyrazolidine derivative, we studied the reaction of **5a**, **6**, and 1-bromo-4iodobenzene **7e**, which afforded the solid pyrazolidine derivative **1ae** with a bromine atom in the structure, indicating that the C–I bond was chemoselectively cleaved. The absolute configurations of **1ae** were determined by X-ray diffraction studies,<sup>15,16</sup> which led to the tentative assignment

## **Table 2.** Cu- and Pd-Catalyzed Three-Component AsymmetricTandem Addition-Cyclization Reactions of

2-(2',3'-Dienyl)- $\beta$ -keto-esters **5a** and Dibenzyl Azodicarboxylate (DBAD) **6** with Different Organohalides



entry	RI	7	yield (%)	dr(R,R/ R,S)	ee% (R,R)	ee% (R,S)
1	PhI	7a	85 (1 <b>aa</b> )	33:67	98	98
2	I—	7b	95 (1 <b>a</b> b)	34:66	98	98
3	I—————————————————————————————————————	7c	92 (1ac)	33:67	98	98
4	ІСООМе	7d	96 (1 <b>ad</b> )	38:62	97	98
5	IBr	7 <b>e</b>	94 (1ae)	36:64	97	>98
6		7f	97 ( <b>1af</b> )	40:60	>98	99
7	COOMe	7g	98 (1ag)	45:55	97	98
8	,Ph I	7h	86 (1 <b>ah</b> )	37:63	99	99
9	/ <sup>C4H9</sup>	7i	75 (1 <b>ai</b> )	34:66	97	98
10	⟨ <sub>s</sub> ⟩∟,	7j	71 ( <b>1aj</b> )	36:64	91	99

of the absolute configurations of other products. It is obvious that the reaction favors the formation of *trans*-pyrazolidine derivative.

In the reaction, it is valuable to note the following: (1) The reaction that allows a great increase in complexity is atom economic. (2) Although two different metals were used to catalyze the reaction, the operation is convenient with a one-pot operation and the yield is very high. (3) Two diastereomers were easily separated with high ee.

In conclusion, we have demonstrated the Cu- and Pdcatalyzed asymmetric one-pot tandem addition-cyclization reaction of  $2-(2',3'-\text{dienyl})-\beta$ -keto esters, organic halides, and dibenzyl azodicarboxylate leading to pyrazolidine derivatives with high enantioselectivity. The advantages of this method are the easy availability and diversity of the starting compounds. Although the diastereoselectivity needs further attention, this study will open a new area for the transition metal-catalyzed chemistry of allenes. Further studies on the scope and synthetic applications of this reaction are being pursued in our laboratory.

Acknowledgment. Financial support from the Major State Basic Research Development Program (Grant G20000-77500), the National Natural Science Foundation of China, and Shanghai Municipal Committee of Science and Technology are greatly appreciated.

**Supporting Information Available:** Typical experimental procedure and analytical data for all products. This material is available free of charge via the Internet at http://pubs.acs.org.

OL0493498

<sup>(15)</sup> CCDC 232606. Crystal data for compound *cis*-**1ae** (3*R*,5*R*): C<sub>32</sub>-H<sub>31</sub>N<sub>2</sub>O<sub>7</sub>Br, MW = 635.50, crystal system, space group orthorhombic, *P*2(1)2(1)2(1), Mo K\alpha, final *R* indices  $[I > 2\sigma(I)]$ ,  $R_1 = 0.0469$ , w $R_2 = 0.0813$ , a = 8.9635 (9) Å, b = 10.2834 (11) Å, c = 32.786 (4) Å,  $\alpha = 90^{\circ}$ ,  $\beta = 90^{\circ}$ ,  $\gamma = 90^{\circ}$ , V = 3022.1 (6) Å<sup>3</sup>, T = 293 (2) K, Z = 4, reflections collected/unique: 18 496/7023 ( $R_{int} = 0.0975$ ), no observation  $[I > 2\sigma(I)]$  3265, parameters 425, absolute structure parameter: -0.001 (7).

<sup>(16)</sup> CCDC 232607. Crystal data for compound *trans*-**1ae** (3*R*,5*S*): C<sub>32</sub>H<sub>31</sub>N<sub>2</sub>O<sub>7</sub>Br, MW = 635.50, crystal system, space group orthorhombic, *P*2(1)2(1)2(1), Mo K $\alpha$ , final *R* indices  $[I > 2\sigma(I)]$ ,  $R_1 = 0.0629$ ,  $wR_2 = 0.1517$ , a = 9.0453 (7) Å, b = 9.5125 (8) Å, c = 35.549 (3) Å,  $\alpha = 90^{\circ}$ ,  $\beta = 90^{\circ}$ ,  $\gamma = 90^{\circ}$ , V = 3058.7 (4) Å<sup>3</sup>, T = 293 (2) K, Z = 4, reflections collected/unique: 18 653/7107 ( $R_{int} = 0.0710$ ), no observation [ $I > 2\sigma(I)$ ] 2796, parameters 357, absolute structure parameter: 0.014 (14).