

Synthesis of Swallowtail-Substituted Multiporphyrin Rods

Patchanita Thamyongkit and Jonathan S. Lindsey*

Department of Chemistry, North Carolina State University, Raleigh, North Carolina 27695-8204

jlindsey@ncsu.edu

Received April 17, 2004

Abstract: The availability of multiporphyrin arrays with defined architectures and good solubility in organic solvents is essential for a wide variety of physical studies. Herein the synthesis of linear multiporphyrin arrays (triads, tetrad, pentad) bearing solubilizing 7-tridecyl (swallowtail) groups is presented. The rodlike arrays are composed of zinc porphyrins at the termini and 1, 2, or 3 free base porphyrins at the core. The free base porphyrins in the tetrad and pentad are joined to each other via *p*-phenylene linkers whereas the zinc porphyrins via 1,4-diphenylethyne linkers. The arrays are designed for studies of interporphyrin electronic communication.

A wide variety of molecular constructs containing multiple porphyrins have been prepared to serve as lightharvesting arrays, optoelectronic gates, charge-separation units, and charge-storage reservoirs.¹⁻⁶ The design of molecular architectures with prescribed electronic properties (energy transfer, electronic switching, charge separation/storage) requires a firm understanding of the mechanisms of electronic communication between interacting constituents. We previously prepared a series of multiporphyrin arrays for studies of ground-state electronic communication. The method for examining groundstate electronic communication involved hole-hopping in partially oxidized arrays. The arrays examined include (1) dimers of zinc porphyrins,^{7,8} (2) linear or right-angle triads with zinc porphyrins at the termini of the linear array and a free base porphyrin at the core,⁹ and (3) starshaped pentads with four zinc porphyrins at the periphery and a free base porphyrin at the core.⁷ Despite the larger size of the pentads, hole-hopping occurred between

- (1) Harvey, P. D. In *The Porphyrin Handbook*; Kadish, K. M., Smith, K. M., Guilard, R., Eds.; Academic Press: San Diego, CA, 2003; Vol. 18, pp 63–250.
- (2) (a) Kim, D.; Osuka, A. J. Phys. Chem. A 2003, 107, 8791–8816.
 (b) Aratani, N.; Osuka, A. Macromol. Rapid Commun. 2001, 22, 725–740.
- (3) Holten, D.; Bocian, D. F.; Lindsey, J. S. Acc. Chem. Res. 2002, 35, 57–69.
- (4) Burrell, A. K.; Officer, D. L.; Plieger, P. G.; Reid, D. C. W. *Chem. Rev.* **2001**, *101*, 2751–2796.
- (5) Gust, D.; Moore, T. A.; Moore, A. L. Acc. Chem. Res. 2001, 34, 40-48.
- (6) Wasielewski, M. R. Chem. Rev. 1992, 92, 435-461.
- (7) Seth, J.; Palaniappan, V.; Johnson, T. E.; Prathapan, S.; Lindsey, J. S.; Bocian, D. F. *J. Am. Chem. Soc.* **1994**, *116*, 10578–10592.
- (8) (a) Strachan, J.-P.; Gentemann, S.; Seth, J.; Kalsbeck, W. A.;
 (8) (a) Strachan, J.-P.; Gentemann, S.; Seth, J.; Kalsbeck, W. A.;
 Lindsey, J. S.; Holten, D.; Bocian, D. F. *J. Am. Chem. Soc.* **1997**, *119*, 11191–11201. (b) Yang, S. I.; Lammi, R. K.; Seth, J.; Riggs, J. A.; Arai, T.; Kim, D.; Bocian, D. F.; Holten, D.; Lindsey, J. S. *J. Phys. Chem. B* **1998**, *102*, 9426–9436.

(9) Seth, J.; Palaniappan, V.; Wagner, R. W.; Johnson, T. E.; Lindsey, J. S.; Bocian, D. F. *J. Am. Chem. Soc.* **1996**, *118*, 11194–11207.

SCHEME 1



zinc porphyrins in a triad motif owing to the star-shaped architecture.

A chief problem encountered upon examination of the linear triads was the limited solubility of the compounds. Indeed, adequate solubility for electrochemical studies could only be achieved in mixtures of chlorinated solvents (e.g., CH₂Cl₂ and CHCl₃).⁹ The previous arrays incorporated porphyrins bearing mesityl groups at all nonlinking meso positions (Triad-1, Chart 1). We recently showed that 7-tridecyl ("swallowtail") groups impart solubility to meso-substituted porphyrin building blocks without affecting the physical properties of the macrocycles.¹⁰ Herein we present the synthesis of a series of rodlike arrays composed of multiple swallowtail-substituted porphyrins, including four triads, one tetrad, and one pentad. The triads enable in-depth studies of electronic communication while the tetrad and pentad enable examination of whether hole-hopping persists over longer distances than that in the triad arrays.

The four triads (**Triad-2**–**5**) are shown in Chart 1. **Triad-2** bears swallowtail substituents at nonlinking meso positions of the terminal porphyrins and mesityl substituents at nonlinking meso positions of the central free base porphyrin. **Triad-3** is a zinc-chelated derivative of **Triad-2**. **Triad-4** bears swallowtail groups at the terminal porphyrins while the free base porphyrin bears pentafluorophenyl groups to tune the energy levels of the porphyrin molecular orbitals. **Triad-5** bears swallowtail and mesityl groups in the reversed pattern of that in **Triad-2**. The tetrad or pentad each incorporates zinc porphyrins at the termini and two or three free base porphyrins as the intervening unit (Chart 2). The linker between the free base porphyrins is a *p*-phenylene unit,

10.1021/jo049348t CCC: \$27.50 © 2004 American Chemical Society Published on Web 07/28/2004

⁽¹⁰⁾ Thamyongkit, P.; Speckbacher, M.; Diers, J. R.; Kee, H. L.; Kirmaier, C.; Holten, D.; Bocian, D. F.; Lindsey, J. S. *J. Org. Chem.* **2004**, *69*, 3700–3710.

JOC Note

CHART 1







CHART 2





CHART 3



which affords enhanced electronic coupling between the porphyrins compared with that of the diphenylethyne unit.³ Each array is pseudocentrosymmetric along the long axis (ignoring torsional motions and NH tautomerism) and the two terminal porphyrins are identical with one another.

The general approach for synthesis of the triads entailed a Sonogashira reaction with a difunctional core porphyrin and 2 equiv of the monofunctional terminal porphyrin. Porphyrin building blocks **Zn-1**,¹⁰ **Zn-2**,¹⁰ **Zn-3**,¹¹ **4**,¹² and **Zn-4**¹² are known compounds (Chart 3). Condensation of dipyrromethane **5**¹³ and 4-(2-(trimeth-

J. Org. Chem, Vol. 69, No. 17, 2004 5797

JOC Note

SCHEME 2 (1) EtMgBr, THF (2) (3) DDQ (3) DD

SCHEME 3



11

ylsilyl)ethynyl)benzaldehyde with catalysis by TFA¹⁴ followed by oxidation with DDQ afforded porphyrin **6** in 16% yield. Subsequent treatment with TBAF in THF gave ethynyl porphyrin **7**, a free base precursor for the synthesis of **Triad-4**, in 75% yield (Scheme 1). Acylation of dipyrromethane **8**¹⁰ with pyridyl benzothioate **9**¹⁵ led to 1-acyldipyrromethane **10**. Reduction of **10** and self-

SCHEME 4



condensation¹⁵ of the resulting carbinol in CH_2Cl_2 containing Yb(OTf)₃¹⁶ followed by oxidation with DDQ gave porphyrin **11**, required for the synthesis of **Triad-5**, in 18% yield (Scheme 2). The condensation of a dipyrromethane-carbinol can be achieved with a mild Lewis acid such as Yb(OTf)₃ in $CH_2Cl_2^{15}$ or by trifluoroacetic acid in CH_3CN ,¹⁶ though the former conditions generally are superior in terms of diminished propensity to acidolysis of the dipyrromethane and greater ease of purification of the resulting porphyrin.

⁽¹¹⁾ Wagner, R. W.; Johnson, T. E.; Li, F.; Lindsey, J. S. J. Org. Chem. **1995**, 60, 5266–5273.

⁽¹²⁾ Li, J.; Ambroise, A.; Yang, S. I.; Diers, J. R.; Seth, J.; Wack, C. R.; Bocian, D. F.; Holten, D.; Lindsey, J. S. *J. Am. Chem. Soc.* **1999**, *121*, 8927–8940.

⁽¹³⁾ Laha, J. K.; Dhanalekshmi, S.; Taniguchi, M.; Ambroise, A.; Lindsey, J. S. *Org. Proc. Res. Dev.* **2003**, *7*, 799–812.

⁽¹⁴⁾ Littler, B. J.; Ciringh, Y.; Lindsey, J. S. *J. Org. Chem.* **1999**, *64*, 2864–2872.

⁽¹⁵⁾ Rao, P. D.; Littler, B. J.; Geier, G. R., III; Lindsey, J. S. *J. Org. Chem.* **2000**, *65*, 1084–1092.

⁽¹⁶⁾ Geier, G. R., III; Callinan, J. B.; Rao, P. D.; Lindsey, J. S. J. Porphyrins Phthalocyanines 2001, 5, 810-823.

)CNote

SCHEME 5



The synthesis of the triads is summarized in Scheme 3. The Sonogashira reaction was performed under copperfree conditions in toluene/TEA (5:1) at 35 °C in the presence of $Pd_2(dba)_3$ and $P(o-tol)_3$.¹⁷ In each case, the desired triad was isolated by a straightforward combination of adsorption chromatography and size exclusion chromatography. The yields ranged from 37% to 87%. Triad-2, Triad-3, and Triad-4, which bear swallowtail groups at the porphyrin termini, exhibit satisfactory solubility (>5 mg/mL) in common organic solvents (e.g., toluene, CH₂Cl₂, CHCl₃, THF) for routine handling, characterization, and physical studies. By contrast, Triad-5, which has swallowtail groups at the central porphyrin, was less soluble in the same solvents. In general, the swallowtail-substituted porphyrin arrays are more soluble than the corresponding mesityl-substituted porphyrin arrays (e.g., Triad-1).

The synthetic approach toward the target tetrad and pentad similarly entailed Sonogashira coupling of a core unit of *p*-phenylene-linked free base porphyrins and the terminal swallowtail-substituted zinc porphyrin. The reaction of 4-iodobenzaldehyde with excess pyrrole containing InCl₃¹³ afforded dipyrromethane **12** in 66% yield (Scheme 4). Diacylation of 12 was carried out with 3,5di-tert-butylbenzoyl chloride $(13)^{18}$ under catalysis by SbCl₅.¹⁹ Treatment of the mixture with dibutyltin dichloride¹⁰ and TEA gave the corresponding diacyldipyrromethane-tin complex 14 in 39% yield. Reduction of 14 with NaBH₄ in THF/methanol for 4 h gave the dipyrromethane-dicarbinol. Reaction of the latter with 1,4bis(dipyrromethyl)benzene (15)^{20,21} in CH₂Cl₂ containing

Yb(OTf)₃¹⁶ followed by oxidation with DDQ afforded the difunctional dimer **16** in 8% yield. Although the overall yield of **16** was low, the starting materials are readily available in sizable quantities.

The coupling reactions to form the tetrad and pentad were performed under the standard conditions for Pdmediated reactions of iodo and ethynyl porphyrins. The reaction of dimer 16 and porphyrin Zn-2 was complicated by the incomplete dissolution of the dimer (16) at the start of the reaction. Tetrad-1 was obtained in 19% yield (Scheme 4). The low yield is attributed to the low solubility of 16 and the formation of trimeric byproducts (as shown by analytical SEC). On the other hand, the Pd-mediated coupling of triad 17²² and porphyrin Zn-2¹⁰ afforded **Pentad-1** in 71% yield (Scheme 5). As with the triads, the tetrad and pentad were purified by straightforward combination of adsorption and size exclusion chromatography. Both Tetrad-1 and Pentad-1 exhibit satisfactory solubility (>5 mg/mL) in a variety of common organic solvents. Studies of electronic communication in the various rodlike arrays can now be undertaken and will be reported elsewhere.

Acknowledgment. This research was supported by a grant from the Division of Chemical Sciences, Office of Basic Energy Sciences, Office of Energy Research, U.S. Department of Energy.

Supporting Information Available: Full Experimental Section including characterization data (NMR spectra, mass spectra) for all new compounds, as well as analytical SEC traces for the arrays. This material is available free of charge via the Internet at http://pubs.acs.org.

JO049348T

(22) Yu, L.; Lindsey, J. S. Tetrahedron 2001, 57, 9285-9298.

⁽¹⁷⁾ Wagner, R. W.; Ciringh, Y.; Clausen, C.; Lindsey, J. S. Chem. Mater. 1999, 11, 2974-2983

^{(18) (}a) Korshunov, M. A.; Bodnaryuk, F. N.; Fershtut, E. V. Zh. (16) (a) Korshinov, M. A., Bonnaryu, F. N., Fershut, E. V. Zh.
 Org. Khim. **1967**, *3*, 140–143. (b) Alberti, A.; Benaglia, M. J. Organomet. Chem. **1992**, 434, 151–158.
 (19) Tamaru, S.-I.; Yu, L.; Youngblood, W. J.; Muthukumaran, K.; Taniguchi, M.; Lindsey, J. S. J. Org. Chem. **2004**, 69, 765–777.

⁽²⁰⁾ Lee, C.-H.; Lindsey, J. S. *Tetrahedron* 1994, *50*, 11427–11440.
(21) Bennis, V.; Gallagher, J. F. *Acta Crystallogr.* 1998, *C54*, 130– 132