This article was downloaded by: [North Dakota State University] On: 17 October 2014, At: 20:53 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lcar20

A New Approach to The Synthesis of Enantiomerically Pure 4-Deoxy Sugars

Romualdo Caputo, Annalisa Guaragna, Giovanni Palumbo, Silvana Pedatella & Francesco Solla

^a Dipartimento di Chimica Organica e Biologica , Università di Napoli Federico II Via Mezzocannone , 16 I-80134 Napoli, Italy E-mail:

Published online: 07 Mar 2008.

To cite this article: Romualdo Caputo , Annalisa Guaragna , Giovanni Palumbo , Silvana Pedatella & Francesco Solla (2000) A New Approach to The Synthesis of Enantiomerically Pure 4-Deoxy Sugars, Journal of Carbohydrate Chemistry, 19:4-5, 631-634, DOI: 10.1080/07328300008544104

To link to this article: <u>http://dx.doi.org/10.1080/07328300008544104</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/</u>page/terms-and-conditions

COMMUNICATION

A NEW APPROACH TO THE SYNTHESIS OF ENANTIOMERICALLY PURE 4-DEOXY SUGARS¹

Romualdo Caputo, Annalisa Guaragna, Giovanni Palumbo,* Silvana Pedatella, Francesco Solla

Dipartimento di Chimica Organica e Biologica, Università di Napoli Federico II Via Mezzocannone,16 I-80134 Napoli, Italy – e-Mail: ctsgroup@unina.it

Received September 17, 1999 - Final Form February 15, 2000

In a recent paper² we have reported the design and synthesis of 3-C-lithiated 5,6dihydro-1,4-dithiin-2-yl[(4-methoxybenzyl)oxy]methane (1) which can be utilized as an allylic alcohol anion equivalent and leads to three-carbon elongations of various electrophiles by introduction of a fully protected hydroxypropenyl moiety. The latter contains a double bond, which can be unravelled to the *cis* configuration by diastereoselective removal³ of the dimethylene-disulfur bridge, as well as a protected primary hydroxyl group that, depending on the deprotection conditions used (DDQ/NaBH₄ or DDQ), may either lead to the free allylic alcohol or to an α , β unsaturated aldehyde.



Scheme 1. Coupling reaction with (R)-benzyl glycidyl ether.

CAPUTO ET AL.

We report now a new versatile synthesis of a precursor of 4-deoxy-L-sugars that is in fact initiated by coupling of 1 with (R)-benzyl glycidyl ether.

The coupling product 2 derived from (R)-benzyl glycidyl ether, after acetylation of the free hydroxyl group to give 3, was selectively deprotected *via* oxidative removal of the *p*-methoxybenzyl ether by DDQ to afford carbaldehyde 4 as shown in Scheme 2. Cyclization of carbaldehyde by treatment with TMSOTf and TEA in methanol led to a diastereomeric mixture of dihydropyrans 5 and 6. The latter was then desulfurised by treatment with Raney-Ni (W2) in glacial acetic acid to give the final compound 7. This is a formal 2,3-dehydro-2,3,4-trideoxy L-sugar which can be hydroxylated in a stereocontrolled manner⁴ to give 4-deoxy-L-sugars.



(i) DDQ in CH₂Cl₂/H₂O; (ii) TMSOTf, NEt₃ in MeOH; (iii) Raney-Ni in glacial acetic acid
Scheme 2. Formal synthesis of 4-deoxy-L-sugars.

The same procedure reported in Schemes 1 and 2 applied to (S)-benzyl glycidyl ether led, as expected, to the formation of the D analog of 7. Work is now in progress to achieve 4-deoxy L- and D-sugars by stereocontrolled hydroxylations of the double bond in 7 and its D analog.

EXPERIMENTAL

General methods. ¹H NMR spectra were recorded in CDCl₃ solutions: chemical shifts were reported in ppm (δ) downfield from internal tetramethylsilane (TMS). Optical rotations were measured in CHCl₃ solutions (1.0 dm cell). Thin-layer chromatography (TLC) analyses were performed on silica gel Merck 60 F₂₅₄ plates (0.2 mm layer thickness). Column chromatography was carried out with Merck Kieselgel 60 (70-230 mesh).

(5*R*,7*R*)- and (5*S*,7*R*)-[(Benzyloxy)methyl]-5-methoxy-3,5,7,8-tetrahydro-2H-[1,4]dithiino[2,3-c]pyran (5 and 6). To a stirred solution of aldehyde 4² (1.0 g; 2.8 mmol) in methanol (5 mL) at room temperature, TEA (1.95 mL; 14.0 mmol) and TMSOTf (2.2 mL; 14.0 mmol) were added slowly over 1 h. After 2 h, most of the solvent was evaporated under reduced pressure and replaced by AcOEt (10 mL). The organic phase was washed with brine until neutral, then dried (Na₂SO₄) and concentrated under reduced pressure. Chromatography of the crude residue on silica gel (hexane-AcOEt, 8:2) afforded the pure diastereomeric dihydropyrans 5 and 6 (0.83 g, 90% yield; diastereomer ratio 1.6:8.4). Higher R_f compound 5: (0.14 g), oil; $[\alpha]_D^{25}$ +18 (*c* 0.9); ¹H NMR (200 MHz) δ 1.98 (dd, 1H, J_{8a,7} = 3.3Hz, J_{8a,8b} = 16.6 Hz, H-8a), 2.44 (dd, 1H, J_{8b,7} = 11.4 Hz, J_{8b,8a} = 16.6 Hz, H-8b), 3.00-3.38 (m, 2H, H-2), 3.18-3.31 (m, 2H, H-3), 3.54 (dd, 1H, J_{9a,7} = 4.2 Hz, J_{9a,9b} = 10.3 Hz, H-9a), 3.57 (dd, 1H, J_{9b,7} = 5.1 Hz, J_{9b,9a} = 10.3 Hz, H-9b), 3.83 (s, 3H, OCH₃), 4.26-4.37 (m, 1H, H-7), 4.57 (d, 1H, J_{Ha,Hb} = 12.1 Hz, Ha_{Bn}), 4.61 (d, 1H, J_{Hb,Ha} = 12.1 Hz, Hb_{Bn}), 4.85 (s, 1H, H-5), 7.35-7.39 (m, 5H, H_{Ar}).

Anal. Calcd for $C_{16}H_{20}O_{3}S_{2}$ (324.46): C, 59.23; H, 6.21. Found C, 59.58; H, 6.17. Lower R_{f} compound 6: (0.69 g), oil; $[\alpha]_{D}^{25}$ +39 (*c* 1.4); ¹H NMR (400 MHz) δ 1.93 (dd, 1H, $J_{8a,7} = 3.4$ Hz, $J_{8a,8b} = 16.5$ Hz, H-8a), 2.39 (dd, 1H, $J_{8b,7} = 11.6$ Hz, $J_{8b,8a} = 16.5$ Hz, H-8b), 3.05-3.15 (m, 2H, H-2), 3.18-3.31 (m, 2H, H-3), 3.43 (s, 3H, OCH₃), 3.53 (dd, 1H, $J_{9a,7} = 4.6$ Hz, $J_{9a,9b} = 10.6$ Hz, H-9a), 3.58 (dd, 1H, $J_{9b,7} = 5.2$ Hz, $J_{9b,9a} = 10.6$ Hz, H-9b), 4.23-4.32 (m, 1H, H-7), 4.57 (d, $J_{Ha,Hb} = 12.1$ Hz, 1H, Ha_{Bn}), 4.61 (d, 1H, $J_{Hb,Ha} =$ 12.1 Hz, Hb_{Bn}), 4.77 (s, 1H, H-5), 7.31-7.38 (m, 5H, H_{Ar}); ¹³C NMR (400 MHz) ppm 27.8, 28.9 (C-2, C-3), 33.8 (C-8), 55.8 (CH₃O), 67.3 (C-9), 72.4 (C-Bn), 74.0 (C-7), 99.1 (C-5), 110.0 (C-4a and C-8a).

Anal. Calcd for C₁₆H₂₀O₃S₂ (324.46): C, 59.23; H, 6.21. Found C, 59.49; H 6.26.

(2S,6R)-2-Methoxy-6-[(benzyloxy)methyl]-3,6-dihydro-2H-pyran (7). A

solution of dihydropyran 6 (0.5 g; 1.5 mmol) in glacial acetic acid (10 mL) was added in one portion to a stirred suspension of Raney-Ni (W2) (5.5 g, wet) in the same solvent (10 mL) at 0° C and under argon stream. The suspension was stirred for 2 min (TLC monitoring). Then the solid was filtered off and washed with glacial acetic acid, water, and AcOEt. The filtrate was neutralized with saturated aq Na₂CO₃ and extracted with AcOEt. The combined organic layers were washed with water until neutral, dried (Na₂SO₄), and concentrated under reduced pressure to afford a crude residue. Chromatography of the latter on silica gel (CH₂Cl₂) gave pure 7 (0.26 g, 70% yield). ¹H NMR (200 MHz) δ 2.05-2.35 (m, 2H, H-5), 3.48 (s, 3H, OCH₃), 3.50-3.65 (m, 2H, H-7), 4.10-4.31 (m, 1H, H-6), 4.57 (d, 1H, J_{Ha,Hb} = 12.0 Hz, Ha_{Bn}), 4.63 (d, 1H, J_{Hb,Ha} = 12.0 Hz, Hb_{Bn}), 5.02 (d, 1H, J_{6.5} = 6.5 Hz, H-2), 5.66 (dd, 1H, J_{5.6} = 6.5 Hz, J_{5.4} = 9.1 Hz, H-3), 5.92-6.15 (m, 1H, H-4), 7.15-7.32 (m, 5H, H_{Ar}).

Anal. Calcd for C₁₄H₁₈O₃ (234.30): C, 71.77; H, 7.74. Found: C, 71.56; H, 7.80.

ACKNOWLEDGMENTS

Financial support by Regione Campania (Ricerca Scientifica) to G.P. is gratefully acknowledged.

REFERENCES AND NOTES

- 1. Presented as a poster at the First Euroconference on Carbohydrates in Drug Research, Sardinia, September 16-19, 1999.
- 2. R. Caputo, A. Guaragna, G. Palumbo and S. Pedatella, J. Org. Chem., 62, 9369 (1997).
- 3. R. Caputo, G. Palumbo and S. Pedatella, *Tetrahedron*, 50, 7265 (1994).
- a) H.C. Kolb, M.S. VanNieuwenhze and K.B. Sharpless, *Chem. Rev.*, 94, 2483 (1994); b) M.S. VanNieuwenhze and K.B. Sharpless, *Tetrahedron Lett.*, 35, 843 (1994); c) C.Y. Park and K.B. Sharpless, *Tetrahedron Lett.*, 35, 2495 (1994).