

Magnesium perchlorate as efficient Lewis acid for the Knoevenagel condensation between β -diketones and aldehydes

Giuseppe Bartoli, Marcella Bosco, Armando Carlone, Renato Dalpozzo,
Patrizia Galzerano, Paolo Melchiorre, Letizia Sambri*

Università di Bologna, Dipartimento di Chimica Organica 'A. Mangini', v.le Risorgimento 4, 40136 Bologna, Italy

Received 17 January 2008; revised 15 February 2008; accepted 18 February 2008

Available online 21 February 2008

Abstract

A new protocol for the Knoevenagel condensation between β -diketones and aliphatic and aromatic aldehydes promoted by $\text{Mg}(\text{ClO}_4)_2$ under mild conditions is reported.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: Knoevenagel condensation; Magnesium perchlorate; β -Diketones; Aldehydes; Mild reaction conditions

The Knoevenagel condensation is a well-known organic reaction largely employed in order to form C–C bonds for the synthesis of important derivatives in perfume, polymer, and pharmaceutical applications.¹ The Knoevenagel adducts, in fact, are useful intermediates for further transformations, such as Diels–Alder and Michael additions.²


The classical Knoevenagel condensation (Scheme 1) has been carried out by reacting a methylene active compound with an aldehyde or a ketone in the presence of a base.^{1,3} Recently, alternative procedures employed heterogeneous catalysts, such as zeolites,⁴ clays,⁵ layered double hydroxides (LDHs),⁶ and hydrotalcites.⁷ Various protocols carried out in ionic liquids⁸ have also been developed. On

the contrary, relatively few examples of condensation mediated by Lewis acids such as TiCl_4 ,⁹ ZnCl_2 ,¹⁰ $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ / NaI ,¹¹ and HClO_4 – SiO_2 ¹² have been reported. Frequently, high reaction temperatures^{4a,5,12,13} or microwave irradiation^{12,14} are necessary to promote the reaction independently from the employed catalytic system.

Methylene active compounds carrying two electron-withdrawing groups, such as malononitrile, cyanoacetates, malonates, and β -ketoesters, are generally used in the known condensations. However, only few examples of β -diketones as starting materials are reported.^{12,15} Very likely, such compounds are less reactive than the other ones since their attitude to form a stable cyclic enol.

Since our interest in the last years in the use of metal perchlorates as Lewis acids,¹⁶ and since their ability to coordinate 1,3-bidentate substrates,¹⁷ in order to extend their applicability we investigated the possibility to promote a Knoevenagel condensation between β -diketones with aldehydes in the presence of a perchlorate salt.

On the other hand, perchlorate salts of alkaline and alkaline earth metals showed to be very active in promoting a large variety of reactions,¹⁸ in some cases being more powerful than metal triflates.^{16b,17c,19} Concerning the risk connected with the use of such salts, it has been demonstrated that such perchlorates are not dangerous chemicals



Scheme 1.

* Corresponding author. Tel.: +39 0512093624; fax: +39 0512093654.
E-mail address: letizia.sambri@unibo.it (L. Sambri).

if not employed under highly acidic conditions and not exposed to high temperatures (>300–500 °C).²⁰


To test the reactivity of β -diketones we chose the reaction of acetoacetone **4a** with benzaldehyde **5a** in the presence of various catalytic systems, under solvent free conditions at room temperature. We checked the conversion by NMR at maximum after 70 h, (Table 1).

The best reaction conditions required the use of a 10 mol % of $\text{Mg}(\text{ClO}_4)_2$, and a 20 mol % of MgSO_4 (Table 1, entry 1) to obtain a 65% of conversion in the desired product without any traces of by-products. On the other hand, an increase in temperature (40 °C) led to the formation of the side product **7aa**, derived from the attack of acetoacetone on the Knoevenagel adduct, right after 30 h (Table 1, entry 2). The addition of MgSO_4 as a catalyst was necessary very likely to adsorb the water formed during the condensation. In fact the reaction carried out only in the presence of $\text{Mg}(\text{ClO}_4)_2$ was slower (55% vs 65% of conversion, Table 1, entries 1 and 3). As in other reactions,^{16a,17d} $\text{Mg}(\text{ClO}_4)_2$ showed to be more effective than $\text{Zn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ (Table 1, entries 1 and 4). The attempt to increase the conversion by the addition of a catalytic amount of Et_3N was unsuccessful: after 48 h the conversion in the desired product was 55% but a 5% of the side product **7aa** was also detected (Table 1, entry 5).

Notably, the reaction works well under neat conditions, the addition of a solvent slows the conversion rate.

The protocol was applied to other substrates.²¹ Results are reported in Table 2. Acetoacetone **4a** smoothly reacted with both aromatic and aliphatic aldehydes **5a–h** (Table 2, entries 1–8). The presence of an electron-withdrawing group on the aromatic aldehyde, such as NO_2 and CN , increases the reactivity and very good yields in the desired product **6** were obtained (Table 2, entries 2–4).

Table 1
Reaction of acetoacetone **4a** with benzaldehyde **5a** (1.2 equiv) under various conditions


			
Entry	Catalyst	Time (h)	Conversion ^a in 6aa (7aa)
1	$\text{Mg}(\text{ClO}_4)_2$ 10 mol % MgSO_4 20 mol %	70	65 (0)
2	$\text{Mg}(\text{ClO}_4)_2$ 10 mol % MgSO_4 20 mol %	30 ^b	65 (3)
3	$\text{Mg}(\text{ClO}_4)_2$ 10 mol %	70	55 (0)
4	$\text{Zn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ 10 mol % MgSO_4 20 mol %	70	52 (0)
5	$\text{Mg}(\text{ClO}_4)_2$ 10 mol % MgSO_4 20 mol % Et_3N 10 mol %	48	55 (5)

^a Calculated from ^1H NMR.

^b Reaction carried out at 40 °C.

Table 2

Reaction of dicarbonyl compounds **4** with aldehydes **5** (1.2 equiv) in the presence of $\text{Mg}(\text{ClO}_4)_2$ (10 mol %) and MgSO_4 (20 mol %) at rt

						
Entry	R ¹	R ²	R ³	Time (h)	Yields ^a (%) (E/Z ratio)	Product
1	Me	Me	C_6H_4	70	55	6aa
2	Me	Me	$p\text{-NO}_2\text{-C}_6\text{H}_4$	70	86 ^b	6ab
3	Me	Me	$m\text{-NO}_2\text{-C}_6\text{H}_4$	48	78 ^b	6ac
4	Me	Me	$p\text{-CN-C}_6\text{H}_4$	24	70 ^b	6ad
5	Me	Me	$p\text{-Br-C}_6\text{H}_4$	70	65	6ae
6	Me	Me	$n\text{-C}_3\text{H}_7$	48	70	6af
7	Me	Me	$n\text{-C}_5\text{H}_{11}$	48	67	6ag
8	Me	Me	$\text{Ph}(\text{CH}_2)_2$	90	73 ^c	6ah
9	Et	Et	$p\text{-NO}_2\text{-C}_6\text{H}_4$	70	62 ^b	6bb
10	Et	Et	$p\text{-CN-C}_6\text{H}_4$	90	80 ^b	6bd
11	Et	Et	5- NO_2 -2-furyl	24	81 ^b	6bi
12	Et	Et	$\text{Ph-C}\equiv\text{C}$	24	63	6bj
13	<i>i</i> -Bu	Me	$p\text{-NO}_2\text{-C}_6\text{H}_4$	90	75 (1:7)	6cb
14	OEt	Me	$p\text{-NO}_2\text{-C}_6\text{H}_4$	54	88 ^b (1:3)	6db
15	OEt	Me	$p\text{-CN-C}_6\text{H}_4$	70	76 ^b (1:2.5)	6dd
16	Me	Me	trans-PhCH=CH	90	30	6ak

^a Isolated yields.

^b Reaction carried out in THF (3.5 M).

^c A 5% of the isomer **8ah** was also detected.

The reaction with aliphatic aldehydes is generally faster than those with aromatic ones. In particular, despite the presence of α -protons in the aliphatic aldehydes only the α,β -unsaturated systems **6** are obtained and no isomerization to β,γ -unsaturated²² ones was detected, (Table 2, entries 6 and 7). Only the reaction of 3-phenylpropanal **5h** required longer reaction times, and traces of by-product **8ah** (Fig. 1), an isomer of the desired product **6ah**, was observed.

Increasing the bulkiness of the diketone system ($\text{R}^1 = \text{R}^2 = \text{Et}$, **4b**), the reactivity essentially did not change (Table 2, entries 9–12) and the expected products were obtained in good yields. The reaction of **4b** with the substituted heteroaromatic aldehyde **5i** was faster than those with an aromatic one giving **6bi** in very good yield. Moreover, the reaction of **4b** with an unsaturated aldehyde like **5j** gave only **6bj** as product, and no 1,4-addition product was detected. On the other hand, an asymmetric and bulkier system like 6-methylheptane-2,4-dione **4c** took longer time to give **6cb** in good yields even if two isomeric products are formed, the Z-isomer being predominant, (Table 2, entry 13).

The protocol works also with active systems generally used in classical Knoevenagel condensation different from diketones. Ethyl 3-oxobutanoate **4d** reacted with aromatic




Fig. 1.

aldehydes to give a mixture of *Z*- and *E*-isomeric alkenes, (Table 2, entries 14 and 15).

We tried to extend the procedure to α,β -unsaturated aldehydes, but unfortunately not very satisfactory results were obtained, (Table 2, entries 16). In fact in the reaction of **4a** with cinnamaldehyde the desired product **6ak** was obtained in 30% yields, together with a mixture of starting materials and unidentified by-products.

In some cases, when the aldehyde is solid and does not dissolve in the diketone, the addition of a minimum amount of solvent is necessary to solubilize the reagents. THF proved to be the best choice, and the reactions were carried out under concentrated conditions (3.5 M).

In conclusion, we have demonstrated another interesting application of $\text{Mg}(\text{ClO}_4)_2$ to act as a Lewis acid in promoting the synthesis of trisubstituted functionalized alkenes via a Knoevenagel condensation between poorly reactive β -diketones and aliphatic and aromatic aldehydes. The protocol works also with more reactive systems like β -ketoesters. The reaction conditions are very mild, in fact the condensation works at room temperature, in the absence or with a minimum amount of solvent. Notably, this procedure led only to the Knoevenagel products, any side-product derived from a subsequent Michael addition of the β -diketone was never detected. In addition, the reaction with aliphatic aldehydes gave generally only the desired product, in contrast to other reported examples.

Acknowledgments

This research was carried out within the framework of the National Project 'Stereoselezione in Sintesi Organica' supported by MIUR, Rome, and FIRB National Project 'Progettazione, preparazione e valutazione farmacologica di nuove molecole organiche quali potenziali farmaci innovativi'.

References and notes

- (a) Jones, G. *Org. React.* **1967**, *15*, 204–599; (b) Tietze, L. F.; Beifuss, U. In *Comprehensive Organic Synthesis*; Pergamon: New York, 1991; Vol. 2, pp 341–394.
- Tietze, L. F. *Chem. Rev.* **1996**, *96*, 115–136.
- Knoevenagel, E. *Berichte* **1898**, *31*, 2585–2595.
- (a) Saravanamurugan, S.; Palanichamy, M.; Hartmann, M.; Murugesan, V. *Appl. Catal. A* **2006**, *298*, 8–15; (b) Reddy, T. I.; Varma, R. S. *Tetrahedron Lett.* **1997**, *38*, 1721–1724.
- Bigi, F.; Chesini, L.; Maggi, R.; Sartori, G. *J. Org. Chem.* **1999**, *64*, 1033–1035.
- Lakshmi Kantam, M.; Ravindra, A.; Venkat Reddy, Ch.; Sreedhar, B.; Choudary, B. M. *Adv. Synth. Catal.* **2006**, *348*, 569–578.
- Lakshmi Kantam, M.; Choudary, B. M.; Venkat Reddy, Ch.; Koteswara Rao, K.; Figueras, F. *Chem. Commun.* **1998**, 1033–1034.
- (a) Ranu, B. C.; Jana, R. *Eur. J. Org. Chem.* **2006**, 3767–3770; (b) Hangarge, R. V.; Jarikote, D. V.; Shingare, M. S. *Green Chem.* **2002**, *4*, 266–268; (c) Harjani, J. R.; Nara, S. J.; Salunkhe, M. M. *Tetrahedron Lett.* **2002**, *43*, 1127–1130.
- (a) Lehnert, W. *Tetrahedron* **1974**, *30*, 301–305; (b) Green, B.; Crane, R. I.; Khaideim, I. S.; Leighton, R. S.; Newaz, S. S.; Smyser, T. E. *J. Org. Chem.* **1985**, *50*, 640–644.
- Shanthan Rao, P.; Venkataratnam, R. V. *Tetrahedron Lett.* **1991**, *32*, 5821–5822.
- Bartoli, G.; Beleggia, R.; Giuli, S.; Giuliani, A.; Marcantoni, E.; Massaccesi, M.; Paletti, M. *Tetrahedron Lett.* **2006**, *47*, 6501–6504.
- Kantevari, S.; Bantu, R.; Nagarapu, L. *J. Mol. Catal. A: Chem.* **2007**, *269*, 53–57.
- (a) Bigi, F.; Conforti, M. L.; Maggi, R.; Piccino, A.; Sartori, G. *Green Chem.* **2000**, *2*, 101–103; (b) Hangarge, R. V.; Sonwane, S. A.; Jarikote, D. V.; Shingare, M. S. *Green Chem.* **2001**, *3*, 310–312; (c) Kaupp, G.; Naimi-Jamal, M. R.; Schmeyers, J. *Tetrahedron* **2003**, *59*, 3753–3760.
- Yadav, J. S.; Subba Reddy, B. V.; Basak, A. K.; Visali, B.; Narsaiah, A. V.; Nagaiah, K. *Eur. J. Org. Chem.* **2004**, 546–551.
- Su, C.; Chen, Z.-C.; Zheng, Q.-G. *Synthesis* **2003**, 555–559.
- (a) Bartoli, G.; Babiuch, K.; Bosco, M.; Carlone, A.; Galzerano, P.; Melchiorre, P.; Sambri, L. *Synlett* **2007**, 2897–2901. and references herein cited.; (b) Bartoli, G.; Bosco, M.; Carlone, A.; Dalpozzo, R.; Locatelli, M.; Melchiorre, P.; Sambri, L. *J. Org. Chem.* **2006**, *71*, 9580–9588.
- (a) Chakraborti, A. K.; Sharma, L.; Gulhane, R.; Shivani *Tetrahedron* **2003**, *59*, 7661–7668; (b) Bartoli, G.; Bosco, M.; Locatelli, M.; Marcantoni, E.; Melchiorre, P.; Sambri, L. *Synlett* **2004**, 239–242; (c) Bartoli, G.; Bosco, M.; Dalpozzo, R.; Marcantoni, E.; Massaccesi, M.; Sambri, L. *Eur. J. Org. Chem.* **2003**, 4611–4617; (d) Bartoli, G.; Bosco, M.; Locatelli, M.; Marcantoni, E.; Melchiorre, P.; Sambri, L. *Org. Lett.* **2005**, 427–430.
- Bartoli, G.; Locatelli, M.; Melchiorre, P.; Sambri, L. *Eur. J. Org. Chem.* **2007**, 2037–2049. and references herein cited.
- (a) Shivani; Pujala, B.; Chakraborti, A. K. *J. Org. Chem.* **2007**, *72*, 3713–3722; (b) Bhagat, S.; Chakraborti, A. K. *J. Org. Chem.* **2007**, *72*, 1263–1270.
- (a) Long, J. *Chem. Health Safety* **2002**, *9*, 12–18; (b) El-Awad, A. M.; Gabr, R. M.; Girgis, M. M. *Thermochim. Acta* **1991**, *184*, 205–212.
- General procedure for the Knoevenagel condensation:** In a two necked flask equipped with a magnetic stirring bar, $\text{Mg}(\text{ClO}_4)_2$ (0.10 mmol), MgSO_4 (0.20 mmol), β -keto derivative **4** (1.0 mmol) and aldehyde **5** (1.2 mmol) were added. If the system was completely solid, THF (0.29 ml) was added to dissolve the reagents. The mixture was stirred at rt. CH_2Cl_2 was added to the crude reaction mixture. Filtration on Celite and removal of the solvent by rotary evaporation gave the crude product. The substituted olefin **6** was purified by flash chromatography on silica gel with the appropriate mixture of hexane/ Et_2O .
Spectroscopic data for selected compounds follow: **3-Butyridenepentane-2,4-dione (6af)**: ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) = 0.90 (t, $J_{\text{HH}} = 7.4$, 3H), 1.42–1.51 (m, 2H), 2.11–2.18 (m, 2H), 2.25 (s, 3H), 2.26 (s, 3H), 6.63 (t, $J = 7.69$ Hz, 1H). ^{13}C NMR (CDCl_3 , 100 MHz): δ (ppm) = 13.7 (CH_3), 21.8 (CH_2), 25.9 (CH_3), 31.4 (CH_2), 31.6 (Me), 145.2 (CH), 146.6 (C), 197.1 (C), 203.5 (C). MS (ESI+): $m/z = 177$.
4-((5-Nitrofur-2-yl)methylene)heptane-3,5-dione (6bi): ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) = 1.11 (t, $J_{\text{HH}} = 7.1$, 3H), 1.22 (t, $J_{\text{HH}} = 7.1$, 3H), 2.67–2.75 (m, 4H), 6.82 (d, $J_{\text{HH}} = 3.8$, 1H), 6.15 (s, 1H), 7.28 (d, $J_{\text{HH}} = 3.8$, 1H). ^{13}C NMR (CDCl_3 , 100 MHz): δ (ppm) = 7.1 (CH_3), 7.7 (CH_3), 32.0 (CH_2), 37.1 (CH_2), 112.7 (CH), 117.7 (CH), 121.5 (CH), 143.4 (C), 150.5 (C), 197.8 (C), 205.9 (C). MS (ESI+): $m/z = 274$.
4-(3-Phenylprop-2-ynylidene)heptane-3,5-dione (6bj): ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) = 1.12 (t, $J_{\text{HH}} = 7.2$, 3H), 1.20 (t, $J_{\text{HH}} = 7.2$, 3H), 2.67 (q, $J_{\text{HH}} = 7.3$, 2H), 2.88 (q, $J_{\text{HH}} = 7.3$, 1H), 6.89 (s, 1H), 7.34–7.40 (3H, m), 7.43–7.46 (2H, m). ^{13}C NMR (CDCl_3 , 100 MHz): δ (ppm) = 7.6 (CH_3), 7.9 (CH_3), 32.6 (CH_2), 36.7 (CH_2), 85.0 (C), 105.4 (C), 120.2 (CH), 121.6 (C), 128.6 (CH), 129.9 (CH), 132.0 (CH), 149.8 (C), 198.1 (C), 204.8 (C). MS (ESI+): $m/z = 263$.
- (a) Wilson, B. D. *J. Org. Chem.* **1963**, *28*, 314–320; (b) Hiramatsu, H.; Harada, K.; Kojima, Y.; Fujiwara, K. *Nippon Kagaku Kaishi* **1989**, *4*, 714–721.