

Total Synthesis of Plagiochin D, A Macrocyclic Bis(bibenzyl) from Liverworts by Intramolecular Still-Kelly Reaction

Yoshiyasu Fukuyama,* Hideyuki Yaso, Kazuhiko Nakamura and Mitsuaki Kodama


Institute of Pharmacognosy, Faculty of Pharmaceutical Sciences, Tokushima Bunri University,
Yamashiro-cho, Tokushima 770, Japan

Received 17 September 1998; revised 21 October 1998; accepted 23 October 1998

Abstract: Plagiochin D (**4**), a unique macrocyclic bis(bibenzyl) having both a biphenyl ether and a biaryl units isolated from the liverwort *Plagiochila acanthophylla*, was synthesized. The key 16-membered ring closure of a dibromoperrottetin A derivative **13** was realized by Pd(0) catalyzed intramolecular Still-Kelly reaction.


© 1998 Elsevier Science Ltd. All rights reserved.

Macrocyclic Bis(bibenzyls) are plant metabolites occurring exclusively in liverworts, and over 30 congeners have been isolated so far.¹ Their unique structures and diversity of their biological activities have attracted much attention.^{1,2} They are categorized into two structural types 1 and 2, which are made up of macrocyclic rings linked via two biphenyl ether C–O bonds, and one biphenyl ether C–O and one biaryl C–C bonds, respectively. These macrocyclic rings are presumably derived by oxidative C–O or C–C coupling of the corresponding open chain bis(bibenzyls) such as perrottetin A (**5**).³ Among the type 2 of 16-membered ring bis(bibenzyls), plagiochin A (**1**)⁴ exhibits an interesting neurotrophic activity in the culture of fetal rat cerebral hemisphere.⁵ In order to establish a practical way for the preparation of plagiochin A (**1**), we have investigated the synthesis of plagiochin D (**4**), the simplest member of the plagiochins, as a part of studies on macrocyclic bis(bibenzyls).^{6,7} The synthesis of **4** reported by Nogradi *et al.*⁸ involved the Wurtz-type radical coupling at the position **a** for the macrocyclization.




Although we were succeeded in the final ring closure at the position **a** before Nogradi's report,⁹ all the attempted procedures for the ring closure at this position suffered considerably poor yield and reproducibility. Therefore, we turned our attention to employ intramolecular C–C cross-coupling at the position **b** in dihalogenated perrottetin A derivatives **6** by using organotransition metals. In this paper, we are pleased to report the synthesis of plagiochin D (**4**) realized by an intramolecular Still-Kelly reaction.¹¹

In accordance to the procedure used for the synthesis of marchantin A and riccardin B⁷ belonging to type I macrocyclic bis(bibenzyls), dibromoperrottetin A (**13**), a key precursor for the crucial C–C ring closure step, was prepared as follows. The A–C ring segment **8**⁷ and the phosphonate **7** readily derivable from *m*-anisaldehyde were effectively combined by Wardworth-Emmons reaction using sodium hydride to give rise to **9** in 94 % yield. The ester group of **9** was then reduced with lithium aluminum hydride to the alcohol, which was brominated with tetrabromomethane followed by treatment of trimethyl phosphite to afford the phosphonate **10** in 81 % yield. The Wardworth-Emmons reaction between **10** and **11**¹² smoothly proceeded to yield the bis(dibromostilbene) **12** in 89 % yield. Next catalytic hydrogenation of the double bonds in **12** was troublesome



Reagents and conditions: a) Br₂, CHCl₃, 70°C; b) NaBH₄, THF, 0°C; c) CBr₄, Ph₃P, MeCN; d) P(OMe)₃, 90°C; e) NaH, THF, 0°C; f) LiAlH₄, THF, 0°C; g) CBr₄, Ph₃P, MeCN; h) P(OMe)₃, 90°C; i) NaH, THF, 0°C; j) H₂, PtO₂, CH₂Cl₂; k) NaH, MOMCl, DMF.

due to being accompanied by an unanticipated debromination. After several attempts, **12** was cleanly hydrogenated using PtO_2 as catalyst and methylene chloride as solvent, thereby giving rise to the key intermediate **13** in 73 % yield after the protection of the liberated hydroxy group on the C ring as a MOM ether. We are now ready for performing the key intramolecular C–C ring closure step for the construction of the 16-membered ring.




As first of all, Ni(0) mediated intramolecular coupling¹³ which was recently utilized for the construction of the 12-membered vancomycin-type biaryl ring system¹⁴ and the 17-membered ring biaryl system¹⁵ was applied to **13** and **14**¹⁶ but failed to give the cyclized product **17**. Only the reduction product **16** was obtained, while the intramolecular Suzuki-Miyaura Pd(0) catalyzed reaction on **13** using bis(pinacolate)diboron¹⁷ made no C–C bond formation. In contrast, the intramolecular version of Still-Kelly reaction led to the crucial cyclization of **13**. A 0.01 M toluene solution of **13** was heated at 120°C with 1.1 equivalent of hexamethylditin and 5 mol% tetrakis(triphenylphosphine)palladium in the sealed tube to give the cyclized product **17** in 17 % yield along with a trimethylstannyl compound **15** (9 %) and the recovery **13** (45 %). The cyclization of diiodide **14** under the same conditions was comparable to that of **13**. The key intermediate for this ring closure was regarded as the isolated trimethylstannyl compound **15**. In fact, a diluted solution of **15** in toluene, upon treatment of 5 mol% tetrakis(triphenylphosphine)palladium in the sealed tube at 120°C, resulted in the formation of biaryl system **17** in 20 % yield. Finally, removal of the MOM groups in **17** by treatment of 47% HBr in MeOH afforded plagiochin D (**4**) in 87 % yield, all spectral data of which were superimposable with those of natural one.

In conclusion, the intramolecular Still-Kelly reaction was demonstrated to be quite useful for the crucial 16-membered ring formation at the position **b** of **13** in moderate and reproducible yield in spite of accompanying the formation of the dimmers (about 15 %). Thus, we have developed a new approach for the construction of the rigid macrocyclic biaryl ring system of the plagiochins. Further work toward total synthesis of the most biologically intriguing plagiochin A (**1**) is in progress.

Acknowledgments. We are grateful to Prof. Y. Asakawa and Dr. T. Hashimoto (Tokushima Bunri University) for kind donation of natural plagiochins A ~ D. This work was supported by a Grant-in-Aid for Scientific Research (No. 09680582) from the ministry of Education, Science, Sports and Culture, Japan.

References and Notes

- Asakawa, Y. *In Progress in the Chemistry of Organic Natural Products* (Hertz, W.; Kirby, G. W.; Moor, R. E.; Setglich, W.; Tamm, Ch. Ed), **1995**, vol. 65, p. 1-562, Springer, Wien.
- (a) Asakawa, Y. *In Bryophyte Development: Physiology and Biochemistry* (Chorpa, R. N.; Bhatla, S. C. Ed), **1990**, p. 259, CRS Press, Boca Raton. (b) Keserü, Gy. M.; Nogradi, M. *Nat. Prod. Rep.* **1995**, *12*, 69-75. (c) Gottsegen, A.; Nogradi, M.; Vermes, B.; Kajtar-Peredy, M.; Bihatsi-Karasai, E. *J. Chem. Soc. Perkin Trans.1* **1990**, 315-320. (d) Iyoda, M.; Sakaitani, M.; Otsuka, H.; Oda, M. *Tetrahedron Lett.* **1985**, *26*, 4777-4780.
- Toyota, M.; Tori, M.; Takikawa, K.; Shiobara, Y.; Kodama, M.; Asakawa, Y. *Tetrahedron Lett.* **1985**, *26*, 6097-6010.
- Hashimoto, T.; Tori, M.; Asakawa, Y.; Fukazawa, Y. *Tetrahedron Lett.* **1987**, *28*, 6925-6928.
- Fukuyama, Y.; Nakamura, K.; Kodama, M. *111th Annual Meeting of Pharmaceutical Society of Japan*, Tokyo, Japan, **1991**, Abstracts 2, p. 4. Plagiochin A (**1**) showed significantly not only acceleration of neurite sprouting but also enhancement of choline acetyl transferase activity in the culture of fetal rat cerebral hemisphere at 10 ~ 0.1 μ M.
- Kodama, M.; Shiobara, Y.; Sumitomo, H.; Matsumura, K.; Tsukamoto, M.; Harada, C. *Tetrahedron Lett.* **1985**, *26*, 877-880.
- Kodama, M.; Shiobara, Y.; Sumitomo, H.; Matsumura, K.; Tsukamoto, M.; Harada, C. *J. Org. Chem.* **1987**, *53*, 72-77.
- Keserü, G. M.; Mezey-Vandor, G.; Nogradi, M.; Vermes, B.; Kajtar-Peredy, M. *Tetrahedron* **1992**, *48*, 913-922.
- Fukuyama, Y.; Nakamura, K.; Kodama, M. *112th Annual Meeting of Pharmaceutical Society of Japan*, Fukuoka, Japan, **1992**, Abstracts 2, p. 78. The only way by which we could achieve the ring closure at the position **a** involved the cyclization through a sulfur atom from **18** to **19** and then the transformation of the formed sulfide linkage in **19** to the C-C bond by the benzyne-Stevens reaction.¹⁰ This procedure, however, had not been able to overcome poor yield (1%) and reproducibility for the transformation of **18** to **4**. All other trials for the C-C bond formation at the position **a** failed.



- Boekelheide, V.; Otsubo, T. *Tetrahedron Lett.* **1975**, 3881-3884.
- Still, J. K.; Groh, B. L. *J. Am. Chem. Soc.* **1987**, *109*, 813-817. Kelly, T. R.; Li, Q.; Bhushan, V. *Tetrahedron Lett.* **1990**, *31*, 161-164.
- Fleming, I.; Woolias, M. *J. Chem. Soc. Perkin I* **1979**, 829-837.
- Semmelhack, M. F.; Helquist, P.; Jones, L. D.; Keller, L.; Mendelson, L.; Ryono, L. S.; Smith, J. G.; Stauffer, R. D. *J. Am. Chem. Soc.* **1981**, *103*, 6460-6471.
- Nicolaou, K. C.; Chu, X.-J.; Ramanjulu, J. M.; Natarajan, S.; Bräse, S.; Rübsum, F.; Boddy, C. N. C. *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 1539-15540
- Carbonnelle, A.-C.; Zamora, E. G.; Beugelmans, R.; Roussi, G. *Tetrahedron Lett.* **1998**, *39*, 4471-4472.
- The diiodide **14** was prepared by the lithiation of **13** with *n*-BuLi, followed by quenching with I₂ at -78°C.
- Ishiyama, T.; Murata, M.; Miyaura, N. *J. Org. Chem.* **1995**, *60*, 7508-7510.