

Pyridine *N*-Sulphides. Some Simple *N*-Arylthiopyridinium Salts and a Possible Source of Arylsulphenium Ions

By RUDOLPH A. ABRAMOVITCH,* ADRIAN L. MILLER, and JACEK PILSKI†

(Department of Chemistry and Geology, Clemson University Clemson, SC 29631)

Summary *NN'*-Dipyridinium disulphide dichloride (1) and 1-(4-nitrophenylthio)pyridinium chloride (5) and its 4-NMe₂ derivative (2) have been synthesized and their structures established unambiguously; they undergo nucleophilic attack at sulphur rather than in the pyridine ring and the tetrafluoroborate from (5) serves as a possible source of 4-nitrophenylsulphenium ion which can attack anisole readily to give (7).


As we pointed out in an earlier paper,¹ virtually nothing is known about pyridine *N*-sulphides and *N*-thiopyridinium salts. Though the intermediacy of such compounds has been postulated a number of times² they have been isolated and characterised definitively in only a very few cases, such as in the case of isothiazolo[2,3-*a*]pyridinium salts.^{1,3} Two other possible pyridine *N*-sulphide derivatives have been reportedly isolated but not formulated as such. The reaction of pyridine with sulphur monochloride was said to give a compound S₂Cl₂.2C₅H₅N which was ionic.⁴ Similarly, pyridine and sulphur dichloride in carbon tetrachloride have been said to give a crystalline product SCl₂.2C₅H₅N.⁵ We have established (n.m.r. spectroscopy, microanalysis) that the former has structure (1) and is attacked by water at sulphur to give pyridine (89%), sulphur (19%), and SO₂.⁶ We also report the synthesis and properties of the first fully isolated and characterised simple *N*-arylthiopyridinium salts.

Reaction of 4-dimethylaminopyridine with *p*-nitrobenzenesulphenyl chloride in dry CCl₄ and work-up of the precipitate formed under N₂ and anhydrous conditions (dry box) gave colourless 4-dimethylamino-1-(4-nitrophenylthio)pyridinium chloride (2) (72%) which still contained one molecule of water of crystallisation! On drying

in vacuo at 60 °C it lost half a molecule of water of crystallisation, and at 80 °C it was obtained anhydrous.†§ On exposure to air the compound turned yellow, probably owing to some hydrolysis to 4,4'-dinitrodiphenyl disulphide (3).

Similar reaction of *p*-nitrobenzenesulphenyl chloride with an excess of dry pyridine (dried over CaH₂ and distilled under N₂ in a dry box) under conditions as anhydrous as we could achieve gave 1-(4-nitrophenylthio)pyridinium chloride (5) (82%), m.p. 62–64 °C (decomp.),‡¶ as yellow-brownish crystals containing one molecule of water of crystallisation. When the compound was dried further in a nitrogen stream at room temperature it analysed for 3/4 H₂O of crystallisation. It could not be dried at higher temperatures because it underwent thermolysis readily. Indeed, sublimation of (5) *in vacuo* at 120 °C gave pyridine (62%), (3) (37%), and *p*-nitrobenzenesulphenyl chloride (4) (65%) [contaminated with sublimed (5)]. Compound (4) was recovered unchanged under similar conditions so that (3) must result from the thermolysis of (5) and not from (4).

Treatment of (5) with water led to the formation of pyridine hydrochloride (88%), (3) (47%), and (6) (49%).⁷ Attempted nitration of (5) led only to the isolation of (6)



† On leave of absence from Akademia Rolnicza, Krakow, Poland.

‡ All new compounds gave microanalytical and spectral data consistent with their proposed structures.

§ ¹H N.m.r. (CDCl₃) δ 8.15 (d, 2H, *J*_{α,β} 8 Hz, H_α), 8.09 (d, 2H, *J*_{o,m} 6.5 Hz, H_o), 7.57 (d, 2H, *J*_{α,β} 8 Hz, H_β), 6.78 (d, 2H, *J*_{o,m} 6.5 Hz, H_m), 3.25 (s, 6H, CH₃).

¶ ¹H N.m.r. (CDCl₃) δ 8.65 (d, 2H, *J*_{α,β} 4 Hz, H_α), 8.25 (d, 2H, *J*_{o,m} 8.3 Hz, H_o), 7.7–7.2 (m, 5H, H_γ, H_β, H_m). The spectrum was different from the sum of those of the starting materials.

(91%). Reaction of (5) with sodium azide, or with triethylamine in dry acetonitrile, or attempted photolysis in the presence of anisole gave only (3), some of which at least could have arisen from (5) or other products during work-up. Reaction of (5) with KCN in acetonitrile at room temperature gave *p*-nitrophenylthiocyanate (85%). Anion exchange with tetrafluoroborate or 2,4,6-trinitrobenzenesulphonate occurred readily but the products hydrolysed too easily to be isolated in pure form. The above results confirm that both proton bases and carbon nucleophiles prefer to attack the sulphur atom in (5) rather than the pyridine ring.

It was hoped that (5), like *p*-nitrophenoxypyridinium tetrafluoroborate which undergoes unimolecular thermolysis to generate the *p*-nitrophenyloxonium ion,⁸ would undergo heterolysis to pyridine and the *p*-nitrophenylsulphenium

ion, $\text{O}_2\text{NC}_6\text{H}_4\text{S}^+$. When (5) was heated in anisole at 180–200 °C for 2 h only (3) could be isolated on work-up. On the other hand, when (5) in MeCN was first treated with AgBF_4 and then with anisole at room temperature, 4-nitrophenyl 4'-methoxyphenyl sulphide (7)⁹ (85%) and pyridinium tetrafluoroborate (89%) were isolated. It would appear, therefore, as though (5) (as the tetrafluoroborate) might indeed serve as a source of the sulphenium ion. Since, however, a more convenient and unambiguous source of this ion was found¹⁰ no further applications of (5) were sought.

We thank the N.S.F. and the National Institutes of Health for the support of this work.

(Received, 6th February 1981; Com. 141.)

¹ R. A. Abramovitch, M. N. Inbasekaran, A. L. Miller, and J. M. Hanna, Jr., *J. Heterocycl. Chem.*, submitted for publication.

² L. Goodman and N. Kharasch, *J. Am. Chem. Soc.*, 1955, **77**, 6541; G. K. Helmkamp, D. C. Owsley, W. M. Barnes, and H. N. Cassey, *ibid.*, 1968, **90**, 1635; V. J. Traynelis and J. N. Rieck, *J. Org. Chem.*, 1973, **38**, 4334.

³ G. G. Abott and D. Leaver, *J. Chem. Soc., Chem. Commun.*, 1973, 150.

⁴ R. C. Paul, J. Kishore, D. Singh, and K. C. Malhotra, *Indian J. Chem.*, 1970, **8**, 829; H. G. Heal and J. Kane, *J. Inorg. Nucl. Chem.*, 1967, **29**, 1539.

⁵ U. Wannagat and G. Schindler, *Angew. Chem.*, 1957, **69**, 784.

⁶ Cf. the behaviour of S_2Cl_2 : Gmelins, 'Handbuch der anorganischen Chemie.' B/3, 8 Auflage, Verlag Chemie GMBH, Weinheim/Bergstr., 1963, p. 1770.

⁷ Th. Zincke and S. Lenhardt, *Liebigs Ann. Chem.*, 1913, **400**, 2.

⁸ R. A. Abramovitch, M. Inbasekaran, and S. Kato, *J. Am. Chem. Soc.*, 1973, **95**, 5428; R. A. Abramovitch, G. Alvernhe, and M. N. Inbasekaran, *Tetrahedron Lett.*, 1977, 1113.

⁹ S. L. Bass and T. B. Johnson, *J. Am. Chem. Soc.*, 1930, **52**, 1146.

¹⁰ R. A. Abramovitch and J. Pilski, *J. Chem. Soc., Chem. Commun.*, 1981, 704, following Communication.