## negative charge<sup>[5]</sup> ("excess-electron transport"). A similar hopping model was also applied to explain long-distance charge transport through proteins.<sup>[6]</sup> Hopping is especially favored if the amino acids carry side chains that can be easily oxidized (tyrosine, tryptophan, and histidine). For instance, a multistep electron-hole transport cascade involving three tryptophan units was proposed for the photoactivation process in the enzyme photolyase.<sup>[6a]</sup> Presumably, not only tryptophan but also tyrosine and histidine are involved in the mechanism of radical initiation in ribonucleotide reductase, where an electron hole is transported over a distance of 35 Å.<sup>[6e]</sup> In these systems, transient intermediates occurring in the multistep charge transport have not yet been detected, and it is not clear whether the electron transfer is best described as proceeding through space or through the peptide

In order to examine electron-hole transport in peptides, we synthesized the series of model systems 1a-d bearing three different redox sites, that is, a modified tetrahydrofuran as a charge-injection system,<sup>[8]</sup> 2,4,6-trimethoxyphenylalanine<sup>[9]</sup> as an intermediate charge carrier, and tyrosine as a final electron donor. An oligoproline spacer of variable length (n = 0, 1, 3, 3)5) was chosen to separate the tyrosine residue from the 2,4,6trimethoxyphenylalanine moiety. As is known from related studies of electron-hole transfer in DNA, irradiation of the pivaloylated tetrahydrofuran unit in compounds like 1a-d will generate the dihydrofuranosyl radical cations, 2a-d, through Norrish  $\alpha$  cleavage and subsequent  $\beta$  elimination of the phosphate leaving group.<sup>[10]</sup> These radical cations have

## OMe OMe MeC MeC ÔMe = 308 nm ťBu II O oPO(OPh)₂ 2a-d 0), **1b** (*n* = 1), **1c** (*n* = 3), **1d** (*n* = 5) OMe OMe MeC MeO ÒМе - H 3a 4a-d

Scheme 1. Mechanism for the photolysis of 1 a-d and the subsequent electron-transfer reactions.

[\*] Prof. Dr. B. Giese, Dr. M. Napp, Dr. O. Jacques, A. M. Taylor Department of Chemistry, University of Basel St. Johanns-Ring 19, 4056 Basel (Switzerland) Fax: (+41) 61-267-1105 E-mail: bernd.giese@unibas.ch Dipl.-Chem. H. Boudebous, Prof. Dr. J. Wirz Department of Chemistry, University of Basel Klingelbergstrasse 80, 4056 Basel (Switzerland) [\*\*] This work was supported by the Swiss National Science Foundation. Supporting Information for this article is available on the WWW under http://www.angewandte.org or from the author.

Angew. Chem. Int. Ed. 2005, 44, 4073-4075

strong oxidizing properties<sup>[11]</sup> and can oxidize the neighboring 2,4,6-trimethoxyphenylalanine side chain (Scheme 1).<sup>[12]</sup>

Compounds 1a-d were dissolved in acetonitrile/water (3:1) and irradiated with 308-nm pulses from a XeCl excimer laser,<sup>[13]</sup> which allowed selective excitation of the pivaloyl chromophore. The transient absorption spectra observed after a delay of 50 ns with respect to the laser pulse indicated the presence of radical cations 3a-d with an absorption maximum at 550 nm (Figure 1a shows the spectrum for system **1c** as a typical example).<sup>[14,15]</sup> This signal decayed

bonds.<sup>[7]</sup>

DOI: 10.1002/anie.200500391

## **Electron Transfer in Peptides**

## **Multistep Electron Transfer in Oligopeptides: Direct Observation of Radical Cation** Intermediates\*\*

Bernd Giese,\* Matthias Napp, Olivier Jacques, Hassen Boudebous, Alexander M. Taylor, and Jakob Wirz

For the transport of electrons or electron holes over long distances, nature often uses DNA double strands or proteins. With DNA, experiments<sup>[1]</sup> and calculations<sup>[2]</sup> have clearly shown that these processes can occur by a hopping mechanism, where the charge transport takes place in a multistep reaction.<sup>[3]</sup> Purines act as carriers of positive charge<sup>[4]</sup> ("electron-hole transport"), and pyrimidines are carriers of



**Figure 1.** Transient absorption spectra observed after a delay of a) 50 ns and b) 1  $\mu$ s with respect to the irradiation of **1 c** in degassed acetonitrile/water (3:1) at a concentration of 8 mm.

within 1  $\mu$ s and was replaced by the characteristic signal of the tyrosyl radical with a maximum absorbance at 410 nm (Figure 1 b).<sup>[16]</sup>

Kinetic traces of the absorbance decay at 550 nm and of the growth observed at 410 nm obeyed the first-order rate law well, and the rate constants determined by least-squares fitting at the two wavelengths were the same within the limits of error. A typical example is shown in Figure 2.



**Figure 2.** Kinetic traces of the transient absorptions at 410 and 550 nm for **1c** (7 mM) in degassed acetonitrile/water (3:1). The decay rate constant at 550 nm is  $3.1(\pm 0.2) \times 10^6 \text{ s}^{-1}$  and the absorbance growth constant at 410 nm is  $3.5(\pm 0.3) \times 10^6 \text{ s}^{-1}$ .

However, the rate constants observed with model peptides **1a–d** increased linearly with the concentration of these compounds. This pointed to competition of intra- and intermolecular electron transfer. The slopes of these plots correspond to the intermolecular rate contributions  $k_{\text{inter}}$ , and the values so obtained ( $k_{\text{inter}} \approx 4 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ ) agree well with the rate constants of diffusion expected for compounds of this size. The intramolecular contributions, namely the rate constants  $k_{\text{intra}}$ , for electron transfer from the tyrosine side chain to the 2,4,6-trimethoxyphenylalanine radical cation moiety (**3a–d**→**4a–d**) were determined from the intercepts (Table 1).<sup>[17]</sup>

The rate constants  $k_{intra}$  (Table 1) decrease by a factor of only about two when the spacer between the aromatic redox centers is elongated by addition of two proline units (compare **1b** with **1c** and **1c** with **1d**). For an electron transfer through the peptide bonds one would expect a rate decrease of about 500 for two additional proline units.<sup>[18]</sup> This indicates that the electron transfer does not proceed by a through-bond mechanism. We attribute the small influence of additional proline spacers upon the  $k_{intra}$  value to the conformational

**Table 1:** Rate constants for intramolecular  $(k_{intra})$  and intermolecular  $(k_{intra})$  electron transfer, determined from the intercepts and slopes of rate-versus-concentration plots. The number of proline units between the aromatic side chains is given by *n*, and  $\sigma_l$  is the length of the throughbond pathway, based on the number of covalent bonds between the aromatic redox centers.<sup>[6f]</sup>

	n	k <sub>intra</sub> [s <sup>-1</sup> ]	k <sub>inter</sub> [м <sup>-</sup> 1 s <sup>-1</sup> ]	$\sigma_{\!\scriptscriptstyle  }[{ m \AA}]$
1a	0	8.6(±0.6)×10 <sup>6</sup>	$\leq$ 5 $\times$ 10 <sup>8</sup>	9.8
1 b	1	1.1(±0.1)×10 <sup>6</sup>	4.3 (±0.4) × 10 <sup>8</sup>	14.0
1c	3	$5.1(\pm 0.3) \times 10^{5}$	$3.0(\pm 0.1) \times 10^8$	22.4
٦d	5	$2.8(\pm 0.3) \times 10^{5}$	$3.2(\pm 0.1) \times 10^8$	30.8

flexibility of the aromatic amino acids.<sup>[19]</sup> The rate does increase considerably when the proline spacer is absent, as in **1a**, where the aromatic side chains are in close contact with each other.

The radical cation of 1,3,5-trimethoxybenzene<sup>[12]</sup> should not be capable of oxidizing phenol to the phenol radical cation (oxidation potential  $E_{ox}$ (PhOH/PhOH<sup>++</sup>) = 1.50 V versus a NHE).<sup>[20]</sup> But oxidation of phenol to the phenoxy radical in aqueous solution (pH 7) has a much lower potential of  $E_{ox}$ (PhOH/PhO<sup>+</sup>) = 0.86 V versus a NHE.<sup>[21]</sup> Thus, oxidation of tyrosine in the systems **3a–d** can easily occur, if the electron transfer is coupled with proton transfer to the solvent. We measured a kinetic deuterium isotope effect  $k_{\rm H}/$  $k_{\rm D}$  of 1.8(±0.2).<sup>[22]</sup> This value is in good agreement with the isotope effect of 2.0–2.5 for the oxidation of tyrosine by a ruthenium–tris(bipyridine) complex in aqueous media, where a proton-coupled electron transfer occurs.<sup>[23]</sup>

In summary, electron-hole transport in peptide radical cations 3a-d occurs by a hopping mechanism between the aromatic side chains of the amino acids, which is similar to the long-distance hole transport through DNA.<sup>[1-4]</sup>

Received: February 1, 2005 Published online: May 25, 2005

**Keywords:** charge transport · electron transfer · flash photolysis · peptides · radicals

- E. Meggers, M. E. Michel-Beyerle, B. Giese, J. Am. Chem. Soc. 1998, 120, 12950.
- [2] J. Jortner, M. Bixon, T. Langenbacher, M. E. Michel-Beyerle, Proc. Natl. Acad. Sci. USA 1998, 95, 12759.
- [3] Recent reviews: "Long-Range Charge Transfer in DNA": Top. Curr. Chem. 2004, 236 and 237.
- [4] B. Giese, J. Amaudrut, A.-K. Köhler, M. Spormann, S. Wessely, *Nature* 2001, 412, 318.
- [5] a) S. Breeger, U. Hennecke, T. Carell, J. Am. Chem. Soc. 2004, 126, 1302; b) T. Ito, S. E. Rokita, Angew. Chem. 2004, 116, 1875; Angew. Chem. Int. Ed. 2004, 43, 1839.
- [6] a) C. Aubert, M. H. Vos, P. Mathis, A. P. M. Eker, K. Brettel, Nature 2000, 405, 586; b) J. R. Winkler, Curr. Opin. Chem. Biol. 2000, 4, 192; c) E. G. Petrov, V. May, J. Phys. Chem. A 2001, 105, 10176; d) C. C. Page, C. C. Moser, P. L. Dutton, Curr. Opin. Chem. Biol. 2003, 7, 551; e) J. Stubbe, D. G. Nocera, C. S. Yee, M. C. Y. Chang, Chem. Rev. 2003, 103, 2167; f) H. B. Gray, J. R. Winkler, Q. Rev. Biophys. 2003, 36, 341; g) T. Morita, S. Kimura, J. Am. Chem. Soc. 2003, 125, 8732.

- [7] The question of whether the electron travels through the peptide bonds by a bridge-mediated superexchange mechanism or by a hopping process has been the subject of several investigations:
  a) A. K. Mishra, R. Chandrasekar, M. Faraggi, M. H. Klapper, J. Am. Chem. Soc. 1994, 116, 1414; b) K. Bobrowski, J. Poznanski, J. Holcman, K. L. Wierzchowski, J. Phys. Chem. B 1999, 103, 10316; c) R. A. Malak, Z. Gao, J. F. Wishart, S. S. Isied, J. Am. Chem. Soc. 2004, 126, 13888; d) K. Yanagisawa, T. Morita, S. Kimura, J. Am. Chem. Soc. 2004, 126, 12780; e) F. Polo, S. Antonello, F. Formaggio, C. Toniolo, F. Maran, J. Am. Chem. Soc. 2005, 127, 492.
- [8] The synthesis of the charge-injecting system was performed following procedures reported earlier: a) R. Glatthar, M. Spichty, A. Gugger, R. Batra, W. Damm, M. Mohr, H. Zipse, B. Giese, *Tetrahedron* 2000, 56, 4117; b) A. Marx, P. Erdmann, M. Senn, S. Körner, T. Jungo, M. Petretta, P. Imwinkelried, A. Dussy, K. J. Kulicke, L. Macko, M. Zehnder, B. Giese, *Helv. Chim. Acta* 1996, 79, 1980. (See the Supporting Information for further details.)
- [9] We used the racemate: D. Dauzonne, R. Royer, *Synthesis* 1987, 399.
- [10] A. Gugger, R. Batra, P. Rzadek, G. Rist, B. Giese, J. Am. Chem. Soc. 1997, 119, 8740.
- [11] When 2,3-dihydrofuran is used as a simple model system, the reduction potentials of radical cations like 2a-d are estimated to be larger than 1.27 V versus a normal hydrogen electrode (NHE): K. Bernhard, J. Geimer, M. Canle-Lopez, J. Reynisson, D. Beckert, R. Gleiter, S. Steenken, *Chem. Eur. J.* 2001, 7, 4640.
- [12] The half-wave oxidation potential of 1,3,5-trimethoxybenzene is 1.25 V versus a NHE: A. Zweig, W. G. Hodgson, W. H. Jura, J. Am. Chem. Soc. 1964, 86, 4124.
- [13] Solutions of 1a-d (0.3-8 mM) were degassed in a quartz cell of 4.5-cm path length and irradiated with laser pulses (Lambda-Physik COMPex 205, λ = 308 nm, pulse width of 25 ns, 100-150 mJ per pulse) at RT (22-24°C). An iStar 720 ICCD camera (Andor) was used for the detection of transient absorption spectra. Transient kinetics were recorded with a 1P28 photomultiplier (Hamamatsu). For a detailed description, see: a) J. Wirz, *Pure Appl. Chem.* 1984, *56*, 1289; b) E. Leyva, M. S. Platz, G. Persy, J. Wirz, *J. Am. Chem. Soc.* 1986, *108*, 3783; c) S. Gerber, J. Wirz, *EPA Newsl.* 1989, *36*, 19.
- [14] P. O'Neill, S. Steenken, D. Schulte-Frohlinde, J. Phys. Chem. 1975, 79, 2773.
- [15] The transient absorption spectrum recorded after oxidation of N-acetyl-2,4,6-trimethoxyphenylalanine methyl ester (5) by

photosensitized electron transfer with 1,4-dicyanonaphthalene as a sensitizer and biphenyl as a codonor is similar to the spectrum shown in Figure 1 a (see the Supporting Information). Furthermore, a compound with the pivaloylated tetrahydrofuran unit directly connected to *N*-acetyl-2,4,6-trimethoxyphenylalanine was synthesized (6). Laser flash photolysis of this compound again gave a spectrum that was practically identical to the one shown in Figure 1 a (see the Supporting Information).

- [16] K. M. Bansal, R. W. Fessenden, Radiat. Res. 1976, 67, 1.
- [17] The irradiations were carried out at concentrations of 0.3–8 mM, which provided  $k_{intra}$  values from the intercepts of rate-versusconcentration plots with a standard error of about 10% (see the Supporting Information). The decay of the radical cation of 2,4,6-trimethoxyphenylalanine in the absence of tyrosine was much slower (dissociation constant  $k_d = 1.1 \cdot 10^4 \text{ s}^{-1}$ ) than the intramolecular electron-transfer reaction **3a–d**–**4a–d** ( $k_{intra} = 0.28 \times 10^6$ –8.6 × 10<sup>6</sup> s<sup>-1</sup>). These intramolecular rates are 10–100 times faster than the observed electron-transfer rates between a tryptophan radical cation and tyrosine.<sup>[7a,b]</sup> This is in accord with the higher oxidation potential ( $E_{ox}$ ) of trimethoxyphenylalanine<sup>[12]</sup> compared to tryptophan ( $E_{ox} = 1.02 \text{ V}$  versus a NHE).<sup>[21]</sup>
- [18] By addition of two proline units, the length of the through-bond pathway (σ<sub>1</sub>, 1.4 Å times the number of covalent bonds in the effective path) is increased by 8.4 Å. With a distance-decay constant of β = 0.73 Å<sup>-1</sup>, the k<sub>intra</sub> value is expected to drop by a factor of about 500: D. R. Casimiro, L. L. Wong, J. L. Colon, T. E. Zewert, J. H. Richard, I.-J. Chang, J. R. Winkler, H. B. Gray, J. Am. Chem. Soc. **1993**, 115, 1485.
- [19] By using <sup>1</sup>H-NOESY experiments in  $[D_3]$  acetonitrile, conformation studies of **1c** were performed. The proline spacer adopts the all-*trans* conformation in about 80%. Subsequent molecularmodeling analysis (with the MakroModel 7.2 program) of compounds **1a–d** in a fixed all-*trans* conformation revealed that, due to the conformational flexibility of the aromatic side chains, different distances exist between the redox centers.
- [20] S. Canonica, B. Hellrung, J. Wirz, J. Phys. Chem. A 2000, 104, 1226.
- [21] A. Harriman, J. Phys. Chem. 1987, 91, 6102.
- [22] The rate of intramolecular electron transfer in compound 1c (acetonitrile/D<sub>2</sub>O (3:1)) is  $2.9(\pm 0.2) \times 10^5 \text{ s}^{-1}$ .
- [23] M. Sjödin, S. Styring, B. Åkermark, L. Sun, L. Hammarström, *Philos. Trans. R. Soc. London Ser. B* 2002, 357, 1471.