β -Acyloxysulfonyl Tethers for Intramolecular Diels—Alder Cycloaddition Reactions

LETTERS 2005 Vol. 7, No. 15 3203–3206

ORGANIC

Nataliya Chumachenko,*,† Paul Sampson,*,† Allen D. Hunter,‡ and Matthias Zeller‡

Department of Chemistry, Kent State University, Kent, Ohio 44242, and Department of Chemistry, Youngstown State University, One University Plaza, Youngstown, Ohio 44555

psampson@kent.edu

Received April 26, 2005

ABSTRACT



 β -Hydroxy sulfone-based tethers were employed for the first time to achieve thermally mediated intramolecular Diels–Alder cycloaddition. The reactions proceeded with complete regioselectivity and high (10/1) to complete *endo/exo*-selectivity and resulted in the preferential formation of one of the two possible *endo*-cycloadducts. The yields and stereoselectivities were proportional to the bulk of the R¹ substituent on the β -acyloxysulfonyl tether.

It is known that temporary tethers can be used to control the regio- and stereoselectivity in Diels–Alder cycloaddition reactions.¹ However, to the best of our knowledge, only two types of tethers based on sulfur-containing functional groups have been reported, involving the use of sulfonate esters² and sulfonamides.³ On the other hand, the intermolecular Diels–Alder reactivity of sulfonyl-substituted dienes and dienophiles has been studied. 1-Sulfonyl-1,3-dienes (the data are scarce and restricted to cyclic dienes)^{4–6} and 2-sulfonyl-

1,3-dienes⁷⁻¹³ are prone to self-dimerization^{4,8-10} through Diels–Alder reactions. They also undergo highly regioselective cycloaddition with electron-rich^{5,9-11} dienophiles and less regioselective reactions with electron-deficient^{6,9,10,12,13}

[†] Kent State University.

[‡] Youngstown State University (X-ray crystallography).

⁽¹⁾ Gauthier, D. R.; Zandi, K. S.; Shea, K. J. Tetrahedron 1998, 54, 2289.

^{(2) (}a) Metz, P.; Fleischer, M.; Fröhlich, R. *Tetrahedron* 1995, *51*, 711.
(b) Plietker, B.; Seng, D.; Fröhlich, R.; Metz, P. *Tetrahedron* 2000, *56*, 873–879.

⁽³⁾ Brosius, A. D.; Overman, L. E.; Schwink, L. J. Am. Chem. Soc. 1999, 121, 700.

^{(4) (}a) Hartke, K.; Gleim, H.-U. *Liebigs Ann. Chem.* **1976**, 716. (b) Hartke, K.; Jung, M. H.; Zerbe, H.; Kämpchen, T. *Arch. Pharm. (Weinheim, Ger.)* **1986**, *319*, 890. (c) Bridges, A. J.; Fischer, J. W. J. Chem. Soc., Perkin Trans. 1 **1983**, 2359.

^{(5) (}a) Posner, G. H.; Wettlaufer D. G. *Tetrahedron Lett.* 1986, 27, 667.
(b) Posner, G. H.; Switzer, C. J. Org. Chem. 1987, 52, 1642. (c) Posner, G. H.; Kinter, C. M. J. Org. Chem. 1990, 55, 3967.

⁽⁶⁾ Miller, J. A.; Ullah, G. M. J. Chem. Res. (Miniprint) 1988, 2737.

⁽⁷⁾ Synthesis of 2-sulfonyl-1,3-dienes: (a) Andell, O. S.; Bäckvall, J.-E. *Tetrahedron Lett.* **1985**, 26, 4555. (b) Bäckvall, J.-E.; Juntunen, S. K.; Andell, O. S. *Org. Synth.* **1990**, 68, 148. (c) Bäckvall, J.-E.; Nájera, C.; Yus, M. *Tetrahedron Lett.* **1988**, 29, 1445. See also refs 8–10, 12, and 13. (8) (a) Chou, T.-S.; Hung, S. C.; Tso, H.-H. *J. Org. Chem.* **1987**, 52,

^{3394. (}b) Hoffmann, H. M. R.; Weichert, A.; Slawin, A. M. Z.; Williams, D. J. *Tetrahedron* **1990**, *46*, 5591.

^{(9) (}a) Inomata, K.; Kinoshita, H.; Takemoto, H.; Murata, Y.; Kotake, H. *Bull. Chem. Soc. Jpn.* **1978**, *51*, 3341. (b) Chou, T.-S.; Hung, S.-C. *J. Org. Chem.* **1988**, *53*, 3020.

⁽¹⁰⁾ Bäckvall, J.-E.; Juntunen, S. K. J. Am. Chem. Soc. 1987, 109, 6396.
(11) (a) Bäckvall, J.-E.; Plobeck, N. A.; Juntunen, S. K. Tetrahedron Lett. 1989, 30, 2589. (b) Barnwell, N.; Beddoes, R. L.; Mitchell, M. B.; Joule, J. A. Heterocycles 1994, 37, 175. (c) Bäckvall, J.-E.; Rise, F. Tetrahedron Lett. 1989, 30, 5347.

^{(12) (}a) Cuvigny, T.; Du Penhoat, C. H.; Julia, M. *Tetrahedron* **1986**, 42, 5329. (b) Chou, S.-S. P.; Chao, M.-H. *J. Chin. Chem. Soc.* **1996**, 43, 53. (c) de la Pradilla, R. F.; Montero, C.; Viso, A. *Chem. Commun.* **1998**, 409.

^{(13) (}a) Chou, S.-S. P.; Sun, D.-J. J. Chem. Soc., Chem. Commun. 1988, 1176. (b) Back, T. G.; Lai, E. K. Y.; Muralidharan, K. R. J. Org. Chem. 1990, 55, 4595.

alkenes. If a 2-sulfonyl-1,3-diene bears an electron-donating group at C-3, the regioselectivity of the cycloaddition with α , β -unsaturated carbonyl compounds is increased in the presence of Lewis acids.¹³ However, this is not the case for nonsubstituted 2-sulfonyl butadienes.¹⁰ Connecting a sulfonebearing diene and an electron-deficient dienophile by a tether bearing a stereogenic center might be expected not only to improve the regioselectivity but also to cause the cycload-dition to proceed in a diastereoselective manner. The presence of a sulfone group in the final cycloadduct provides various possibilities for further transformations. We have begun to explore the potential utility of sulfone-based tethers for cycloaddition reactions and present here our preliminary results.¹⁴

We considered the (*Z*)-butadienyl sulfinate anion, readily available from butadiene sulfone (1),¹⁵ as an attractive building block to prepare 1-sulfonyl-1,3-dienes. We found¹⁶ that Zn (*Z*)-sulfinate **2a**, freshly prepared from the corresponding Li (*Z*)-sulfinate in aqueous medium, reacted with ethylene oxide or propylene oxide regioselectively to produce the corresponding β -hydroxy sulfones as (*E*)-/(*Z*)-mixtures.

Isomerization to the (*E*)-stereoisomers occurred readily in the presence of DMAP to give cleanly (*E*)-butadienyl β -hydroxy sulfones **3** and **4** in 30% isolated yield from **1** (Scheme 1).



As an alternative approach to β -hydroxy sulfone tethers, we attempted to employ α -halo ketones as electrophiles to trap the (*Z*)-butadienyl sulfinate anion. However, Zn, Li, or K butadienyl sulfinates, freshly prepared from **1** in THF, failed to react with α -bromo ketones, which were recovered unreacted. Bouchez and Vogel reported¹⁷ that allyl or β -carboxyalkyl sulfinate anions, produced in situ from the corresponding silyl sulfinates, reacted with α -bromo esters to generate the corresponding sulfones. To our delight, this approach worked well for the reaction of (*Z*)-butadienyl silyl sulfinate **2b** with α -bromo ketones to furnish (*Z*)- β -oxo sulfones **5** and **6**, which were reduced in high yield to

(16) Chumachenko, N.; Sampson, P. Submitted.

produce the corresponding (*Z*)- β -hydroxy sulfones **7** and **8**. DMAP-mediated isomerization required longer times than were employed for the preparation of sulfones **3** and **4** but still provided quantitative yields of the corresponding (*E*)- β -hydroxy sulfones **9** and **10** (Scheme 2).



The esterification of (E)- β -(dienesulfonyl) alcohols **3**, **4**, **9**, and **10** with methacrylic acid (**11**) and acrylic acid (**12**) proceeded in good yield using conventional DCC/DMAP methods to afford the corresponding acrylate esters **14a**-**f**,**i** (Scheme 3, Table 1, method A). No alcohol decomposition

Table 1. Esterification of (E)- β -(Dienylsulfonyl) Alcohols 3, 4, 9, and 10 to Afford Acrylate Esters 14a-i

,					
product	\mathbb{R}^1	\mathbb{R}^2	$method^a$	yield $(\%)^b$	
14a	Н	Me	А	85	
14b	Me	Me	Α	93	
14c	Ph	Me	Α	81	
14d	t-Bu	Me	Α	50^{c}	
14e	Me	Н	Α	83	
14f	\mathbf{Ph}	н	Α	81	
14g	Me	\mathbf{Br}	Α	28^c	
14h	Ph	\mathbf{Br}	В	$47^{c} (59^{d})$	
14i	Н	Н	Α	85	

^{*a*} A: DCC, DMAP, CH₂Cl₂, rt. B: 2,4,6-Trichlorobenzoyl chloride, Et₃N, LiCl, THF, rt. ^{*b*} Isolated yields after purification by silica chromatography. ^{*c*} Nonoptimized yields. ^{*d*} Based on consumed β -hydroxy sulfone.

was observed under the reaction conditions. Not surprisingly,¹⁸ esterification using α -bromoacrylic acid (13) was

⁽¹⁴⁾ Chumachenko, N.; Sampson, P. *Abstracts of Papers*, 228th National Meeting of the American Chemical Society, Philadelphia, PA, August 22–26, 2004; American Chemical Society: Washington, DC, 2004; ORGN 621.

 ^{(15) (}a) Polunin, E. V.; Zaks, I. M.; Moiseenkov, A. M.; Semenovskii,
 A. V. *Izvestiya Akademii Nauk SSSR, Ser. Khim.* 1979, *3*, 641. (b) Krug,
 R. C.; Rigney, J. A.; Tichelaar, G. R. *J. Org. Chem.* 1962, *27*, 1305. (c)
 Crumbie, R. L.; Ridley, D. D. *Aust. J. Chem.* 1981, *34*, 1017.

⁽¹⁷⁾ Bouchez, L.; Vogel, P. Synthesis 2002, 225.

⁽¹⁸⁾ To the best of our knowledge, no examples of esterification of α -bromoacrylic acid (13) using an alcohol have been reported. Esterification to afford a phenyl ester (10% yield) was reported in one case: Schmidt, A.; Bach, E.; Schollmeyer, E. *Dyes Pigments* 2003, *56*, 27. Three esters of 13 have been prepared via carboxylate alkylation using (a) benzyl bromide [Osborne, N. F. *J. Chem. Soc., Perkin Trans. 1* 1980, 150], (b) allyl bromide [Bhat, L.; Steinig, A. G.; Appelbe, R.; de Meijere A., *Eur. J. Org. Chem.* 2001, *9*, 1673], and (c) epichlorohydrin [Mukhitdinova, N. A.; Khodzhaeva, F.; Askarov, M. A. *Dokl. Akad. Nauk UzSSR* 1977, 39 (CA 87 134867)].



problematic. The best yield to date was achieved when alcohol **9** was treated with the mixed anhydride derived from α -bromoacrylic acid (**13**) and 2,4,6-trichlorobenzoyl chloride in the presence of LiCl and triethylamine (Scheme 3, Table 1, method B).

Thermally mediated intramolecular Diels-Alder cycloaddition (IMDA) reaction of **14a-h** (Scheme 3, Table 2)

Table 2. Diels-Alder Cycloaddition Reactions of 14a-h							
entry	compd	<i>T</i> (°C)	time (h)	ratio of 15/16/17/14 ª	yield of 15a-h (%)		
1	14a	130	29.5	100//33/20	42 of 15a + 17a		
2	14b	128	32.5	100/25/12/22	47 of $15b + 16b$		
3	14b	140	41	100/25/12/	36		
4	14c	138	28	100/25/10/20	49		
5	14c	145	50	100/25/10/	49		
6	14d	138	47	100/4/12/	65		
7	14e	130	43	100/30//	54 + 18 of $16e$		
8	14f	127	43	100/25//23	57 ± 16 of $\mathbf{16f}$		
9	14g	115	21	100/27/9/ ^b	58		
10	14h	127	21	$100/12/8/^{b}$	63		

^{*a*} Determined from ¹H NMR analysis of the crude reaction mixtures. ^{*b*} Ratio can only be estimated because **16** and **17** are unstable.

occurred with complete regioselectivity and high (10/1) to complete *endo/exo*-selectivity (except when $R^1 = H$ (14a)) and resulted in the preferential formation of one of the two possible endo-cycloadducts. For substrates 14a-d, derived from methacrylic acid (11), the major endo-cycloadducts 15a-d were separated as single diastereomers after crystallization, while the minor exo-cycloadducts 17a,d and endocycloadducts **16b**-**d** were characterized by ¹H and ¹³C NMR analysis of the residual mother liquors. For cycloadditions involving the substrates **14e**,**f**, derived from acrylic acid (**12**), both the major endo-cycloadducts 15e,f and the minor endocycloadducts 16e,f were cleanly separated as single diastereomers by column chromatography. For cycloadditions involving substrates 14g,h, derived from 2-bromoacrylic acid (13), the major *endo*-cycloadducts **15g**,**h** were purified by silica chromatography. Unfortunately, the minor endocycloadducts 16g.h underwent partial dehydrobromination on exposure to silica to afford the corresponding dienes 18g,h. In the case of 16g, we were able to obtain a sample of diastereomerically pure material from the silica column; however, the dehydrobromination of 16h was too fast to permit its isolation. The presence and relative amount of 16h, as well as the *exo*-products **17b**,**c** (derived from methacrylic acid (11)) and 17g,h (derived from 2-bromoacrylic acid (13)), were tentatively determined on the basis of the integration of H^{4a} in the ¹H NMR of the crude reaction mixtures. The absence of the formation of *exo*-products 17e,f in the Diels-Alder cycloaddition of acrylate substrates **14e.f** might have been expected, while the high endo/exo-selectivity observed for methacrylate substrates 14a-d, as well as bromoacrylate substrates 14g,h, was a pleasant surprise. It is welldocumented that dienophiles derived from methacrylic acid (11) give poor stereoselectivity in Diels-Alder reactions and often show a proclivity to undergo *exo*-cycloaddition.^{19,20}

The relative stereochemistry of the major *endo*-cycloadducts **15a,b,g** and minor *endo*-cycloadduct **16g** were confirmed by their X-ray crystal structures.²¹ The relative stereochemistry of the remaining *endo*-cycloadducts was established by NOE experiments and ¹H NMR analysis.

We observed that increasing the bulk of the substituent R^1 on the tether backbone is beneficial for avoiding side reactions during the IMDA reaction. The nonsubstituted substrate **14i**, derived from acrylic acid (**12**), failed to give any intramolecular cycloaddition product and was completely polymerized on heating (120 °C) in toluene. Large amounts of polymerization byproducts were produced in IMDA reactions of the other nonsubstituted (**14a**) and Mesubstituted (**14b,e,g**) substrates, while for Ph- (**14c,f**) and *t*-Bu-substituted (**14d**) substrates, no significant tar formation was observed during the IMDA reaction.

⁽¹⁹⁾ Saito, A.; Ito, H.; Taguchi, T. Org. Lett. 2002, 4, 4619.

^{(20) (}a) Evans, D. A.; Chapman, K. T.; Bisaha, J. J. Am. Chem. Soc. **1988**, 110, 1238. (b) Hayashi, Y.; Nakamura, M.; Nakao, S.; Inoue, T.;
Shoji, M. Angew. Chem., Int. Ed. **2002**, 41, 4079.

^{(21) (}a) Zeller, M.; Hunter, A. D.; Sampson, P.; Chumachenko, N. Acta Crystallogr. 2004, E60, o731. (b) Zeller, M.; Hunter, A. D.; Sampson, P.; Chumachenko, N. Acta Crystallogr. 2004, E60, o729. (c) Zeller, M.; Hunter, A. D.; Sampson, P.; Chumachenko, N. Acta Crystallogr. 2004, E60, o727. (d) Zeller, M.; Hunter, A. D.; Sampson, P.; Chumachenko, N. Acta Crystallogr. 2004, E60, o724.

The *endo*-major (15)/*endo*-minor (16) diastereoselectivity in IMDA reactions involving Me- (14b,e,g) and Phsubstituted (14c,f) substrates was comparable (\sim 4/1), while for *t*-Bu-substituted substrate 14d, the diastereoselectivity was improved to \sim 25/1.

endo/exo-Diastereoselectivity was significantly increased (from 3/1 to 10/1) on introducing a substituent R¹ (Me) in place of a hydrogen atom on the tether backbone (Table 2, compare entries 1 and 2). Interestingly, it remained the same (\sim 9/1 to \sim 14/1) on changing R¹ from a relatively small Me group to the larger Ph and *t*-Bu substituents (see Table 2, entries 2–6, 9, and 10).

The cycloaddition of α -bromoacrylate substrates **14g,h** proceeded faster than for the analogous methacrylate (**14b,c**) and acrylate substrates (**14e,f**) and required ~10 °C lower reaction temperatures.

Furthermore, the fact that the acrylate substrate **14i** failed to give any IMDA reaction products, while the corresponding methacrylate derivative **14a** was converted into IMDA products in 42% yield, contradicts the conventional wisdom that normal-electron demand Diels—Alder reactions of methacrylate dienophiles are usually more difficult than for the corresponding acrylate substrates.^{19,20}

We propose a transition state for the above cycloaddition reactions (Figure 1) in which (i) steric interactions between R^1 and the carbonyl group and (ii) the stereoelectronically preferred s-trans ester conformation¹⁹ are responsible for the observed diastereoselectivity.

In conclusion, the linkage of 1-sulfonyl-1,3-butadiene and various acrylate dienophiles by a two-carbon tether bearing a large Ph or *t*-Bu substituent allowed efficient diastereose-



Figure 1. Proposed transition state for Diels-Alder cycloaddition of **14b-h**.

lective formation of the corresponding *endo*-IMDA cycloadducts. Our current efforts are directed toward the synthesis of enantiomerically pure cycloadducts and exploring their synthetic utility. Our results along these lines will be reported in due course.

Acknowledgment. We thank Kent State University for financial support and Yehor Novikov for valuable discussions. M.Z. was supported by NSF Grant 0111511, and the diffractometer was funded by NSF Grant 0087210, by Ohio Board of Regents Grant CAP-491, and by Youngstown State University.

Supporting Information Available: Full experimental procedures and spectroscopic data for compounds 3–10, 14–17, and 18g, along with copies of ¹H and ¹³C NMR spectra of major cycloadducts 15a–h, minor cycloadducts 16e,f, and mixtures 17a/15a and 16/15(b,c,g). This material is available free of charge via the Internet at http://pubs.acs.org. OL050930T