

TETRAHEDRON LETTERS

Stereoselective Reductive Coupling of Carbonyl Compounds by a Sm/Et2All System

Yutaka Nishiyama*, Eiji Shinomiya, Shouji Kimura, Kazuyoshi Itoh, and Noboru Sonoda

Department of Applied Chemistry, Faculty of Engineering & High Technology Research Center, Kansai University, Suita, Osaka 564, Japan

Received 26 January 1998; accepted 6 March 1998

Abstract: A new method for generation of the samarium (II) species by the reaction of samarium metal with diethylaluminium iodide (Et_2AII) under mild conditions has been developed. Divalent samarium species generated *in situ* from a Sm/Et₂AII system caused the reductive coupling of aromatic ketones with moderate stereoselectivity to afford corresponding *dl-vic*-diols in moderate yields. © 1998 Elsevier Science Ltd. All rights reserved.

The reductive coupling of aldehydes and ketones to glycols (pinacol coupling) is a powerful method for constructing carbon-carbon bonds,¹ and it can be accomplished with reagents such as Mg metal² or various low-valent complexes of Ti,³ Zr,⁴ V,⁵ Nb,^{5b,6} Sn,⁷ Sm⁸ or Ce.⁹ Recently, the stereochemistry of pinacol coupling has been of great interest due to the creation of two new adjacent stereocenters during the same stage. Although numerous reports on the stereoselective intramolecular pinacol couplings have appeared, ^{1e,10} the intermolecular coupling is fairly uncommon.^{3a,3b,11} In particular, there are few reports on highly stereoselective intramolecular reductive coupling of carbonyl compounds using a low-valent samarium species, which is an excellent reagent of pinacol coupling of carbonyl compounds.^{12,13}

We have recently disclosed that the samarium (II) species was easily prepared by the treatment of metallic samarium with organohalosilanes, R_nSiX_{4-n} (X=Br and I), under ambient conditions.¹⁴ In this Sm/ R_nSiX_{4-n} system, the reactivity of the low valent samarium species was affected by the organohalosilanes used (e.g., yield of hydrobenzoin, reductive coupling product of benzaldehyde, 1^{4a} , 1^{4b} and the selectivity of the azo, nitroso, azoxy, and amino products, during the reduction of aromatic nitro compounds^{14c}). From these results on the reactions using the Sm/ R_nSiX_{4-n} system, it is interesting to note that the use of other organometallic reagents instead of organohalosilanes may cause some exchange during the reactions.

In the course of our studies on the development of new preparative methods of the divalent samarium species, we found that the dark green solution similar to the THF solution of SmI₂ obtained by Kagan's method could be easily prepared by the reaction of metallic samarium with diethylaluminum iodide (Et₂AlI) (1). From the studies of the utility of this samarium species in organic reactions, we found that the reductive dimerization of ketones proceeded with high stereoselectivity (Eq. 1).

Sm + Et₂All
$$-$$
 CH₃CN, 50 °C, 3h 0 °C, 30min Ptt Me OH (Eq. 1)

0040-4039/98/\$19.00 © 1998 Elsevier Science Ltd. All rights reserved. PII: S0040-4039(98)00541-3

In a representative procedure the samarium (II) species was easily generated as follows: samarium powder (75 mg, 0.5 mmol) and acetonitrile (2.5 mL) were placed in a three-necked flask. A toluene solution of diethylaluminum iodide (1.5 mL of a 1.0 M solution, 1.5 mmol) was slowly added at 0 °C, and the mixture was stirred at 50 °C for 3 h under a nitrogen atmosphere. The samarium metal was gradually dissolved, and the color of the solution changed to dark green. When acetophenone (60 mg, 0.5 mmol) (2) was added to the resulting dark green solution at 0 °C, 2,3diphenylbutane-2,3-diol (3) was formed in 61 % yield with high stereoselectivity (dl: meso = 4.5 : 1.0) (Run 3 in Table 1).¹⁵ The yield and stereoselectivity of the diol 3 were markedly affected by the reaction temperature (Runs 2-5).16 The maximum yield was obtained at 0 °C, while



Sm⊣	+ Et₂All (1) 50 °C	Ph (0.5 CN 30 ml) , 3h	HO >h	
Run	Sm/Et ₂ All mmol / mmol	Temp/°C	Yield/% ^a	Selectivity ^b <i>dl / meso</i>
1	0.5 / 1.0	0	28 %	
2	0.5 / 1.5	-40	52 %	1.6 : 1.0
3	0.5 / 1.5	0	61 %	4.5 : 1.0
4	0.5 / 1.5	25	30 %	5.3 : 1.0
5	0.5 / 1.5	50	27 %	5.6 : 1.0
6	0.5 / 2.0	0	18 %	

^a Isolated yield. ^b Determined by ¹H NMR.

higher temperatures (25 and 50 °C) led to a decrease in the yield of 3 due to the formation of various byproducts. The highest stereoselectivity was observed at 50 °C (Run 5). The ratio of the organoaluminum reagent to the samarium metal also affected the yield of 3 (Runs 1, 3 and 6). When the samarium species generated from the samarium metal and two equivalents amount of Et 2AlI were reacted with 2, pinacolic product 3 was formed in an unsatisfactory yield (Run 1). In the case of the Sm/Et 2AlI (1/4 mmol) reagent, the color of the solution changed to yellow which is the characteristic color of Sm(III), and the yield of the coupling product decreased (Run 6).

The results of the reactions of various carbonyl compounds with the divalent samarium species generated *in situ* from Sm/Et₂All are shown in Table 2. In the reaction of 4-methylacetophenone, a decrease in the **Table 2**. Reductive Coupling of Various Carbonyl Compounds by Sm/Et₂All Reagent^a

Run	Substrate	Yield/% ^b	Selectivity ^c dl / meso	Run	Substrate	Yield/% ^b	Selectivity ^c dl / meso
	x i			5		43	1.2 : 1.0
1	: X = 4-CH ₃	63	3.8 : 1.0	6 ^e	\sim	trace	_
2 ^d	X = 4-Cl	60	1.3 : 1.0		, Ö		
3 ^d	X = 4-0CH ₃	71	4.8 :1.0	7	П Н	68	1.0 : 1.3
4	X = 2-CH ₃	45	14.1: 1.0				

^a Carbonyl compound (0.5 mmol) was reacted with Sm (0.5 mmol) species generated *in situ* from Sm (0.5 mmol)/Et₂All (1.5 mmol) at 0 °C for 30 min. ^b Isolated yield. ^c Determined by ¹H NMR.

^d Sm (0.75 mmol)/Et₂All (2.25 mmol) was used. ^e The reaction was carried out for 24 h.

stereoselectivity was not observed; however, the substitution of a chloro or methoxy group for hydrogen in the para position reduced the stereoselectivity of the products (Runs 1-3). In the case of 2-methylacetophenone, the highest *dl* selectivity was obtained (dl : meso = 14.1 : 1.0) (Run 4). However, non-aryl ketones such as 2-pentanone did not couple at an appreciable rate (5% after 24 h) (Run 6). The reaction of aromatic aldehydes such as benzaldehyde with the samarium (II) species under the same reaction conditions as that of acetophenone led to a mixture of stereoisomers (dl : meso = 1.0 : 1.3) in 68 % yield (Run 7).

Although we can not explain the reason for the high stereoselectivity at the present time, the stereoselectivity has been rationalized by considering the chelation model 4 in which samarium or aluminum is coordinated to both oxygen atoms (Scheme 1).¹⁸

Scheme 1 $Ph Me \xrightarrow{Sm / Et_2All} \begin{bmatrix} 0 - ML_n \\ Ph OH \\ Ph OH \\ Ph OH \\ HO Ph H \\ (4) \end{bmatrix} \xrightarrow{H^+} Ph H OH \\ HO Ph H \\ (4) \\ d/$

The results described in this paper show that the Sm/Et₂AlI system is not only a new generation method of the Sm (II) species but also a useful method for the highly stereoselective pinacol coupling of ketones. Further application of this reaction system in organic reactions is now in progress.

REFERENCES AND NOTES

- For reviews, see: (a) Wirth, T. Angew. Chem., Int. Ed. Engl. 1996, 35, 61. (b) Dushin, R. G. In Comprehensive Organometallic Chemistry II; Hegedus, L. S., Ed.; Pergamon: Oxford, 1995; Vol 12, pp1071. (c) Furstner, A. Angew. Chem., Int. Ed. Engl. 1993, 32, 164. (d) Robertson, G. M. In Comprehensive Organic Synthesis; Trost, B. M., Ed.; Pergamon: Oxford, 1991; Vol 3, pp563. (e) McMurry, J. E. Chem Rev. 1989, 89, 1513. (f) Kahn, B. E.; Rieke, R. D. Chem. Rev. 1988, 88, 733.
- Furstner, A.; Csuk, R.; Rohrer, C.; Weidmann, H. J. Chem. Soc., Perkin Trans. 1 1988, 1729 and references therein.
- (a) Lipski, T. A.; Hilfiker, M. A.; Nelson, S. G. J. Org. Chem. 1997, 62, 4566. (b) Barden, M. C.; Schwartz, J. J. Am. Chem. Soc. 1996, 118, 5484 and references therein.
- 4. Askham, F. R.; Carroll, K. M. J. Org. Chem. 1993, 58, 7328.
- (a) Hirao, T.; Hasegawa, T.; Maguruma, Y.; Ikeda, I. J. Org. Chem. 1996, 61, 366. (b) Kammermeier, B.; Beck, G.; Jacobi, D.; Jendralla, H. Angew. Chem., Int. Ed. Engl. 1994, 33, 685. (c) Park, J.; Pederson, S. F. Tetrahedron 1992, 48, 2069. (d) Kempf, D. J.; Sowin, T. J.; Doherty, E. M.; Hannick, S. M.; Codavoci, L.; Henry, R. F.; Green, B. E.; Spanton, S. G.; Norbeck. D. W. J. Org. Chem. 1992, 57, 782. (e) Annunziata, R.; Benaglia, M.; Cinquini, M.; Cozzi, F.; Giaroni, P. Ibid. 1992, 57, 782. (f) Raw, A. S.; Pederson, S. F. Ibid. 1991, 56, 830. (g) Park, J.; Pederson, S. F. Ibid. 1990, 55, 5924. (h) Konradi, A. W.; Pederson, S. F. Ibid. 1990, 55, 4506. (i) Annunziata, R.; Cinquini, M.; Cozzi, F.; Giaroni, P. Tetrahedron Assymetry 1990, I, 355. (j) Takahara, P. M.; Freudenberger, J. H.; Konradi, A. W.; Pederson, S. F. Ibid. 1989, 30, 7177. (k) Freudenberger, J. H.; Konradi, A. W.; Pederson, S. F. J. Am. Chem. Soc. 1989, 111, 8014.

- (a) Szymoniak, J.; Besancon, J.; Moise, C. Tetrahedron 1994, 50, 2841. (b) Szymoniak, J.; Besancon, J.; Moise, C. Tetrahedron 1992, 48, 3867.
- 7. Hays, D. S.; Fu, G. C. J. Am. Chem. Soc. 1995, 117, 3867.
- 8. Akane, N.; Hayano, T.; H. Kusui, H.; Nishiyama, Y.; Ishii, Y. J. Org. Chem. 1994, 59, 7902 and references therein.
- 9. Imamoto, T.; Kusumoto, T.; Hatanaka, Y.; Yokoyama, M. Tetrahedron Lett. 1982, 23, 1353.
- (a) Raw, A. S.; Pedersen, S. F.; J. Org. Chem. 1991, 56, 830. (b) Guidot, J. P.; Le Gall, T.; Mioskowski, C. Tetrahedron Lett. 1994, 35, 6671. (c) Hays, D. S.; Fu G. C. J. Am. Chem. Soc. 1995, 117, 7283. (d) Kang, M.; Park, J.; Konradi, A. W.; Pedersen, S. F. J. Org. Chem. 1996, 61, 5528 and references therein.
- (a) Handa, Y.; Inanaga, J. Tetrahedron Lett. 1987, 28, 5717. (b) Raubenheimer, H. G.; Seebach, D. Chimia, 1986, 40, 12. (c) Grugel, C.; Neumann, W. P.; Sauer, J.; Seifert, P. Tetrahedron Lett. 1978, 2847. (d) Pons, J. -M.; Zahra, J. -P.; Santelli, M. Tetrahedron Lett. 1981, 22, 3965. (e) Seebach, D.; Daum, H. J. Am. Chem. Soc. 1971, 93, 2795. (f) Seebach, D.; Oei, H. -A.; Daum, H. Chem. Ber. 1977, 110, 2316. (g) Clerici, A.; Porta, O. Tetrahedron 1982, 38, 1293. (h) Stocker, J. H.; Kern, D. H. J. Org. Chem. 1968, 33, 291. (i) Stocker, J. H.; Kern, D. H. J. Org. Chem. 1968, 33, 1270. (j) Bewick, A.; Brown, D. J. J. Chem. Soc. Perkin Trans 2, 1977, 99. (k) Bewick, A.; Cleghprn, H. P. J. Chem. Soc. Perkin Trans 2, 1973, 1410.
- H. B. Kagan has shown that the reaction of acetophenone with SmI2 in the presence of NiI2 proceeded with moderate stereoselectivity (*meso* : *dl* = 1.0 : 3.8). However, the scope and limitation of this reaction were not described in this paper : Namy, J. L.; Kagan, H. B. *Tetrahedron*, 1996, 52, 14225.
- 13. M. Uemura has recently reported the highly *threo*-selective pinacol coupling of tricarbonylchromium complexes of benzaldehyde with SmI₂ : Taniguchi, N.; Kaneta, N; Uemura, M. J. Org. Chem. **1996**, 61, 6088.
- a) Akane, N.; Kanagawa, Y.; Nishiyama, Y.; Ishii, Y. Chem. Lett., 1992, 2431. b) Nishiyama, Y.; Hatano, T.; Emoto, T.; Muromachi, A.; Ishii, Y. Technology Reports of Kansai Univ., 1995, 38, 113.
- 15. The ratio of *meso* and *dl* isomers was determined by ¹H NMR See, Agahiglam, H.; Moraveck, J. F.; Gauthier, H. Can. J. Chem., **1963**, 41, 194.
- 16. To clarify the features of the Sm/Et2AlI system, the reductive coupling of acetophenone by Sm (II) species generated by other methods (Sm/CH2ICH2I¹⁷ and Sm/Me3SiCl/NaI¹⁴) were investigated. However, these methods yielded an approximately 1 : 1 mixture of *meso* and *dl*-diol [*meso* : *dl* = 1.0 : 1.0 (Sm/CH2ICH2I) and 1.5 : 1.0 (Sm/Me3SiCl/NaI)] in contrast with predominant *dl* selectivity in the case of the Sm/Et2AlI system.
- 17. Girard, P.; Namy, J. L.; Kagan, H. B. J. Am. Chem. Soc. 1980, 102, 2693.
- 18. The high stereoselective pinacolic coupling of aldehydes by titanium or vanadium compounds has already reported. The *dl*-selectivity of the products was explained by the chelation model 5 where metal binded with two oxygens of the carbonyl group.

