



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

A NOVEL AND CONVENIENT METHOD FOR THE SYNTHESIS OF PHENSTATIN

Maojiang Wu^a, Qinggang Ji^a, Chunhao Yang^a & Yuyuan Xie^a

^a State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, SIBS, Chinese Academy of Sciences, Graduate School of the Chinese Academy of Sciences, 555 Zu Chong Zhi Road, Shanghai, 201203, PRCHINA E-mail:

Version of record first published: 06 Feb 2009.

To cite this article: Maojiang Wu, Qinggang Ji, Chunhao Yang & Yuyuan Xie (2005): A NOVEL AND CONVENIENT METHOD FOR THE SYNTHESIS OF PHENSTATIN, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 37:3, 272-275

To link to this article: <http://dx.doi.org/10.1080/00304940509354958>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

5. Kodak, *Soc. Anon Belg.* 595,533 (1960); [*Chem. Abstr.*, **55**, 14140i (1961)].
6. J. Calcines, H. M. Dolan, K. R. Falkensterns, C. D. Kellys, E. P. Murphys, R. L. Klinger, B. E. Kahn, A. G. Difrancesco and R. K. Hailstone, *J. Imaging Science & Technology*, **46**, 101 (2002).
7. K. Abe and T. Sakurai, *Japan Kokai* 77 59 782 (1977); *Chem. Abstr.*, **87**, 137245z (1977).
8. C. Bouillon and J. Maignan, *Chem. Abstr.*, **94**, 46780y (1981),
9. C. A. Audeh, *US Patent Ger. Offen.*, 3,002,878, 4,759,873 (1988); *Chem. Abstr.*, **109**, 131946e (1988).
10. (a) R. Rosen and E. E. Reid, *J. Am. Chem. Soc.*, **44**, 634 (1922); (b) G. M. Bennett and E. M. Whincop, *J. Chem. Soc.*, **120**, 422 (1921).
11. J. E. Shaw, *US Patent*, 5,162,585 (1992); *Chem. Abstr.*, **118**, 41198u (1993).
12. B. N. Acharya, H. Rana, J. Acharya and M. P. Kaushik, *J. Liquid Chrom. and Rel. Technol.*, **27**, 2945 (2004).

A NOVEL AND CONVENIENT METHOD FOR THE SYNTHESIS OF PHENSTATIN


Submitted by Maojiang Wu, Qinggang Ji, Chunhao Yang* and Yuyuan Xie
(02/28/05)

State Key Laboratory of Drug Research
Shanghai Institute of Materia Medica, SIBS
Chinese Academy of Sciences
Graduate School of the Chinese Academy of Sciences
555 Zu Chong Zhi Road, Shanghai. 201203, P.R.CHINA
e-mail: chyang@mail.shcnc.ac.cn

Phenstatin (**4**), a combretastatin A-4 (CA-4) derivative designed from the CA-4 skeleton by replacement of the olefin group with a carbonyl group, was found to be a potent anti-tubulin agent by the Pettit's group and displays excellent anticancer and antimitotic activities comparable to those of CA-4.¹ It is used as standard for antitubulin test and is now under preclinical studies²⁻⁹ with the potential to be developed as a drug.^{10,11} Pettit's group reported the synthesis of phenstatin starting from 3-[(*tert*-butyldimethylsilyl)oxy]-4-methoxybenzaldehyde by

oxidation to the corresponding carboxylic acid, followed by conversion to the acid chloride and treatment with morpholine. Reaction of the amide with the lithium derivative prepared from 3,4,5-trimethoxybromobenzene and *tert*-butyllithium at -78°C followed by deprotection afforded phenstatin in 30% overall yield.¹ Another synthetic method was reported by Liou and coworkers starting from 3,4,5-trimethoxybromobenzene and 3-[(*tert*-butyldimethylsilyl)oxy]-4-methoxybenzaldehyde in 3 steps (no yields given).⁵

This paper reports a new approach to **4** in a more economical and efficient route (60% overall yield) as shown in *Scheme 1*, in which the rarely used¹² but easily cleaved chloroacetyl function was selected as the protecting group of guaiacol and using polyphosphoric acid (PPA) as condensing agent.



Scheme 1

EXPERIMENTAL SECTION

Melting points were determined on a Büchi 510 melting apparatus, and are not corrected. ^1H NMR spectra were recorded on Bruker-400 NMR spectrometer. Microanalyses were carried out on a Leco CHN-2000 Elemental Analyzer. Mass spectra were obtained on a MAT-95 spectrometer. All reagents were commercial products from China National Medicines Group Shanghai Chemical Reagents Company.

Preparation of 2-Methoxyphenyl Chloroacetate (2).— Guaiacol (10 g, 0.08 mol) and chloroacetyl chloride (13 g, 0.115 mol) were placed into a 100 mL flask. The mixture was heated to 135°C and refluxed for 4 hours¹³ with stirring, then cooled to r.t. Cold water (10 mL) was added carefully to react with excess acid chloride. The mixture was extracted with CH_2Cl_2 (3 x 30 mL). The combined organic phase was washed sequentially with water, saturated NaHCO_3 , brine, then dried over anhydrous MgSO_4 . The solvent was evaporated in *vacuo* and yielded a yellow oil, which was dissolved in hot ethanol (25 mL) and recrystallized to afford 11.5 g (71%) of pure **2**, mp $58-60^{\circ}\text{C}$ (*lit.*¹³ mp. 60°C). The mother liquor was purified by silica gel (eluted petroleum ether-ethyl acetate, 1: 10) to give another 3 g of **2**, resulting in an overall yield 90%. ^1H NMR (CDCl_3): δ 7.24 (1H, m, H-4), 7.07 (1H, dd, $J = 1.6$, $J = 8$, H-6), 6.98 (1H, dd, $J = 1.2$, $J = 8$, H-5), 6.96 (1H, m, H-3), 4.35 (2H, s, CH_2Cl), 3.83 (3H, s, $-\text{OCH}_3$).

Preparation of Phenstatin Chloroacetate (3).- 3,4,5-Trimethoxybenzoic acid (10 g, 0.047 mol) was mixed by means of a mechanical stirrer with 80 g of PPA¹⁴ at 80°C and stirred for 1 hour, then **2** (3 g, 0.015 mmol) was added the mixture was stirred for an additional 4 hours. It was then added slowly into ice-water with vigorous stirring. The brown solid which precipitated was collected and washed with water (50 mL) and saturated NaHCO₃ (80 mL). Recrystallization from ethanol afforded 2.5 g (42%) of solid, mp. 150°C. The alcohol mother liquor was purified by silica gel (eluted petroleum ether-ethyl acetate, 1:1) to give an additional 1.3 g of **3**, resulting in an overall yield of 65%. Unreacted 3,4,5-trimethoxybenzoic acid (5.0 g) could be recovered from the filtrate above by acification to pH 5. ¹H NMR (CD₃COCD₃): δ 7.80 (1H, dd, J_{6/2} = 2.1, J_{6/5} = 8.7, H-6), 7.63 (1H, d, J_{2/6} = 2.1, H-2), 7.31 (1H, d, J_{5/6} = 8.7, H-5), 7.05 (2H, s, H-2',6'), 4.61 (2H, s, -CH₂Cl), 3.96 (3H, s, 4, -OCH₃), 3.85 (6H, s, 3',5'-OCH₃), 3.81 (3H, s, 4'-OCH₃). MS (*m/z*, %): 396 (27, M+2), 395 (18, M+1), 394 (83, M), 318 (100), 303 (24), 287 (20), 275 (24), 195 (33), 151 (32), 124 (48), 109 (22).

Anal. Calcd for C₁₉H₁₉ClO₇: C, 57.80; H, 4.85. Found: C, 57.74; H, 4.78

Preparation of Phenstatin (4).- Compound **3** (3 g, 0.008 mol) and sodium acetate trihydrate (5 g, 0.036 mol), were dissolved in 30 mL CH₃OH and refluxed for 3 hours. The solvent was evaporated and the residue was washed with water, dried in air, to give 2.37 g (98%) white solid, mp. 150°C, *lit.*¹ mp. 150°C. ¹H NMR (CDCl₃): δ 7.44 (1H, d, J_{2/6} = 2.1, H-2), 7.39 (1H, dd, J_{6/2} = 2.1, J_{6/5} = 8.2, H-6), 7.03 (2H, s, H-2',6'), 6.92 (1H, d, J_{5/6} = 8.2, H-5), 5.70 (1H, s, -OH), 3.99 (3H, s, 4-OCH₃), 3.93 (3H, s, 4'-OCH₃) 3.89 (6H, s, 3',5'-OCH₃); MS (*m/z*, %): 318 (100, M), 303 (18), 287 (14), 275 (25), 195 (24), 151 (32).

Anal. Calcd for C₁₇H₁₈O₆: C, 64.14; H, 5.70. Found: C, 64.07; H, 5.76

REFERENCES

1. G. R. Pettit, B. Toki, D. L. Herald, P. Verdier-Pinard, M. R. Boyd, E. Hamel and R. K. Pettit, *J. Med. Chem.*, **41**, 1688 (1998).
2. G. R. Pettit, M. P. Grealish, D. L. Herald, M. R. Boyd, E. Hamel and R. K. Pettit, *J. Med. Chem.*, **43**, 2731 (2000).
3. J. P. Liou, C. W. Chang, J. S. Song, Y. N. Yang, C. F. Yeh, H. Y. Tseng, Y. K. Lo, Y. L. Chang, C. M. Chang and H. P. Hsieh, *J. Med. Chem.*, **45**, 2556 (2002).
4. G. R. Pettit, M. P. Grealish, M. K. Jung, E. Hamel, R. K. Pettit, J. C. Chapuis and T. M. Schmidt *J. Med. Chem.*, **45**, 2534 (2002).
5. J. P. Liou, J. Y. Chang, C. W. Chang, C. Y. Chang, N. Mahindroo, F. M. Kuo and H. P. Hsieh, *J. Med. Chem.*, **47**, 2897 (2004).
6. H. Prinz, Y. Ishii, T. Hirano, T. Stoiber, J. A. C. Gomez, P. Schmidt, H. Dussmann, A. M. B. X. J. H. M. Prehn, E. G. Gunther, E. Unger and K. Umezawa, *J. Med. Chem.*, **46**, 3382 (2003).

7. J. P. Liou, Y. L. Chang, F. M. Kuo, C. W. Chang, H. Y. Tseng, C. C. Wang, Y. N. Yang, J. Y. Chang, S. J. Lee and H. P. Hsieh, *J. Med. Chem.*, **47**, 4247 (2004).
8. H. P. Hsieh, J. P. Liou, Y. T. Lin, N. Mahindroo, J. Y. Chang, Y. N. Yang, S. S. Chern, U. K. Tan, C. W. Chang, T. W. Chen *et al.*, *Bioorg. Med. Chem. Lett.*, **13**, 101 (2003).
9. N. H. Nam, *Current Medicinal Chemistry*, **10**, 1697 (2003).
10. M. L. Brown, J. M. Rieger and T. L. MacDonald, *Bioorg. Med. Chem.*, **8**, 1433 (2000).
11. B. Vernon and S. Powell, WO 2004009127; *Chem. Abstr.*, **140**, 133864 (2004).
12. Z. S. Tong, T. S. Zhong and M. L. Huang, *Yaoxue xuebao*, **6**, 164 (1958); *Chem. Abstr.*, **53**, 14039 (1959)
13. H. V. Krannichfeldt, *Ber.*, **46**, 4018 (1913).
14. L. H. Klemm and G. M. Bower, *J. Org. Chem.*, **23**, 344 (1958).

**EXPLOITATION OF CO-OPERATIVE DIRECTED *ortho*-METALLATION (DoM) BY
1,3-RELATED –OMOM GROUPS IN THE DEVELOPMENT OF A FULLY REGIO-
CONTROLLED SYNTHESIS OF ATRANOL FROM ORCINOL**

Submitted by Martin G. Banwell* and Satish Chand
(01/05/05)

*Research School of Chemistry, Institute of Advanced Studies
The Australian National University
Canberra, ACT 0200, AUSTRALIA
e-mail: mgb@rsc.anu.edu.au*

In connection with efforts to develop a total synthesis of the fungal metabolite diversionol¹ we required access to atranol (**5**, *Scheme 1*). The latter compound and its chloro derivative, both of which are powerful allergens, are themselves natural products and found as major components in oak moss absolute, an extract of the lichen *Everia prunatri* and an important ingredient in the perfumery industry.² Atranol also represents a key sub-structure associated with a wide-range of other lichen-derived natural products of polyketide origin. It has been prepared, albeit in very low yield, through Vilsmeier-Haack formylation of commercially available orcinol (1). The major product of this reaction is, in fact, the regioisomeric orcyraldehyde (**2**)³ which we