Synthesis and Characterisation of Porphyrazinoctamine Derivatives: X-Ray Crystallographic Studies of [2,3,7,8,12,13,17,18-Octakis(dibenzylamino)porphyrazinato]magnesium(II) and {2,3,7,8,12,13,17,18-Octakis[allyl(benzyl)amino]porphyrazinato}nickel(II)

Neelakandha S. Mani, ^a L. Scott Beall, ^b Todd Miller, ^a Oren P. Anderson, ^a Håkon Hope, ^c Sean R. Parkin, ^c David J. Williams, ^b Anthony G. M. Barrett*^b and Brian M. Hoffman*^d

^a Department of Chemistry, Colorado State University, Fort Collins, CO 80523, USA

Department of Chemistry, Imperial College of Science Technology & Medicine, London, UK SW7 2AY

• Department of Chemistry, University of Čalifornia, Davis, CA 95616, USA

^d Department of Chemistry, Northwestern University, Evanston, IL 60208, USA

2,3-Bis(dibenzylamino)- and 2,3-bis[allyl(benzyl)amino]-2Z-butene-1,4-dinitriles are macrocyclised by reaction with magnesium propoxide in propanol to provide the corresponding (porphyrazinato)magnesium(II) derivatives, the structures of which are confirmed by X-ray crystallographic studies of the title magnesium and nickel(II) complexes; cyclisation of alternative diaminomaleonitrile derivatives gives the corresponding porphyrazinoctamines.

Recently, we described the synthesis of the porphyrazine octathiolate derivative 2 by macrocyclisation of 2,3-bis-(benzylthio)maleonitrile 1 using magnesium propoxide in propanol, transmetallation and reductive debenzylation (Scheme 1).¹ The octathiolate 2 exhibited unusual coordination chemistry. Complexes were isolated with four metal cations peripherally bonded to the eight thiolate residues either via quadruple bidentate (S-S)² or tridentate (S-meso-N-S)¹ coordination. This peripheral multimetallation is in addition to metal-ion complexation within the porphyrazine cavity and thus provides novel pentametallic arrays. These complexes, denoted star porphyrazines, are of considerable interest for the study of indirect metal-metal interactions via the porphyrazine π -system. Herein, we report the synthesis of the analogous porphyrazinoctaamine derivatives. This novel† class of ligands is also expected to show rich and varied multimetallic complexation.

Commercial, inexpensive diaminomaleonitrile 3 was converted into the N, N, N', N'-tetraalkyl derivatives 4a, 4b and 5a-5c following the methods reported by Sheppard and coworkers. Thus, tetramethylation and benzylation of diamine 3 gave adducts 4a³ and 4b[‡] (53%). Dimethylation and diallylation of N, N'-dibenzyldiaminomaleonitrile³ (5, $\mathbb{R}^1 =$ Bn, $R^2 = H$) gave diamines 5a (64%) and 5c (80%) respectively. Finally, sequential double reductive alkylation³ of 3 using pyridine-2-carboxaldehyde and methylation gave dinitrile 5b (51%) (Scheme 2). All five dinitriles 4a, 4b and 5a-5c were readily cyclised using the magnesium propoxide method introduced by Linstead.⁴ Heating under reflux of propanol solutions of the dinitriles with magnesium propoxide gave the porphyrazines **6a-6e** (Scheme 3, Table 1). In each case, the porphyrazines were isolated and purified by chromatography on neutral alumina. All five porphyrazines were isolated as blue-black solids with purple reflections.

The magnesium complexes **6a**, **6c** and **6e** were demetallated by reaction with trifluoroacetic acid to produce porphyrazines **7a**, **7c** and **7e**. Subsequent reaction with nickel(π) acetate⁵ resulted in selective metallation within the macrocyclic cavity to provide the corresponding nickel complexes **8a**, **8c** and **8e**. Spectroscopic data were in full agreement with the proposed



formulations and the data for the octakis-dibenzylamino derivative **6c** are representative.§ Compound **6c** showed an electronic absorption spectrum with the Q band at 707 nm, a less intense peak at 574 nm and the Soret band⁶ at 368 nm. As the definitive confirmation of the structures, single crystal X-ray determinations were carried out on the magnesium porphyrazine **6c**¶ (Fig. 1) and the nickel porphyrazine **8e**∥ (Fig. 2), recrystallised from chlorobenzene–methanol and ethyl acetate–hexanes, respectively.

In this work we have outlined our initial studies on the synthesis of characterisation of a structurally novel class of macrocycles: 2,3,7,8,12,13,17,18-porphyrazineoctaamines.



Scheme 2 Reagents and conditions: i, R^1 CHO, MeOH; NaBH₄, THF, MeOH; R^1 CHO, MeOH; NaBH₄, THF, MeOH; R^2 X, NaH, THF; ii, R^1 X ($R^1 = R^2$), NaH, THF



Scheme 3 Reagents and conditions: i, $Mg(OPr)_2$, PrOH, heat; ii, CF_3CO_2H ; iii, $Ni(OAc)_2$, PhCl, DMF, heat

 Table 1^a Preparation of porphyrazines 6, 7 and 8

Entry	R ¹ , R ²	4 or 5 (%)	6(%)	7(%)	8(%)
1	Me, Me	4a (ref. 3)	6a (29, 48)	7a (21)	8a (43)
2	Bn, Me	5a (64)	6b (25)	` ´ ´	` ´
3	Bn, Bn	4b (53)	6c (31)	7c (86)	8c (59)
4	2-pvr. Me	5b (51)	6d (25)		<u> </u>
5	Bn, allyl	5c (80)	6e (20)	7e (82)	8e (92)

^a 2-pyr = 2-pyridylmethyl, yields of **5a** and **5c** based upon the known **5** ($R^1 = Bn, R^2 = H$, see ref. 3).



Fig. 1 X-ray crystal structure of [2,3,7,8,12,13,17,18-octakis(dibenzylamino)porphyrazinato]magnesium(II)



Fig. 2 X-ray crystal structure of {2,3,7,8,12,13,17,18-octakis[allyl-(benzyl)amino]porphyrazinato}nickel(u)

Clearly, the synthetic strategy is flexible and should be suitable for the synthesis of diverse analogues. Further studies on the preparation of related molecules and multimetallic complexation will be reported in due course.

We thank Glaxo Group Research Ltd., for the most generous endowment (to A. G. M. B.), the Wolfson Foundation for establishing the Wolfson Centre for Organic Chemistry in Medical Science at Imperial College, and the National Science Foundation (CHE-9107589 and DMR-9119832) for generous support of our studies.

Received, 1st June 1994; Com. 4/03173C

J. CHEM. SOC., CHEM. COMMUN., 1994

Footnotes

^{\dagger} Kopranenkov *et al.* have reported the preparation of [2,3,7,8,12,13,17,18-octakis(*tert*-butylamino)porphyrazinato]magnesium(11). However, these authors do not provide convincing data in support of their structural assignment. See ref. 7.

 ‡ All new compounds were fully characterised by spectroscopic data and microanalysis and/or HRMS. Porphyrazine was directly converted into the nickel complex 8c for characterisation.
 § Selected data for 6c: mp 185–190 °C (EtOAc-hexanes); TLC 0.86

§ Selected data for **6c**: mp 185–190 °C (EtOAc-hexanes); TLC 0.86 (EtOAc: hexanes 2:3); IR (CH₂Cl₂) v/cm⁻¹ 2957, 1450, 1430, 1314, 1285, 1233, 944, 881, 711, 675, 655; UV-VIS (CH₂Cl₂) $\lambda_{max}/nm (\log \epsilon)$ 368 (4.75), 574 (4.45), 707 (4.51); ¹H NMR (300 MHz, CDCl₃) δ 7.3 (m, 32H), 7.1 (m, 48H), 5.25 (s, 32H); ¹³C NMR (75 MHz, CDCl₃) δ 152.5, 140.3, 137.7, 129.1, 128.4, 126.9, 55.3; HRMS (FAB) *m/z* calc. for C₁₂₈H₁₁₂MgN₁₆; (M + H⁺), 1897.9185. Found: (M + H⁺), 1897.9124. Satisfactory elemental analysis was obtained.

¶ Crystal data for 6c: $\dot{C}_{138}H_{134}N_{16}O_2Mg$, M = 2072.92, monoclinic, a = 18.939(4), b = 27.813(6), c = 22.582(5) Å, $\beta = 99.56(3)^\circ$, V = 11730(4) Å³, space group $P2_1/n$, Z = 4, $D_c = 1.17$ g cm⁻³, μ (Cu-K α) = 0.60 mm⁻¹, F(000) = 4408. Data were measured on a Siemens rotating anode crystallographic system at the University of California, Davis, with Cu-K α radiation (Ni filter). The structure was solved with the direct methods program SHELXS-90⁸ and refined (all non-H atoms anisotropic) by full-matrix least-squares analysis on F^2 (all data) by using the program SHELXL-93,⁹ to give R = 0.074 (all data, on F) and wR = 0.229 (all data, on F^2) for the 14359 unique reflections measured. A disordered molecule of solvent was situated about the origin, but the disorder was so severe that a chemically sensible model for this occluded molecule could not be obtained.

|| Crystal data for 8e: C₉₆H₉₆N₁₆Ni·C₆H₅Cl, M = 1645.1, monoclinic, a = 17.453(2), b = 10.634(2), c = 25.256(3) Å, $\beta = 109.11(2)^\circ$, V = 4429 Å³, space group $P2_1/n$, Z = 2 (the molecule has crystallographic C_1 symmetry), $D_c = 1.23$ g cm⁻³, μ (Cu-K α) = 10.4 cm⁻¹, F(000) = 1740. A clear prism of dimensions 0.07 × 0.40 × 0.60 mm was used. Data were measured on a Siemens P4/PC diffractometer with Cu-K α radiation (graphite monochromator) using ω -scans. The structure was solved by direct methods and the non-hydrogen atoms refined anisotropically to give R = 0.062, $R_w = 0.065$ for 4759 independent observed reflections $[|F_o| > 4\sigma(|F_o|), 2\theta ≤ 120^\circ]$. The structure contains two pairs of disordered and partially overlapping 0.25 occupancy C₆H₅Cl molecules per porphyrazine unit.

Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre. See Information for Authors, Issue No. 1.

References

- C. S. Velázquez, W. E. Broderick, M. Sabat, A. G. M. Barrett and B. M. Hoffman, J. Am. Chem. Soc., 1990, 112, 7408; C. S. Velázquez, G. A. Fox, W. E. Broderick, K. A. Andersen, O. P. Anderson, A. G. M. Barrett and B. M. Hoffman, J. Am. Chem. Soc., 1992, 114, 7416.
- 2 C. S. Velázquez, T. F. Baumann, M. M. Olmstead, H. Hope, A. G. M. Barrett and B. M. Hoffman, J. Am. Chem. Soc., 1993, 115, 9997.
- 3 R. W. Begland, D. R. Hartter, F. N. Jones, D. J. Sam, W. A. Sheppard, O. W. Webster and F. J. Weigert, *J. Org. Chem.*, 1974, 39, 2341.
- 4 R. P. Linstead and M. Whalley, J. Chem. Soc., 1952, 4839.
- 5 J. W. Buchler, in *The Porphyrins*, ed. D. Dolphin, Academic, New York, 1978, vol. 1, ch. 10.
- 6 M. Gouterman, in *The Porphyrins*, ed. D. Dolphin, Academic, New York, 1978, vol. 2, ch. 1.
- 7 V. N. Kopranenkov, L. S. Goncharova and E. A. Luk'yanets, J. Org. Chem. USSR, 1979, 15, 962.
- 8 G. M. Sheldrick, Program SHELXS-90, Universität Göttingen, 1990.
- 9 G. M. Sheldrick, Program SHELXS-93, Universität Göttingen, 1993.