


Regioselective Ortho Lithiation of 3-Aryl
and 3-Styryl FuransMaria Tofi, Thomas Georgiou, Tamsyn Montagnon, and
Georgios Vassilikogiannakis*Department of Chemistry, University of Crete, 71409 Iraklion, Crete, Greece
vasil@chemistry.uoc.gr

Received May 23, 2005

ABSTRACT



An unusual regioselectivity pattern for the ortho lithiation of 3-aryl and 3-styryl furans has been uncovered wherein lithiation occurs preferentially at the sterically encumbered 2-position. The results are attributed, at least in part, to stabilization of the intermediate furyl anion by through-space donation of π -electron density from the substituent appended at the 3-position to the lithium cation. This ortho lithiation reaction may be applied as a useful synthetic tool for accessing 2,3-disubstituted furans.

The large number of natural and designed furans that constitute synthetic targets indicates that the ability to construct variously substituted furans¹ at will is a highly sought after skill. In this paper, we report the results of a study investigating the selective ortho lithiation of 3-aryl and 3-styryl furans at the 2-position, and, thereby, we introduce a method that could assist those seeking to synthesize 2,3-substituted furans that are relatively difficult to access by other means.

Heteroatom-directed lithiations at the 2-position of 3-substituted furans² (including 3-furoic acids,³ 3-arenesulfonyl furans,⁴ 3-furanmethanols,⁵ 3-furanmethanol derivatives⁶ and 3-bromofurans⁷) are known in the literature. The intramolecular [1,4] O→C silyl migrations of 3-(silyloxy)methyl furans and trialkylsilyl esters of 3-furoic acids to provide

2-trialkylsilyl-3-(hydroxymethyl) furans and 2-trialkylsilyl-3-furoic acids, respectively, are also known.⁸

On the other hand, the ortho lithiation of 3-alkylfurans is reported to afford a mixture of 2,3- and 2,4-disubstituted furans with the latter always being the major product,⁹ presumably for steric reasons. In most cases, the reported ratio of 2,4- vs 2,3-disubstituted furans varied between from 2:1 to 3:1.

During our studies¹⁰ toward the synthesis of a family of marine secondary metabolites, the prunolides,¹¹ we attempted to silylate 3-(4-methoxyphenyl)furan (**3a**) under standard ortho lithiation conditions (Scheme 1B). Despite our initial expectations for the predominant silylation occurring at the less sterically hindered 5-position, this reaction afforded

(1) For a review in regioselective syntheses of substituted furans, see: Hou, X. L.; Cheung, H. Y.; Hon, T. Y.; Kwan, P. L.; Lo, T. H.; Tong, S. Y.; Wong, H. N. C. *Tetrahedron* **1998**, *54*, 1955–2020.

(2) Gschwend, H. W.; Rodriguez, H. R. *Org. React.* **1979**, *26*, 1.

(3) (a) Knight, D. W.; Knott, A. P. *J. Chem. Soc., Perkin Trans. 1* **1981**, 1125. (b) Beese, G.; Keay, B. A. *Synlett* **1991**, 33.

(4) McCobie, S. W.; Shankar, B. B.; Ganguly, A. K. *Tetrahedron Lett.* **1987**, *28*, 4123–4126.

(5) (a) Goldsmith, D.; Liotta, D.; Saindane, M.; Waykole, L.; Bowen, P. *Tetrahedron Lett.* **1983**, *24*, 5835–5838. (b) Tanis, S. P.; Head, D. B. *Tetrahedron Lett.* **1984**, *25*, 4451–4454.

(6) Honda, T.; Kobayashi, Y.; Tsubuki, M. *Tetrahedron* **1993**, *49*, 1211–1222.


(7) (a) Ly, N. D.; Schlosser, M. *Helv. Chim. Acta* **1977**, *60*, 2085. (b) Weyerstahl, P.; Schenk, A.; Marschall, H. *Liebigs Ann. Chem.* **1995**, 1849.

(8) (a) Bures, E.; Spinazzé, P. G.; Beese, G.; Hunt, I. R.; Rogers, C.; Keay, B. A. *J. Org. Chem.* **1997**, *62*, 8741–8749. (b) Bures, E. J.; Keay, B. A. *Tetrahedron Lett.* **1988**, *29*, 1247–1251. (c) Bures, E. J.; Keay, B. A. *Tetrahedron Lett.* **1987**, *28*, 5965–5968.

(9) (a) Kernan, M. R.; Faulkner, D. J. *J. Org. Chem.* **1988**, *53*, 2773–2776. (b) Wang, E. S.; Yuen, M. C.; Wong, H. N. C. *Tetrahedron* **1996**, *52*, 12137–12158.

(10) Sofikiti, N.; Tofi, M.; Montagnon, T.; Vassilikogiannakis, G.; Stratakis, M. *Org. Lett.* **2005**, *7*, 2357–2359.


(11) (a) Carroll, A. R.; Healy, P. C.; Quinn, R. J.; Tranter, C. J. *J. Org. Chem.* **1999**, *64*, 2680–2682. (b) Ortega, M. J.; Zubía, E.; Ocaña, J. M.; Naranjo, S.; Salvá, J. *Tetrahedron* **2000**, *56*, 3963–3967.

Scheme 1. Regioselective Ortho Lithiation of 3-Arylfurans


mainly the 2-silylated product (**4a:5a** = 3:1).¹² To investigate the generality of this surprising and interesting observation we decided to scrutinize the result further by examining a range of substrates bearing different substituents at the para position of the appended phenyl ring (Scheme 1).


Akin to **3a**, substrates **3b** and **3c** were easily synthesized by employing a Suzuki coupling to unite 3-furanboronic acid (**1**) with the appropriate para-substituted bromobenzene (**2b,c**, Scheme 1A). Following the ortho lithiation/trimethylsilyl chloride quench sequence of these new substrates (**3b** and **3c**), it quickly became obvious that the more electron-withdrawing the substituent on the phenyl ring, the greater the enhancement was in the observed regioselectivity (in favor of the 2,3-disubstituted product, Scheme 1).

In an attempt to deconvolute the role played by conjugation, between the furan and the phenyl moieties, in securing the unusual regioselectivity that we had been observing, substrates **7a** and **7b** were prepared using a Suzuki-coupling protocol (Scheme 2). When the benzylfurans **7a** and **7b** were subjected to the established ortho lithiation/silylation condi-

Scheme 2. Ortho Lithiation of 3-Benzylfurans


tions, the 2,3- and 2,4-substituted furans **8a,b** and **9a,b** (respectively) were obtained in ratios of **8a:9a** = 1:1.3 (for the reaction of **7a**) and **8b:9b** = 1.1:1 (for the reaction of **7b**). These ratios, representing nearly equal substitution at the 2- and 5-positions of the furan substrate, not only differ significantly from those obtained for the previously tested conjugated substrates where the 2-position was favored but also differ slightly from the reported results for 3-alkylfurans⁹ where the 5-position was favored.


The next task we identified for our investigation was an examination of furan substrates where conjugation was achieved not by an aromatic substituent at the 3-position but by an olefin appended at this site instead. To this end, furan substrates **11** and **13** were synthesized, as shown in Scheme 3, using a combination of Wittig olefinations and a Suzuki

Scheme 3. Ortho Lithiation of 3-Alkenylfurans **11 and **13****


coupling. For substrates **11** and **13** the selectivity obtained in the ortho lithiation reaction was lower (ranging between 1.1 and 2.0:1) than we had previously observed with 3-arylfurans; however, once again, the more sterically hindered 2-position was favored (Scheme 3). Furthermore, and, once again, electron-donating groups decreased the selectivity (see (*E*)-**11** vs (*E*)-**13**). (*E*)-Substrates must be compared when we are talking about this electronic effect because, in the case of the (*Z*)-substrates, other factors come into play. On going from (*E*)-**11** to (*Z*)-**11**, the observed selectivity for the reaction at the 2-position dropped from 2.0:1 to 1.4:1 for steric reasons; however, on going from (*E*)-**13** to (*Z*)-**13**, the selectivity increased possibly due to the stabilizing influence of coordination of the methoxy group to the lithium cation (II, Figure 2).

Bearing all the results obtained thus far in mind, we decided next to examine the effect of extended conjugation at the 3-position of the furan ring. For this purpose, we targeted the 3-styrylfurans **17a–c** (both the (*E*)- and (*Z*)-isomers). These new substrates were easily prepared using standard Wittig protocols (Scheme 4). Crucially, careful

Scheme 4. Regioselective Ortho Lithiation of 3- β -Styrylfurans



chromatographic separation of the mixture of geometrical isomers obtained from the Wittig reactions allowed us access to pure samples of each isomer. Separate ortho lithiation, under our established conditions, of each of the (*E*)- or (*Z*)-configured isomers of **17a–c** afforded a set of interesting new results in which the selectivity in favor of the 2,3-disubstituted product was universally higher than had been recorded for any of our previously tested substrates (Scheme 4).

Within this latest set of results, a similar increase in the observed regioselectivity was recorded in the cases of the unsubstituted and *p*-CF₃-substituted substrate when compared to the *p*-OMe-substituted substrate ((*E*)-**17a** vs (*E*)-**17b** vs (*E*)-**17c**). The most impressive feature for this batch of results, however, was the high selectivity (for the 2,3-disubstituted product) obtained for the (*Z*)-substrates ((*Z*)-**17a–c**) in which the 2-position was severely sterically encumbered by the proximal phenyl ring (Scheme 4).

To investigate the postulate that conjugation of the 3-substituent (aryl, alkenyl, or styryl) with the furan is pivotal to the unexpected regioselectivity (of the ortho lithiation reaction of these substrates), two more unconjugated substrates were prepared (**20** and **21**, Figure 1). In 3-prenylfuran (**20**), the olefin was included as a potential through-space π -donor for the lithium cation in order to test whether an alternative interaction of this sort was important in producing the observed regioselectivity. Likewise, in 3-(2-phenyl)ethyl furan (**21**), the aryl group was included to act similar to a potential π -donor.

3-Prenylfuran (**20**; Figure 1) was synthesized by lithium halogen exchange (between 3-bromofuran and *n*-BuLi) followed by alkylation of the resulting furyl anion with prenyl-Br. 3-(2-Phenyl)ethyl furan (**21**), on the other hand,




Figure 1. Sterically governed ortho lithiation of 3-prenyl and 3-(2-phenyl)ethyl furans.

was easily prepared by hydrogenation (H₂, Pd/C) of **17b** (a mixture containing both geometrical isomers was used). In both cases, the standard ortho lithiation reaction (*n*-BuLi, TMSCl, -25 \rightarrow 25 $^{\circ}$ C) resulted in formation of the less sterically hindered 2,4-disubstituted furan in stark contrast to the results for the previously examined conjugated substrates. It had, therefore, become obvious that simple hydrogenation of the double bond of **17b** to furnish **21** produced a change in the regioselectivity in the subsequent ortho lithiation reaction by a factor of approximately 11 and led to the preferential formation of the opposite regioisomer.

At this point we should note that TMSCl was used as the quenching electrophile in all the ortho lithiation reactions for the purpose of directly comparing the results obtained. However, numerous other electrophiles may be employed to give access to a broad range of synthetically useful 2,3-disubstituted furans (starting from 3-aryl and 3-styryl furans). Further elaboration in the case of 2-substituted 3-styrylfurans might readily be achieved by functionalization (i.e., oxidative cleavage) of the double bond.

In an effort to further explain our ortho lithiation results, the $^1J(^{13}\text{C}-\text{H})$ coupling constants for both ortho positions of some representative substrates were measured. It is well-known that this coupling constant can be associated with the kinetic acidity of the corresponding hydrogens.¹³ It is important to note here that the full assignment of the relevant ^{13}C NMR peaks was accomplished using HMQC experiments. The results obtained from this NMR study are summarized in Table 1.

Table 1. $^1J(^{13}\text{C}-\text{H})$ for C-2 and C-5 of Selected Substrates

substrate	$^1J(^{13}\text{C}2-\text{H})$	$^1J(^{13}\text{C}5-\text{H})$
3a	200.1	202.1
3b	201.0	203.4
7a	199.5	201.3
(<i>E</i>)- 17b	201.2	202.2
(<i>Z</i>)- 17b	201.2	202.2

The $^1J(^{13}\text{C}-\text{H})$ coupling constants measured reveal that for substrates **3a**, **3b**, and **7a**, the 5-position hydrogen would

(12) All product ratios reported in Schemes 1–4 were measured by ^1H NMR of the crude reaction mixtures, while yields refer to isolated yields after chromatographic purification.

(13) Pedersen, E. B. *J. Chem. Soc., Perkin Trans. 2* **1977**, 473

appear to be more acidic than the other ortho hydrogen at the 2-position (the C-5 coupling constant is higher by ~ 2 Hz than the corresponding C-2 coupling constant). Despite this fact, the deprotonation of **3a** and **3b** occurs predominantly at the 2-position. In cases where the kinetic acidity of the two ortho protons is closer (such as substrates (*E*)-**17b** and (*Z*)-**17b** with ~ 1 Hz difference) the observed regioselectivity in favor of the 2-position hydrogen is even higher. The kinetic acidity of the ortho hydrogens is not, therefore, by itself sufficient to explain the observed regioselectivity. However, an increase in the acidity of the 2-position proton compared to the position-5 proton (on going from conjugated substrates **3a** and **3b** to (*E*)-**17b** and (*Z*)-**17b**) is in accord with higher regioselectivity in favor of the 2-position. While the data also make it obvious (Table 1) that having conjugated substituents at the 3-position increases the kinetic acidity of the ortho furan protons (see **7a** vs **3a**, **3b**, and **17b**), electron-donating groups appended to the conjugated 3-substituent decrease the kinetic acidity (**3a** vs **3b**). The latter effect can easily be rationalized as being the result of destabilization of the furyl anion by the electron-donating substituent.

A plausible mechanistic explanation for how this complicated situation arises has to do with the presence of a stabilizing interaction between the π -electronic system of the conjugated 3-aryl, 3-alkenyl, and 3-styryl substituents (**I**, **III**, and **IV**, Figure 2) and the lithium cation. These through-space interactions, known as π -stacking interactions,¹⁴ are not included in the measured kinetic acidity. The stabilization of a late transition state in the deprotonation step, due to interactions of this nature, may be of key importance. For this hypothesis to be correct, we need to assume that the π -donating group of **21** is positioned at too great a distance to participate strongly in such a stabilizing coordination. Likewise, the 3-benzyl furans (**7a** and **7b**) and 3-prenylfuran (**20**) readily adopt different conformations (due to their extra degrees of rotational freedom) many of which do not facilitate the stabilizing coordination between the π -system and the lithium cation; thus, overall, the stabilization in these substrates is of reduced strength.

(14) Ma, J. C.; Dougherty, D. A. *Chem. Rev.* **1997**, *97*, 1303–1324.




Figure 2. Through-space stabilizing interactions.

In summary, we have revealed an unusual regioselectivity pattern in the ortho lithiation of 3-aryl and 3-styryl furans (lithiation at the sterically encumbered 2-position is favored over lithiation at the 5-position) that may be attributed to stabilization of the intermediate furyl anion by through-space donation of π -electron density from the substituent appended at the 3-position to the lithium cation. The means by which to access sterically encumbered 2,3-disubstituted furans in preference to the alternative 2,4-disubstituted furans that we have uncovered is a useful synthetic tool.

Acknowledgment. This project was cofunded by the European Social Fund and National Resources (B EPEAEK, Pythagoras Program). This research was also supported by a Marie Curie Intra-European Fellowship (T.M.) within the 6th European Community Framework Program and the European Science Foundation (Cost D-28 program). We thank Dr. Iain Simpson for helpful discussions.

Supporting Information Available: Experimental procedures and ^1H , ^{13}C , and GC-MS spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL051205L