

# Radical Transfer Hydrosilylation/ Cyclization Using Silylated Cyclohexadienes


Stephan Amrein, Andreas Timmermann, and Armido Studer\*

Fachbereich Chemie der Universität Marburg, Hans-Meerwein-Strasse,  
D-35032 Marburg

studer@mail.uni-marburg.de

Received May 23, 2001

## ABSTRACT



A new method for mild metal-free hydrosilylation is described. Silylated cyclohexadienes are used as radical transfer hydrosilylating reagents for various double and triple bonds. A trialkylsilane is transferred from a cyclohexadiene moiety to an alkene. The hydrosilylation can be combined with a C–C bond formation as shown for the preparation of silylated cycloalkanes from the corresponding dienes.

The hydrosilylation/cyclization reaction of dienes is a well-known process. These reactions are generally conducted using transition metals as catalysts. Cationic palladium complexes have been successfully used to catalyze these transformations.<sup>1</sup> Furthermore, neodymium<sup>2</sup> and more importantly yttrium<sup>3</sup> metallocene complexes catalyze these hydrosilylation/cyclization reactions.

From an ecological point of view, it would be highly desirable to conduct these reactions without using transition metals. However, there are only a few reports on *metal-free* hydrosilylation reactions. In general, radical hydrosilylation of alkenes cannot be conducted using trialkylsilanes. This is due to the rather strong Si–H bond in the trialkylsilanes.<sup>4</sup> The alkyl radical formed after initial silyl radical addition

onto the alkene is not reduced with a trialkylsilane. Modified silanes, such as tris(trimethylsilyl)silane, have been successfully used in radical hydrosilylations.<sup>5</sup> In addition, Roberts presented some examples using the concept of polarity reversal catalysis.<sup>6</sup>

Recently, we introduced silylated cyclohexadienes as new tin-free radical reducing reagents.<sup>7</sup> Various typical radical reactions, such as dehalogenations, deselanations, deoxygenations, and intermolecular additions, were performed using these new reagents. The cyclohexadiene CH<sub>2</sub> moiety acts as the H-donor in these radical chain reactions. Reduction of radical R<sup>2</sup> with **1** affords cyclohexadienyl radical **2**. Rearomatization of **2** then provides the corresponding silyl radical, which is able to propagate the chain by reaction with the starting halide, xanthate, or phenylselenide R<sup>2</sup>–X (Scheme 1). As byproduct, methylated resorcin diether **3** is formed. We conceived that the silylated cyclohexadienes can also be used as radical hydrosilylating reagents: If the silyl radical formed in the rearomatization of **2** is allowed to react with an alkene,  $\beta$ -silyl radical **4** will be generated. Reduction of **4** with the cyclohexadiene **1** would then afford **2** and the

(1) Widenhoefer, R. A.; DeCarli, M. A. *J. Am. Chem. Soc.* **1998**, *120*, 3805. Stengone, C. N.; Widenhoefer, R. A. *Tetrahedron Lett.* **1999**, *40*, 1451. Perch, N. S.; Widenhoefer, R. A. *J. Am. Chem. Soc.* **1999**, *121*, 6960. Pei, T.; Widenhoefer, R. A. *Org. Lett.* **2000**, *2*, 1469. Perch, N. S.; Pei, T.; Widenhoefer, R. A. *J. Org. Chem.* **2000**, *65*, 3836. Wang, X.; Chakrapani, H.; Stengone, C. N.; Widenhoefer, R. A. *J. Org. Chem.* **2001**, *66*, 1755.

(2) Onozawa, S.; Sakakura, T.; Tanaka, M. *Tetrahedron Lett.* **1994**, *35*, 8177.

(3) Molander, G. A. *Chemtracts* **1998**, *11*, 237 and references therein. Molander, G. A.; Dowdy, E. D. In *Topics in Organometallic Chemistry*; Kobayashi, S., Ed.; Springer-Verlag: New York, 1999; Vol. 2, pp 120–154.


(4) Chatgililoglu, C. *Chem. Rev.* **1995**, *95*, 1229.

(5) Chatgililoglu, C. *Acc. Chem. Res.* **1992**, *25*, 188.

(6) Roberts, B. P. *Chem. Soc. Rev.* **1999**, *28*, 25.


(7) Studer, A.; Amrein, S. *Angew. Chem., Int. Ed.* **2000**, *39*, 3080.

**Scheme 1.** Silylated Cyclohexadienes as Radical Reducing Reagents




hydrosilylation product **5**. Radical **2** will eventually fragment a silyl radical, thereby propagating the chain (Scheme 2).

**Scheme 2.** Silylated Cyclohexadienes as Radical Hydrosilylating Reagents



This hydrosilylation of the alkene can formally be regarded as a *transfer-hydrosilylation*, since reagent **1** is transformed to the corresponding arene in a reverse hydrosilylation process. The driving force of this *transfer-hydrosilylation* is the resonance energy of the arene. Herein, we report the first applications of the silylated cyclohexadienes in metal-free hydrosilylation reactions.


The synthesis of cyclohexadiene **6** has previously been described.<sup>7</sup> The dimethylphenyl silyl derivative **7**, the triisopropyl compound **8**, and cyclohexadiene **9** were prepared in analogy (Figure 1).



**Figure 1.** Silylated cyclohexadienes **6–9**.

Hydrosilylation of allyl acetate could easily be conducted with reagent **6** in hexane at 80–85 °C (sealed tube) using  $\alpha,\alpha'$ -azoisobutyronitrile (AIBN) as initiator. Best results were obtained with 1.5 equiv of the silylated cyclohexadiene and 0.3 equiv of AIBN. Under these conditions, the hydrosilylation product **10** was isolated in 54% yield (Scheme 3).


**Scheme 3.** Hydrosilylation of Various Double Bonds with Reagent **6**



Hydrosilylation of 4-phenyl-1-butene worked equally well and **11** was isolated in 55% yield. Terminal substituted double bonds can also be hydrosilylated, as shown for the reaction of cyclohexene with **6** to form tetraalkylsilane **12** (60%). Hydrosilylation of  $\beta$ -pinene afforded the ring-opened cyclohexene derivative **13** in 70% yield. The initially formed  $\beta$ -silyl radical undergoes fast ring opening to form a tertiary radical, which is subsequently reduced with reagent **6** to provide compound **13**.

We next studied the hydrosilylation of various dienes which are easily prepared as described in the Supporting Informations. In these reactions the initially formed  $\beta$ -silyl radical undergoes a 5-*exo* radical cyclization to afford the corresponding primary radical, which after reduction provides a silylated cycloalkane. Treatment of diene **14** with reagent **6** provided the hydrosilylation/cyclization product **19** in 80% yield as a 4.3:1 (*cis:trans*) mixture of diastereoisomers (Table 1). The diastereoselectivity was determined by GC analysis. The relative configuration of the major isomer was assigned by comparison of the NMR data with literature values.<sup>8</sup> All other new compounds were assigned in analogy. The predominant formation of the *cis*-isomer is in agreement with the Beckwith–Houk model for 5-*exo* cyclization reactions.<sup>9</sup>

(8) Miura, K.; Oshima, K.; Utimoto, K. *Bull. Chem. Soc. Jpn.* **1993**, 66, 2348.


**Table 1.** Hydrosilylation/Cyclization of Various Dienes


X	starting material	yield (%)	product	dr (cis:trans)
C(CO <sub>2</sub> Et) <sub>2</sub>	<b>14</b>	80	<b>19</b>	4.3 : 1
O	<b>15</b>	71	<b>20</b>	2.5 : 1
NTos	<b>16<sup>a</sup></b>	76	<b>21</b>	2.0 : 1
	<b>17<sup>b</sup></b>	62	<b>22</b>	2.3 : 1
	<b>18</b>	72	<b>23</b>	2.6 : 1

<sup>a</sup> Reaction took 7 h for completion. <sup>b</sup> 3 equiv of **6**, 0.6 equiv of AIBN.

Ether **15**, tosylamide **16**, and ketal **18** were readily transformed to the corresponding cycloalkanes **20**, **21**, and **23** in good yields with moderate selectivities. For tosylamide **16**, reaction took longer (7 vs 4 h). Under our standard conditions, hydrosilylation of diol **17** did not go to completion. However, with 3 equiv of Si reagent **6** and 0.6 equiv of initiator, high conversion was obtained and compound **22** was isolated in 62% yield (dr = 2.3:1).

Since the silyl group in the Si reagents can be readily varied (see Figure 1), basically any silyl group can be introduced via radical hydrosilylation using our approach. To demonstrate this fact, we studied the hydrosilylation of diene **24** using the Si reagents **6–9**. Hydrosilylation with reagent **6** under the standard conditions provided carbocycle **25** in 84% yield (Table 2). Hydrosilylation/cyclization with the bulkier dimethylphenylsilyl **7** and the triisopropylsilyl reagent **8** provided the corresponding products **26** (57%) and **27** (44%, not shown) in lower yields. However, upon using


**Table 2.** Hydrosilylation/Cyclization of Diene **24** with Si Reagents **6–9**


reagent	R	yield (%)	product	dr (cis:trans)
<b>6</b>	SiMe <sub>2</sub> t-Bu	84	<b>25</b>	3.8 : 1
<b>7</b>	SiMe <sub>2</sub> Ph	57	<b>26</b>	4.4 : 1
<b>8<sup>a</sup></b>	Si(i-Pr) <sub>3</sub>	82	<b>27</b>	2.6 : 1
<b>9<sup>a</sup></b>	SiMe <sub>3</sub>	81	<b>28</b>	3.1 : 1

<sup>a</sup> *t*-BuOO-*t*-Bu (0.5 equiv) was used as initiator at 140 °C.


di-*tert*-butylperoxide as initiator at 140 °C in hexane (sealed tube), high yields can be obtained even with these bulky reagents, as shown for the hydrosilylation/cyclization with reagent **8** (82%, see Table 2). It turned out that the trimethylsilyl derivative **9** lacking the methoxy substituents is slightly less reactive.<sup>10</sup> The hydrosilylation could not be initiated using AIBN. However, with di-*tert*-butylperoxide as initiator smooth reaction occurred in hexane at 140 °C and the hydrosilylation/cyclization product **28** was isolated in 81% yield.

We also studied hydrosilylation reactions comprising a 6-exo cyclization (Scheme 4). Reaction of diene **29** with

**Scheme 4.** Radical Hydrosilylation Comprising a 6-Exo Cyclization

reagent **6** afforded cyclohexane derivative **30** in 61% yield as 1:1 mixture of diastereoisomers.<sup>11</sup> As expected, the silyl radical added regioselectively at the unsubstituted terminal double bond of bisalkene **29**.

Hydrosilylation of bisalkyne **31** under our standard conditions (Scheme 5) afforded double hydrosilylation product **32**

**Scheme 5.** Hydrosilylation of Bisalkyne **31**

in 11% yield along with unreacted starting material and mono silylation/cyclization compound **33**.<sup>12</sup> Under optimized conditions (4 equiv of **6**, *t*-BuON=NOt-Bu as initiator),<sup>13</sup> 55% of **32** was isolated. Reduction of the allyl radical **34** formed

(9) Beckwith, A. L.; Schiesser, C. H. *Tetrahedron* **1985**, *41*, 3925. Spellmeyer, D. C.; Houk, K. N. *J. Org. Chem.* **1987**, *52*, 959.

(10) Amrein, S. Unpublished results.


(11) About 5% unreacted starting material remained.

(12) Ojima, I.; Zhu, J.; Vidal, E. S.; Kass, D. F. *J. Am. Chem. Soc.* **1998**, *120*, 6690.

after silyl radical addition to diene **33** occurred regioselectively at the less hindered site (secondary vs tertiary radical).

The hydrosilylation products, if appropriately substituted at silicon, can be readily oxidized to the corresponding alcohols using the well-known Tamao–Fleming oxidation protocol.<sup>14</sup> This certainly expands the utility of our new method. For instance, bisester **26** (*cis:trans* = 4:1) was transformed to alcohol **35** in 73% yield (*cis:trans* = 4:1, Scheme 6).<sup>15</sup>

**Scheme 6.** Tamao–Fleming Oxidation



In conclusion, we report a new method for radical hydrosilylation using silylated cyclohexadienes as *metal-free*

*transfer-hydrosilylating* reagents. Double as well as triple bonds can readily be hydrosilylated with the new method. 1,5- and 1,6-dienes undergo the hydrosilylation/cyclization reaction to afford silylated cycloalkanes. To the best of our knowledge, this is the first report of a transfer-hydrosilylation reaction.

**Acknowledgment.** We thank the Swiss Science National Foundation (2100-055280.98/1) for funding our work. The work described herein is part of the planned dissertation of S.A.

**Supporting Information Available:** Full experimental details and spectroscopic and analytical data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL016160C

(13) Mendenhall, G. D. *Tetrahedron Lett.* **1983**, 24, 451.

(14) Jones, G. R.; Landais, Y. *Tetrahedron* **1996**, 52, 7599.

(15) The same oxidation has already been reported: Widenhoefer, R. A.; Stengone, C. N. *J. Org. Chem.* **1999**, 64, 8681.