Total Synthesis of (-)-Discodermolide

Amos B. Smith, III,* Yuping Qiu, David R. Jones, and Kaoru Kobayashi

> Department of Chemistry Monell Chemical Senses Center, and Laboratory for Research on the Structure of Matter University of Pennsylvania Philadelphia, Pennsylvania 19104

Received August 28, 1995

In 1990 discodermolide emerged as an extremely promising immunosuppressive agent, comparable with FK506 and rapamycin.^{1,2} The exceptional pharmacological potential and extreme scarcity of the natural material [0.002% (w/w) from frozen marine sponge] have stimulated intensive synthetic effort,³ including a total synthesis of both the unnatural and natural antipodes [i.e., (-)- and (+)-1] by Schreiber and co-workers,⁴ which also elucidated the absolute stereochemistry.

From the retrosynthetic perspective, we and others recognized the repeating stereochemical triad embedded in the discodermolide backbone (Scheme 1). This observation suggested dissection of the skeleton into fragments A, B, and C, each to

Scheme 1



derive from a common precursor 2. Union of the fragments at the olefinic linkages and introduction of the terminal diene moiety would exploit a combination of stereocontrolled σ - and π -bond constructions, the former noteworthy as an innovative unifying strategem in our immunosuppressant synthetic program.⁵ Herein we report a highly convergent and stereoselective total synthesis of (-)-discodermolide (1).⁶

(1) (a) Schreiber, S. L. Science **1991**, 251, 283. (b) Rosen, M. K.; Schreiber, S. L. Angew. Chem., Int. Ed. Engl. **1992**, 31, 384. (2) Gunasekera, S. P.; Gunasekera, M.; Longley, R. E. J. Org. Chem.

1990, 55, 4912. Correction: Ibid. 1991, 56, 1346.

(3) For other synthetic approaches to discodermolide, see: (a) Paterson, (3) For other synthetic approaches to discodermolide, see: (a) Paterson,
I.; Wren, S. P. J. Chem. Soc., Chem. Commun. 1993, 1790. (b) Clark, D.
L.; Heathcock, C. H. J. Org. Chem. 1993, 58, 5878. (c) Golec, J. M. C.;
Jones, S. D. Tetrahedron Lett. 1993, 34, 8159. (d) Evans, P. L.; Golec, J.
M. C.; Gillespie, R. J. Ibid. 1993, 34, 8163. (e) Golec, J. M. C.; Gillespie,
R. J. Ibid. 1993, 34, 8167. (f) Yang, G.; Myles, D. C. Ibid. 1994, 35, 1313.
(g) Yang, G.; Myles, D. C. Ibid. 1994, 35, 2503. (h) Paterson, I.; Schlapbach, A. Synlett 1995, 498.

(4) Nerenberg, J. B.; Hung, D. T.; Somers, P. K.; Schreiber, S. L. J. Am. Chem. Soc. **1993**, 115, 12621. Hung, D. T.; Nerenberg, J. B.; Schreiber, S. L. Chem. Biol. 1994, 1, 67.

(5) (a) FK506 formal synthesis: Smith, A. B., III; Chen, K.; Robinson, D. J.; Laakso, L. M.; Hale, K. J. Tetrahedron Lett. 1994, 35, 4271. Also see: Smith, A. B., III; Hale, K. J. Ibid. 1989, 30, 1037. Smith, A. B., III; Hale, K. J.; Laakso, L. M.; Chen, K.; Riéra, A. Ibid. 1989, 30, 6963. (b) Rapamycin and demethoxyrapamycin total syntheses: Smith, A. B., III; Condon, S. M.; McCauley, J. A.; Leazer, J. L., Jr.; Leahy, J. W.; Maleczka, R. E., Jr. J. Am. Chem. Soc. 1995, 117, 5407. Also see: Smith, A. B., III; Condon, S. M.; McCauley, J. A.; Leahy, J. W.; Leazer, J. L., Jr.; Maleczka, R. E., Jr. Tetrahedron Lett. 1994, 35, 4907. Smith, A. B., III; Maleczka, R. E., Jr.; Leazer, J. L., Jr.; Leahy, J. W.; McCauley, J. A.; Condon, S. M. Ibid. 1994, 35, 4911.

Scheme 2



As our point of departure, ester (-)-3 (Scheme 2) was protected as the PMB ether;⁷ LAH reduction, Swern oxidation,^{8a} Evans syn aldol addition of (+)-5,9 and Weinreb transamidation¹⁰ then afforded the common precursor (+)-2.¹¹ The overall yield for this five-step sequence, conveniently executed on a 50-g scale, was 59%.

Elaboration of the C(15-21) fragment A began with DDQ oxidation^{8b} of (+)-2 (Scheme 3). LAH reduction gave aldehyde (-)-6,¹¹ which underwent syn aldol addition of (+)-5⁹ to furnish (+)-7;¹¹ the structure and stereochemistry were confirmed by single-crystal X-ray analysis.¹² Hydroxyl protection as the TBS ether, reductive (LiBH₄)¹³ removal of the chiral auxiliary without epimerization, and iodination via a variation of the Corey protocol¹⁴ [I₂, PPh₃, imidazole, PhH/Et₂O (1:2)] then afforded $(-)-A.^{11}$

Scheme 3



The C(9-14) fragment **B**, a vinylic halide initially conceived as the bromide 9, likewise derived from (+)-2, beginning with protection (TBS) and DIBAL reduction¹⁵ (Scheme 4). Treatment of the resultant aldehyde (+)- 8^{11} with the Wittig reagent $Ph_3PCBrCO_2Et^{16}$ gave predominantly (8.5:1) the expected Z unsaturated ester. A second DIBAL reduction, mesylation, and displacement with Super-Hydride then furnished vinyl bromide (-)-9.¹¹ After considerable experimentation, the corresponding vinyl iodide (-)-**B**¹¹ proved to be a more effective substrate for cross coupling (vide infra). The iodide was generated from (-)-9 with catalytic Zn, NiBr₂, and excess KI (HMPA/DMF,

(6) At the start of our work, the absolute stereochemistry was unknown. (7) Nakajima, N.; Horita, K.; Abe, R.; Yonemitsu, O. Tetrahedron Lett. 1988, 29, 4139.

(8) (a) Mancuso, A. J.; Swern, D. Synthesis 1981, 165. (b) Parikh, J. R.; Doering, W. von E. J. Am. Chem. Soc. 1967, 89, 5505. (c) Oikawa, Y.;

 Doering, W. Von E. J. Am. Chem. Soc. 1907, 89, 5305. (c) Okawa, Y.;
 Yoshioka, T.; Yonemitsu, O. Tetrahedron Lett. 1982, 23, 889.
 (9) Gage, J. R.; Evans, D. A. Org. Synth. 1990, 68, 77.
 (10) (a) Levin, J. I.; Turos, E.; Weinreb, S. M. Synth. Commun. 1982,
 12, 989. (b) Evans, D. A.; Bender, S. L.; Morris, J. J. Am. Chem. Soc. 1988, 110, 2506.

(11) All synthetic compounds were purified by flash chromatography on silica gel. The structure assigned to each new compound is in accord with its infrared, 500-MHz ¹H NMR, and 125-MHz ¹³C NMR spectra, as well as appropriate parent ion identification by high-resolution mass spectrometry. In addition, compounds (+)-2, (+)-7, (+)-10, (-)-14, and)-A gave satisfactory combustion analyses.

(12) Carroll, P. J., University of Pennsylvania, unpublished results.
(13) Penning, T. D.; Djuric, S. W.; Haack, R. A.; Kalish, V. J.; Miyashiro, J. M.; Rowell, B. W.; Yu, S. S. Synth. Commun. 1990, 20, 307.
(14) Corey, E. J.; Pyne, S. G.; Su, W.-G. Tetrahedron Lett. 1983, 24, 1000 4883. We obtained superior results with the less polar solvent system PhH/ Et₂O (1:2).

(15) Nahm, S.; Weinreb, S. M. Tetrahedron Lett. 1981, 22, 3815

(16) Boeckman, R. K., Jr.; Ko, S. S. J. Am. Chem. Soc. 1980, 102, 7146.

0002-7863/95/1517-12011\$09.00/0 © 1995 American Chemical Society

Scheme 4



90 °C, 86% yield);¹⁷ alternatively **B** could be prepared in one step from aldehyde (+)-8 (41%, 6:1 Z/E) via the procedure of Zhao.18

Preparation of the C(1-8) fragment C began with hydroxyl protection (TBSOTf) and removal of the PMB group in common precursor 2 [H₂, Pd(OH)₂/C, EtOH], followed by oxidation with $SO_3 \cdot py^{8c}$ and conversion to dithiane (+)-10^{11,19} (Scheme 5). DIBAL reduction, formation of the dimethoxy acetal, and addition of the derived dithiane anion to benzyl (R)-(-)-glycidyl ether (11) then furnished (-)-12.¹¹ The latter transformation manifests a central strategy of our immunosuppressant synthetic program, the stereospecific generation of protected aldol linkages via dithiane – epoxide coupling.⁵ Dithiane cleavage,²⁰ Evans directed reduction $[Me_4NBH(OAc)_3]$,²¹ and chemoselective cyclization led to the methyl pyranosides **13**.^{11,22} Silylation, debenzylation (H₂, Pd/C), ethyl thioacetal formation,²³ and Swern oxidation furnished (+)-C.¹¹ The stereochemical assignments for C(5) and the anomeric center derived from NMR decoupling and NOE studies of the penultimate alcohol.

Scheme 5



With fragments A, B, and C in hand, we turned to assembly of the discodermolide backbone. Initially we explored the cross coupling of fragment (-)-A with bromide (-)-9; under a variety of conditions, the highest yield of $(-)-14^{11}$ was only 14% [MgBr₂, PdCl₂(dppf)].²⁴ In contrast, reaction of vinyl iodide (-)-B with the organozinc derivative of (-)-A (prepared via the alkyllithium)²⁵ and Pd(PPh₃)₄ as catalyst²⁶ afforded (-)-14 in 66% yield (Scheme 6). Selective removal of the PMB group

- (17) Takagi, K.; Hayama, N.; Inokawa, S. Chem. Lett. 1978, 1435.
- (18) Chen, J.; Wang, T.; Zhao, K. Tetrahedron Lett. 1994, 35, 2827.
 (19) Evans, D. A.; Truesdale, L. K.; Grimm, K. G.; Nesbitt, S. L. J. Am. Chem. Soc. 1977, 99, 5009.
- (20) Stork, G.; Zhao, K. Tetrahedron Lett. 1989, 30, 287.
- (21) Evans, D. A.; Chapman, K. T.; Carreira, E. M. J. Am. Chem. Soc. 1988, 110, 3560.
- (22) The 2:1 mixture of O-methyl anomers, although separable, was carried forward; the anomeric ethyl thioacetals were separated.
 (23) Kim, S.; Park, J. H.; Lee, S. *Tetrahedron Lett.* 1989, 30, 6697.
 (24) Hayashi, T.; Konishi, M.; Kobori, Y.; Kumada, M.; Higuchi, T.;
- Hirotsu, K. J. Am. Chem. Soc. 1984, 106, 158.
- (25) To suppress TBS migration to the primary carbanion, the mixture of (-)-A and dried ZnCl₂ in Et₂O was cooled to -78 °C before treatment with t-BuLi.

[DDQ (1.05 equiv), CH₂Cl₂, H₂O] then provided the primary alcohol with less than 2% overoxidation of the benzylic acetal. Conversion to the corresponding iodide proved unexpectedly difficult; eventually our modification of the Corey protocol $[Ph_3P, I_2, imidazole, PhH/Et_2O(1:2)]^{14}$ did provide the unstable iodide,²⁷ which in turn furnished the requisite Wittig reagent (-)-15¹¹ upon heating (80 °C) with Ph₃P (15 equiv) and i-Pr₂NEt (3 equiv).

Scheme 6



Union of fragment (+)-C with the ylide derived from (-)-15 (Scheme 6) and reductive opening of the acetal ring (DIBAL) gave primary alcohol (-)-16,¹¹ which was oxidized (SO₃·py) to the corresponding aldehyde. Yamamoto olefination²⁸ then generated Z diene (-)-17,¹¹ comprising the complete discoder-molide backbone.²⁹ Hg(II)-mediated thioacetal hydrolysis followed by oxidation (DMSO, Ac_2O) introduced the C(1) lactone.³⁰ Completion of the discodermolide synthetic venture entailed removal of the PMB group (DDQ), carbamate introduction via the Kocovsky protocol (Cl₃CCONCO; imide hydrolysis),³¹ and final deprotection (HF, CH₃CN); synthetic (-)-1 was identical to (-)-discodermolide (500-MHz ¹H and 125-MHz ¹³C NMR, IR, HRMS, optical rotation, and TLC in four solvent systems).32

Acknowledgment. Support was provided by the National Institutes of Health (Institute of General Medical Sciences) through Grant GM-29028.

Supporting Information Available: Spectroscopic and analytical data for 1, 2, 6-10, 12-17, and A-C as well as selected experimental procedures (15 pages). This material is contained in many libraries on microfiche, immediately follows this article in the microfilm version of the journal, can be ordered from the ACS, and can be downloaded from the Internet; see any current masthead page for ordering information and Internet access instructions.

JA9529495

- (26) Negishi, E.; Valente, L. F.; Kobayashi, M. J. Am. Chem. Soc. 1980,
- 102, 3298.
 (27) The interesting reactivity of the iodide will be described in our full account of this work
- (28) Ikeda, Y.; Ukai, J.; Ikeda, N.; Yamamoto, H. Tetrahedron 1987, 43, 723.
 - (29) Heathcock^{3b} has also employed the Yamamoto protocol.
 - (30) Kuzuhara, H.; Fletcher, H. G., Jr. J. Org. Chem. 1967, 32, 2531.
 (31) Kocovsky, P. Tetrahedron Lett. 1986, 27, 5521.
- (32) We thank Professor Schreiber (Harvard University) for a generous sample of synthetic (-)-discodermolide.