

Synthesis and Absolute Configurations of the Cytotoxic Polyacetylenes Isolated from the Callus of *Panax ginseng*

Yasuo FUJIMOTO^{*,a} Mitsuru SATOH,^b Naoki TAKEUCHI^b and Makoto KIRISAWA^a


Nihon University, College of Pharmacy,^a 7-7 Narashinodai, Funabashi, Chiba 274, Japan and Showa College of Pharmaceutical Sciences,^b Tsurumaki, Setagaya, Tokyo 154, Japan. Received September 4, 1989

Panaxacol (1) and dihydropanaxacol (2), cytotoxic polyacetylenes isolated from the callus of *Panax ginseng*, were synthesized starting from D-(−)-diethyl tartrate. The absolute configuration of 1 was determined to be 9*R*, 10*R* and the absolute configuration at C-3 of 2 was tentatively assigned as 3*S* by the application of the *R*(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl (MTPA) method.

Keywords *Panax ginseng*; panaxacol; callus; dihydropanaxacol; absolute configuration; polyacetylene; chiral synthesis

In the previous papers,^{1,2)} we have reported the isolation and structural elucidation of three new cytotoxic polyacetylenes (**1**, **2** and **3**) and panaxydol (**4**)³⁾ isolated from the callus of *Panax ginseng* C. A. MEYER. Although we have established the relative configuration at C-9 and C-10 in panaxacol to be *threo* from the results of nuclear Overhauser effect (NOE) experiments on the acetonide of **1**, the absolute configurations of **1** and **2** have not been determined yet. Thus, we planned to establish the absolute configurations of these polyacetylenes by synthesizing them, using L-(+)- or D-(−)-diethyl tartrate as a chiral template.

D-(−)-Diethyl tartrate was treated with 2,2-dimethoxypropane and camphorsulfonic acid (CSA) to give an acetonide (**5**, 68.5%, 15% recovery of starting material), which was then converted into a diol (**6**, 77.3%)^{4a,b)} by the action of LiAlH₄. Tosylation of **6** with *p*-toluenesulfonyl chloride (TsCl, 1.0 eq)-pyridine gave a mixture of the monotosylate (**7**, 54.5%) and ditosylate (**8**, 14.8%)^{4a,b)} which was separated by silica gel column chromatography. After protection of the hydroxyl group with tetrahydropyranyl (THP) ether, **7** was treated with dihexyl copper lithium⁵⁾ and then 2*N* HCl-MeOH (1:4) to afford



a triol (**9**, 66.4%). Tosylation of **9** gave a diol-monotosylate (**10**, 43.6%) which was transformed to an acetonide. Coupling reaction of the acetonide with 1,3-heptadiyne-5-ol THP ether in the presence of BuLi-hexamethylphosphoric triamide (HMPA) failed to give the desired compounds. Therefore, the compound (**10**) was converted into an epoxide (**11**, 75.7%) by usual means and **11** was allowed to react with 1,3-heptadiyne-5-ol THP ether to give the coupling products, which, without purification, were deprotected to afford a diastereomeric mixture of 9*R*,10*R*-dihydropanaxacol (**12**, 40%, $[\alpha]_D +12.3^\circ$ ($c=0.79$, MeOH)). Similarly, a diastereomeric mixture of 9*S*,10*S*-dihydropanaxacol ($[\alpha]_D -17.0^\circ$ ($c=0.36$, MeOH)) was synthesized from L-(+)-diethyl tartrate. The diastereomeric mixture of dihydropanaxacol (**12**) prepared from natural panaxacol (**1**) by NaBH₄ reduction exhibited positive optical rotation ($[\alpha]_D +13.5^\circ$ ($c=1.0$, MeOH)). Thus, the absolute configuration of panaxacol can be presumed to be the same as that of D-(+)-diethyl tartrate. In order to confirm this presumption, **12** was converted into panaxacol as follows.

The glycol moiety of **12** was protected as the acetonide and then the hydroxyl group at C-3 was oxidized by Swern's method. Deprotection of the oxidation product, followed by purification using high-performance liquid chromatography (HPLC) gave a solid which was identical including optical rotation with natural panaxacol. Thus, the absolute configurations at C-9 and C-10 in panaxacol were confirmed as 9*R* and 10*R*.

Finally, the absolute configuration at C-3 in dihydropanaxacol was tentatively assigned⁶⁾ by the application of Mosher's method. The acetonide (**13**) was treated with *R*-(+)- α -methoxy- α -(trifluoromethyl)phenylacetyl chloride (MTPA-Cl)⁷⁾ to give a diastereomeric mixture of *R*-(+)-MTPA esters which was separated into *R*-(+)-MTPA ester (A) (retention time: 9.2 min) and (B) (retention time: 10.6 min) by HPLC. A comparison of the proton nuclear magnetic resonance spectrum (¹H-NMR) of ester (A) with that of ester (B) revealed that the methyl (δ 0.93) and the methylene (δ 1.83) signals of the ester (A) appeared in higher

field than those (δ 1.03 for the methyl and δ 1.98 for the methylene) of the ester (B). Therefore, the absolute configurations at C-3 of the ester (A) and the ester (B) were deduced to be *R* and *S*, respectively, based on the general rule of the MTPA method⁸⁾ (Chart 1).

Similarly, natural dihydropanaxacol was also transformed to the *R*-(+)-MTPA ester after protection of the glycol moiety. The ester obtained here was identical with the MTPA ester (B) in terms of the ¹H-NMR and retention time on HPLC. Thus, the absolute configuration of dihydropanaxacol was deduced to be 3*S*, 9*R*, 10*R*. The biological activities of these polyacetylenes and related compounds will be reported elsewhere.

Experimental

Melting points were determined on a Yanagimoto melting point apparatus and are uncorrected. Spectral data were obtained on the following instruments: Infrared spectrum (IR) on a Shimadzu IR-430; ¹H-NMR on a JEOL GX-400, GX-270 or FX-100 instrument in CDCl₃ containing tetramethylsilane as an internal standard; mass spectra (MS) on a Hitachi RMU-6M or a JEOL JMS-D 300.

(2*S*,3*S*)-1,2,3-Trihydroxydecyloxy-4-tosyloxybutan-1-ol (**7**) TsCl (3.0 g) was added to a stirred solution of **6** (3.2 g) in pyridine (30 ml), and the mixture was allowed to stand overnight at room temperature. Brine (50 ml) was then added, and the reaction mixture was extracted with AcOEt. The organic layer was concentrated under reduced pressure to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=2:1) to give **7** (3.4 g, 54.5%) as an oil, the ditosylate (**8**, 14.8%) and **6** (10% recovery). **7**: Oil, EI-MS m/z (relative intensity): 301 ($M-Me$)⁺ (100), 285 ($M-CH_2OH$)⁺ (9); ¹H-NMR δ : 1.33, 1.39 (3H each, s), 2.45 (3H, s), 3.63 (1H, dd, $J=4.2$, 12.0 Hz), 3.80 (1H, dd, $J=3.7$, 12.0 Hz), 3.81 (1H, m), 3.99 (1H, m), 4.04 (1H, m), 4.11 (1H, m), 7.36 (2H, d, $J=8.3$ Hz), 7.80 (2H, d, $J=8.3$ Hz).

(2*R*,3*R*)-1,2,3-Trihydroxydecane (**9**) A mixture of **7** (1.7 g), 3,4-dihydro-2*H*-pyran (DHP) (670 mg) and CSA (50 mg) in CH₂Cl₂ (10 ml) was stirred for 30 min at room temperature. The mixture was diluted with AcOEt (50 ml) and then washed with saturated NaHCO₃. The organic layer was dried over Na₂SO₄ and then concentrated under reduced pressure to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=3:1) to afford the THP ether of **7** [2.0 g, 93.0%]; EI-MS m/z (relative intensity): 400 (M)⁺ (2), 385 ($M-Me$)⁺ (100); ¹H-NMR δ : 1.35 (3H, s), 1.38 (3H, s), 1.40–1.60 (8H, br m, $W_{1/2}=20$ Hz), 2.45 (3H, s), 3.52 (1H, m), 3.57 (1H, m), 4.0–4.2 (4H, m), 4.61 (1H, m), 7.34 (2H, d, $J=8.3$ Hz), 7.81 (2H, d, $J=8.3$ Hz)]. This compound was submitted to




Chart 1

the following reaction. An Et₂O solution (9.0 ml) of hexyl lithium (1.56 mmol/ml) was added dropwise to a stirred suspension of cuprous iodide (1.34 g) in Et₂O (25 ml) at -40°C. After 15 min, a solution of the above THP ether (930 mg) in Et₂O (8 ml) was added at -30°C and then the mixture was further stirred for 1 h at the same temperature. Saturated NH₄Cl (10 ml) was added to the reaction mixture and the whole was extracted with Et₂O. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=7:1), followed by HPLC (Senshu Pack Silica-3301-N, 8 × 300 mm, hexane:AcOEt=20:1) to give a diastereomeric mixture of coupling products (500 mg, 66.4%). The mixture of coupling products (400 mg) was dissolved in 5 ml of MeOH-2N HCl (4:1) under stirring at room temperature. After 1 h, brine (40 ml) was added and the mixture was extracted with AcOEt. The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated *in vacuo* to leave **9** (206 mg, 88.0%) as colorless crystals. **9**: mp 52–54°C, CI-MS *m/z*: 191 (M+1)⁺, EI-MS *m/z* (relative intensity): 159 (M-CH₂OH)⁺ (17), 129 (M-C₂H₅O₂)⁺ (41), 111 (M-C₂H₇O₃)⁺ (100); IR (CHCl₃): 3400, 2930, 2850 cm⁻¹; ¹H-NMR δ: 0.95 (3H, t, *J*=7.1 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.50 (2H, br m), 3.62 (2H, m), 3.75 (2H, m); ¹³C-NMR δ: 14.1, 22.7, 25.8, 29.4, 29.8, 31.9, 33.6, 64.5, 72.3, 74.4.

(2R,3R)-1-Tosyloxy-2,3-dihydroxydecane (10) A mixture of **9** (190 mg) and TsCl (185 mg) in pyridine (2 ml) was allowed to stand overnight at room temperature. The mixture was diluted with AcOEt (50 ml) and then washed with brine, dried (Na₂SO₄), and evaporated *in vacuo*. The residue was chromatographed on a silica gel column (hexane:AcOEt=3:1) to give **10** as a colorless solid (150 mg, 43.6%), mp 63°C, CI-MS *m/z*: 345 (M+1)⁺; EI-MS *m/z* (relative intensity): 301 (M-C₃H₇)⁺ (3), 215 (M-C₈H₁₇O)⁺ (2), 173 (M-OTs)⁺ (100); IR (CHCl₃): 3550, 2930, 2850, 1730, 1600, 1460 cm⁻¹; ¹H-NMR δ: 0.88 (3H, t, *J*=7.1 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.42 (2H, m), 2.46 (3H, s), 3.72 (2H, br m), 4.05 (1H, dd, *J*=5.6, 12.8 Hz), 4.12 (1H, dd, *J*=5.4, 12.8 Hz), 7.35 (2H, d, *J*=8.3 Hz), 7.80 (2H, d, *J*=8.3 Hz); ¹³C-NMR δ: 14.2, 21.8, 22.8, 25.7, 29.4, 29.6, 31.9, 33.5, 70.9, 71.6, 78.5, 128.0 (two carbons), 130.0 (two carbons), 132.6, 145.2.

(2R,3R)-1,2-Epoxydecan-3-ol THP Ether (11) A mixture of **10** (114 mg) and K₂CO₃ (457 mg) in MeOH (3.5 ml) was stirred for 1 h at room temperature. The mixture was diluted with AcOEt (40 ml), washed with brine, dried (Na₂SO₄), and evaporated *in vacuo* to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=2:1) to give an epoxide (55 mg, 75.7%). ¹H-NMR δ: 0.88 (3H, t, *J*=7.3 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.60 (2H, m), 1.78 (1H, d, *J*=5.9 Hz, OH), 2.72 (1H, dd, *J*=2.7, 4.9 Hz), 2.83 (1H, dd, *J*=3.9, 4.9 Hz), 2.98 (1H, m), 3.44 (1H, m). A mixture of the above epoxide (50 mg), DHP (40 μl) and CSA (trace amount) in CH₂Cl₂ (1.5 ml) was stirred for 30 min at room temperature. The mixture was diluted with AcOEt (20 ml) and washed successively with saturated NaHCO₃ and brine, dried over Na₂SO₄, and evaporated *in vacuo* to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=3:1) to afford **11** (67 mg, 90.0%). **11**: A mixture of diastereomers, two spots (*R*_f 0.40 and 0.47) on a silica gel thin-layer chromatography (TLC) plate (hexane:AcOEt=5:1), MS *m/z* (relative intensity): 256 (M)⁺ (5), 213 (M-C₂H₅O)⁺ (27), 157 (M-C₇H₁₅)⁺ (56), 155 (M-OTHP)⁺ (100), ¹H-NMR δ: 0.88 (t, *J*=7.1 Hz), 1.20–1.40 (br m, *W*_{1/2}=25 Hz), 1.58 (m), 1.76 (m), 1.84 (m), 2.49 (dd, *J*=2.7, 4.9 Hz), 2.67 (dd, *J*=2.7, 4.9 Hz), 2.76 (t, *J*=4.9 Hz), 2.80 (t, *J*=4.9 Hz), 2.95 (m), 3.09 (m), 3.38 (m), 3.50 (m), 3.88 (m), 4.00 (m), 4.70 (t, *J*=4.4 Hz), 4.98 (t, *J*=4.4 Hz).

1,3-Heptadiyn-5-ol THP Ether BuLi in hexane [10 ml (1.5 mmol/ml)] was added dropwise to a stirred solution of diacetylene⁹⁾ in tetrahydrofuran (THF) [15 ml (1 mmol/ml)] at -50°C and then propionaldehyde (840 mg) was added at the same temperature. After 30 min, saturated NH₄Cl (50 ml) was added to the reaction mixture and then the mixture was extracted with AcOEt. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=5:1) to give 1,3-heptadiyn-5-ol [¹H-NMR δ: 1.03 (3H, t, *J*=7.1 Hz), 1.76 (2H, m), 1.95 (1H, br s, OH), 2.20 (1H, d, *J*=1.0 Hz), 4.38 (1H, m); ¹³C-NMR δ: 9.2, 30.4, 63.6, 67.3, 68.2, 68.9, 77.1]. A mixture of 1,3-heptadiyn-5-ol, DHP (1 ml) and CSA (20 mg) in CH₂Cl₂ (10 ml) was stirred for 1 h at room temperature. The mixture was diluted with AcOEt (20 ml) and then washed successively with saturated NaHCO₃ and brine. The organic layer was dried over Na₂SO₄, and evaporated *in vacuo*. The residual material was chromatographed on a silica gel column (hexane:AcOEt=7:1) to afford 1,3-heptadiyn-5-ol THP ether (1.4 g, 50% from diacetylene). [A mixture

of diastereomers; two spots on a silica gel TLC plate (*R*_f 0.60 and 0.68, hexane:AcOEt=5:1), CI-MS *m/z*: 193 (M+1)⁺, ¹H-NMR δ: 1.00 (t, *J*=7.3 Hz), 1.04 (t, *J*=7.3 Hz), 1.56 (m), 1.78 (m), 2.14 (d, *J*=0.9 Hz), 2.15 (d, *J*=0.9 Hz), 3.54 (m), 3.57 (m), 3.80 (ddd, *J*=3.1, 8.6, 11.6 Hz), 4.00 (ddd, *J*=3.1, 9.8, 12.8 Hz), 4.25 (t, *J*=6.4 Hz), 4.41 (t, *J*=6.4 Hz), 4.74 (t, *J*=3.4 Hz), 4.93 (t, *J*=4.0 Hz)].

(9R,10R)-Dihydropanaxacol (12, a Diastereomeric Mixture at C-3) BuLi (0.3 mmol) in hexane (230 μl) and the epoxide (**11**, 25 mg) in THF (100 μl) were successively added dropwise to a stirred solution of 1,3-heptadiyn-5-ol THP ether (77 mg, 0.4 mmol) and HMPA (50 μl) in THF (130 μl) at -30°C. After 2 h, the reaction was quenched with saturated NH₄Cl (10 ml), and then the mixture was extracted with AcOEt. The extracts were washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to leave an oil, which was chromatographed on a silica gel column (hexane:AcOEt=5:1) to give a diastereomeric mixture of the coupling products (two spots on a silica gel TLC plate, hexane:AcOEt=5:1). A mixture of the coupling products (10 mg) and CSA (trace amount) in MeOH (500 μl) was stirred at room temperature. After 1 h, the mixture was diluted with AcOEt (10 ml), then washed with brine, dried over Na₂SO₄, and evaporated *in vacuo*. The residue was purified by HPLC (Senshu Pack Silica-3251-N, 8 × 250 mm, hexane:AcOEt=2:1) to give **12** (4 mg, 64.5%). [*α*]_D +12.3° (*c*=0.79, MeOH), CI-MS *m/z*: 281 (M+1)⁺, IR (CHCl₃): 3500–3300, 2930 (s), 2850 (m), 2260 (w) cm⁻¹, ¹H-NMR δ: 0.89 (3H, t, *J*=7.0 Hz), 1.02 (3H, t, *J*=7.3 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.50 (2H, m), 1.74 (2H, m), 1.88 (1H, d, *J*=5.8 Hz, OH), 2.01 (1H, d, *J*=5.5 Hz, OH), 2.37 (1H, d, *J*=5.8 Hz, OH), 2.56 (1H, dd, *J*=6.4, 16.5 Hz), 2.59 (1H, dd, *J*=6.4, 16.5 Hz), 3.59 (1H, m), 3.64 (1H, m), 4.36 (1H, q, *J*=6.1 Hz); ¹³C-NMR δ: 9.36 (q), 14.09 (q), 22.66 (t), 24.91 (t), 25.62 (t), 29.24 (t), 29.59 (t), 30.74 (t), 31.84 (t), 33.59 (t), 64.04 (d), 66.67 (s), 69.61 (s), 72.24 (d), 73.17 (d), 77.31 (s), 77.51 (s).

(9R,10R)-Dihydropanaxacol Acetonide (13) A mixture of **12** (72 mg), CSA (10 mg) and 2,2-dimethoxypropane (3 ml) was stirred for 30 min at room temperature. The mixture was diluted with AcOEt (30 ml) and washed with saturated NaHCO₃. The organic layer was washed with brine, dried (Na₂SO₄), and concentrated *in vacuo* to leave an oily material, which was chromatographed on a silica gel column (hexane:AcOEt=5:1) to give **13** (80 mg, 97.3%). EI-MS *m/z* (relative intensity): 305 (M-CH₃)⁺ (17.7), 245 (M-C₃H₇O₂)⁺ (12.2), 199 (M-C₈H₉O)⁺ (100); ¹H-NMR δ: 0.89 (3H, t, *J*=6.8 Hz), 1.01 (3H, t, *J*=7.33 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.40 (6H, s), 1.60 (2H, m), 1.74 (2H, m), 1.85 (1H, br s, OH), 2.58 (1H, dd, *J*=5.4, 17.6 Hz), 2.63 (1H, dd, *J*=5.4, 17.6 Hz), 3.74 (1H, m), 3.79 (1H, m), 4.36 (1H, t, *J*=6.6 Hz); ¹³C-NMR: 9.32 (q), 14.09 (q), 22.66 (t), 23.55 (t), 25.97 (t), 26.98 (q), 27.37 (q), 29.16 (t), 29.66 (t), 30.75 (t), 31.82 (t), 32.90 (t), 64.06 (d), 66.66 (s), 69.65 (s), 76.58 (s), 77.18 (s), 78.21 (d), 80.56 (d), 108.76 (s).

Panaxacol Dimethylsulfoxide (20 μl) and **13** (20 mg) in CH₂Cl₂ (100 μl) were successively added dropwise to a stirred solution of oxalyl chloride (12 μl) in CH₂Cl₂ (250 μl) at -50°C. After 1 h, triethylamine (170 μl) was added to the mixture and then stirring was continued for 10 min. The reaction mixture was diluted with AcOEt (40 ml), then washed with brine, dried over Na₂SO₄, and evaporated *in vacuo*. The residual oily material was dissolved in MeOH (2 ml) and then CSA (trace amount) was added at room temperature under stirring. After 2 h, the reaction mixture was diluted with AcOEt (30 ml), washed with brine, dried over Na₂SO₄, and evaporated *in vacuo*. The residue was purified by HPLC (Senshu Pack Silica-3251-N, 8 × 250 mm, hexane:AcOEt=2:1) to give panaxacol (retention time: 13.2 min) (10 mg, 60.0%). [*α*]_D +18.3° (*c*=1.0, MeOH); CI-MS *m/z*: 279 (M+1)⁺; IR (CHCl₃): 3560 (v), 3400 (s), 2930 (s), 2850 (m), 2240 (s), 2150 (w), 1665 (s) cm⁻¹; ¹H-NMR δ: 0.89 (3H, t, *J*=7.1 Hz), 1.15 (3H, t, *J*=7.3 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.51 (2H, m), 1.91 (1H, d, *J*=5.6 Hz, OH), 2.36 (1H, d, *J*=5.9 Hz, OH), 2.60 (2H, q, *J*=7.3 Hz), 2.64 (1H, dd, *J*=6.3, 17.6 Hz), 2.68 (1H, dd, *J*=5.9, 17.6 Hz), 3.59 (1H, m), 3.70 (1H, m).

R-(+)-MTPA Ester (A) and (B) A mixture of **13** (8 mg) and five drops (large excess) of *R*-MTPA-Cl in pyridine (500 μl) was allowed to stand overnight at room temperature. The mixture was diluted with AcOEt (30 ml), washed with brine, dried over Na₂SO₄, and evaporated *in vacuo*. The residue was separated by HPLC (Senshu Pack Silica-3251-N, 8 × 250 mm, hexane:AcOEt=20:1) to give the *R*-(+)-MTPA ester (A) (retention time: 9.2 min) and *R*-(+)-MTPA ester (B) (retention time: 10.6 min).

R-(+)-MTPA Ester (A): ¹H-NMR δ: 0.88 (3H, t, *J*=7.1 Hz), 0.93 (3H, t, *J*=7.3 Hz), 1.20–1.40 (10H, br m, *W*_{1/2}=25 Hz), 1.40 and 1.41 (3H each, s), 1.58 (2H, m), 1.83 (2H, m), 2.58 (1H, dd, *J*=5.1, 17.8 Hz), 2.64 (1H, dd, *J*=5.1, 17.8 Hz), 3.59 (3H, br s), 3.73 (1H, m), 3.84 (1H, m), 5.54

(1H, t, $J=6.6$ Hz), 7.42 (3H, m), 7.53 (2H, m).

R-(+)-MTPA Ester (B): $^1\text{H-NMR}$ δ : 0.88 (3H, t, $J=7.1$ Hz), 1.03 (3H, t, $J=7.3$ Hz), 1.20—1.40 (10H, br m, $W_{1/2}=25$ Hz), 1.40 and 1.41 (3H each, s), 1.58 (1H, m), 1.89 (2H, m), 2.58 (1H, dd, $J=5.1, 18.3$ Hz), 2.64 (1H, dd, $J=5.1, 18.3$ Hz), 3.55 (3H, br s), 3.73 (1H, m), 3.80 (1H, m), 5.51 (1H, t, $J=6.6$ Hz), 7.43 (3H, m), 7.52 (2H, m).

References and Notes

- 1) Y. Fujimoto and M. Satoh, *Phytochemistry*, **26**, 2850 (1987).
- 2) Y. Fujimoto and M. Satoh, *Chem. Pharm. Bull.*, **36**, 4206 (1988).
- 3) J. Poplawski, J. T. Wrobel and T. Glinka, *Phytochemistry*, **19**, 1539 (1980).
- 4) a) M. Carmack and C. J. Kelley, *J. Org. Chem.*, **33**, 2171 (1968); b) K. Mori and S. Tamada, *Tetrahedron*, **35**, 1279 (1979).
- 5) Dihexyl copper lithium was prepared by the reaction of cuprous iodide and hexyl lithium: e.g., C. R. Johnson and G. A. Dutra, *J. Am. Chem. Soc.*, **95**, 7777 (1973).
- 6) Final confirmation of the absolute configuration at C-3 by synthetic correlation using chiral starting material is in progress and the result will be published elsewhere.
- 7) a) J. A. Dale and H. S. Mosher, *J. Am. Chem. Soc.*, **95**, 512 (1973); b) G. R. Sullivan, J. A. Dale and H. S. Mosher, *J. Org. Chem.*, **38**, 2143 (1973).
- 8) S. Yamaguchi, "Asymmetric Synthesis," Vol. 1, ed. by J. D. Morrison, Academic Press, New York, 1983, p. 125 and references cited therein.
- 9) L. Brandsma and H. D. Verkrujsse, "Studies in Organic Chemistry, Vol. 8, Synthesis of Acetylenes, Allenes and Cumulenes," Elsevier Scientific Publishing Co., Amsterdam, 1981, p. 146.