

Iron-catalyzed Pinacol Coupling of Aryl Ketones with a Phenyltitanium Reagent: A New Type of Catalytic Reaction

Tamio Hayashi* and Keigo Sasaki

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502

(Received February 28, 2011; CL-110165; E-mail: thayashi@kuchem.kyoto-u.ac.jp)

A reaction of aryl ketones with phenyltitanium triisopropoxide ($[\text{PhTi}(\text{Oi-Pr})_3]$) in the presence of $[\text{Fe}(\text{acac})_3]$ as a catalyst (1 mol %) gave the corresponding pinacols in high yields. The catalytic cycle of this process involves an iron-catalyzed disproportionation of $[\text{PhTi}(\text{Oi-Pr})_3]$ into biphenyl and a low-valent titanium species.

Since iron is one of the most abundant metals on earth and consequently is one of the most inexpensive and environmentally friendly, there have recently been extensive studies on the use of iron salts and complexes as catalysts for organic transformations.¹ During the course of our studies on iron-catalyzed carbon–carbon bond forming reactions,² we found that a new type of iron-catalyzed transformation takes place for the reaction of aryl ketones with phenyltitanium triisopropoxide^{3–5} ($[\text{PhTi}(\text{Oi-Pr})_3]$), which gives high yields of the pinacol coupling products.

Table 1 demonstrates the unique catalysis by iron salt to promote pinacol coupling. Thus, to a solution of $[\text{Fe}(\text{acac})_3]$ (0.01 mmol, 1 mol %) in THF (2.0 mL) were added a solution of $[\text{PhTi}(\text{Oi-Pr})_3]$, which was generated in a separate flask from $[\text{CITi}(\text{Oi-Pr})_3]$ (1.5 mmol) and PhLi in cyclohexane/ Et_2O (1.6 mmol), and acetophenone (**1a**, 1.0 mmol) successively, and the mixture was stirred at 20 °C for 3 h. Acidic hydrolysis with 1 M HCl followed by silica gel chromatography gave 0.40 mmol (80% yield) of pinacol coupling product, 2,3-diphenylbutane-2,3-diol (**2a**, *dl/meso* = 8/2), together with 0.485 mmol (97% based on **1a**) of biphenyl (**3**) and 0.19 mmol (19%) of 1,1-diphenylethanol (**4a**) (Entry 1). In the absence of $[\text{Fe}(\text{acac})_3]$, the pinacol **2a** or biphenyl (**3**) was not formed under otherwise the same conditions. Instead, the reaction gave 64% yield of alcohol **4a** which should result from nucleophilic attack of the phenyltitanium reagent on ketone **1a** (Entry 2). Thus, the present pinacol coupling is clearly demonstrated to be catalyzed by the small amount of $[\text{Fe}(\text{acac})_3]$. The yield of pinacol **2a** was slightly enhanced by increasing the amount of $[\text{Fe}(\text{acac})_3]$ catalyst to 5 mol % (Entry 3). The use of isolated $[\text{PhTi}(\text{Oi-Pr})_3]$ in place of the in situ generated phenyltitanium also gave pinacol **2a** albeit in a little lower yield (Entry 4). The pinacol coupling was also observed with FeCl_3 as a catalyst (Entry 5). Of other metal complexes examined, $[\text{Co}(\text{acac})_3]$ was catalytically as active as $[\text{Fe}(\text{acac})_3]$ to give a high yield of **2a** (Entry 6). $[\text{Ni}(\text{acac})_2]$ also catalyzed the present pinacol coupling although its activity is lower than that of $[\text{Fe}(\text{acac})_3]$ or $[\text{Co}(\text{acac})_3]$ (Entry 7). Ruthenium, rhodium, or palladium acac complex did not catalyze the pinacol coupling (Entries 8, 9, and 10).

It has been well-documented that trivalent titanium species, generated by reduction of tetravalent titanium complexes with magnesium or zinc metal, function as one-electron reducing reagents of carbonyl compounds to lead to pinacol coupling.^{6,7}

Table 1. Pinacol coupling of acetophenone (**1a**) with $[\text{PhTi}(\text{Oi-Pr})_3]$ in the presence of transition-metal catalysts^a

Entry	Catalyst	Yield/% ^b 2a ^c	Yield/% ^b 4a	Recovered/% ^b 1a
1 ^d	$[\text{Fe}(\text{acac})_3]$	81 (80)	19	<1
2 ^d	none	0	64	36
3 ^e	$[\text{Fe}(\text{acac})_3]$	83 (80)	17	<1
4 ^f	$[\text{Fe}(\text{acac})_3]$	63	32	<1
5	FeCl_3	75 (73)	25	<1
6	$[\text{Co}(\text{acac})_3]$	83	17	<1
7	$[\text{Ni}(\text{acac})_2]$	36	38	3
8	$[\text{Ru}(\text{acac})_3]$	4	53	43
9	$[\text{Rh}(\text{acac})_3]$	4	51	40
10	$[\text{Pd}(\text{acac})_2]$	3	50	43

^aThe reaction was carried out with **1a** (1.0 mmol), $[\text{PhTi}(\text{Oi-Pr})_3]$ (1.5 mmol, generated from $[\text{CITi}(\text{Oi-Pr})_3]$ and PhLi), catalyst (0.01 mmol, 1 mol %) in THF at 20 °C for 3 h. ^bDetermined by ¹H NMR. The numbers in parentheses are isolated yield. ^cThe ratio of *dl/meso* **2a** is 8/2 for all the Entries 1 and 3–10. ^dThe yields of biphenyl (**3**) are 97% and <2% for Entries 1 and 2, respectively. ^eThe reaction with 5 mol % of $[\text{Fe}(\text{acac})_3]$. ^fThe reaction with isolated $[\text{PhTi}(\text{Oi-Pr})_3]$.

Some organometallic reagents such as Grignard reagents have also been used to reduce titanium(IV) complexes for the pinacol coupling. For example, “[$\text{Ti}(\text{Oi-Pr})_3$]” generated by treatment of $[\text{Ti}(\text{Oi-Pr})_4]$ with EtMgBr is reported to be an effective reagent for the coupling of aromatic aldehydes and ketones.⁸ It is interesting that $[\text{PhTi}(\text{Oi-Pr})_3]$, which is a stable titanium(IV) complex, promotes the pinacol coupling in the presence of an iron catalyst in our system. The formation of biphenyl (**3**) observed in the iron-catalyzed pinacol coupling suggests that $[\text{PhTi}(\text{Oi-Pr})_3]$ undergoes disproportionation in the presence of an iron catalyst to give biphenyl (**3**) and a low-valent titanium species, most likely “[$\text{Ti}(\text{Oi-Pr})_3$]”, which will bring about the pinacol coupling of ketones (Scheme 1).

The reaction pathway proposed in Scheme 1 is supported in part by control experiments shown in Scheme 2. Thus, to a solution of $[\text{PhTi}(\text{Oi-Pr})_3]$ (1.0 mmol) in THF, $[\text{Fe}(\text{acac})_3]$ (0.10 mmol) was added at 20 °C, and the mixture was stirred at the same temperature for 10 min. Acidic hydrolysis of the reaction mixture gave 0.29 mmol (58% yield) of biphenyl (**3**),

References and Notes

- For reviews, see: a) *Iron Catalysis in Organic Chemistry: Reactions and Applications*, ed. by B. Plietker, Wiley-VCH, Weinheim, **2008**. b) C. Bolm, J. Legros, J. L. Paih, L. Zani, *Chem. Rev.* **2004**, *104*, 6217. c) M. Nakamura, S. Ito, in *Modern Arylation Methods*, ed. by L. Ackermann, Wiley-VCH, Weinheim, **2009**, p. 155. d) H. Shinokubo, K. Oshima, *Eur. J. Org. Chem.* **2004**, 2081. e) A. Fürstner, R. Martin, *Chem. Lett.* **2005**, *34*, 624. f) A. Correa, O. G. Mancheño, C. Bolm, *Chem. Soc. Rev.* **2008**, *37*, 1108. g) B. D. Sherry, A. Fürstner, *Acc. Chem. Res.* **2008**, *41*, 1500. h) W. M. Czaplik, M. Mayer, J. Cvengroš, A. J. von Wangelin, *ChemSusChem* **2009**, *2*, 396. For one of the recent topics on the iron-catalyzed coupling reactions, see: i) T. Hatakeyama, T. Hashimoto, Y. Kondo, Y. Fujiwara, H. Seike, H. Takaya, Y. Tamada, T. Ono, M. Nakamura, *J. Am. Chem. Soc.* **2010**, *132*, 10674.
- a) T. Nagano, T. Hayashi, *Org. Lett.* **2004**, *6*, 1297. b) T. Nagano, T. Hayashi, *Org. Lett.* **2005**, *7*, 491. c) E. Shirakawa, T. Yamagami, T. Kimura, S. Yamaguchi, T. Hayashi, *J. Am. Chem. Soc.* **2005**, *127*, 17164. d) T. Yamagami, R. Shintani, E. Shirakawa, T. Hayashi, *Org. Lett.* **2007**, *9*, 1045. e) E. Shirakawa, D. Ikeda, T. Ozawa, S. Watanabe, T. Hayashi, *Chem. Commun.* **2009**, 1885. f) E. Shirakawa, D. Ikeda, S. Yamaguchi, T. Hayashi, *Chem. Commun.* **2008**, 1214.
- B. Weidmann, L. Widler, A. G. Olivero, C. D. Maycock, D. Seebach, *Helv. Chim. Acta* **1981**, *64*, 357.
- For a review on organotitanium reagents, see: F. Sato, H. Urabe, S. Okamoto, *Chem. Rev.* **2000**, *100*, 2835.
- For examples of the use of [PhTi(Oi-Pr)₃] in transition-metal-catalyzed reactions, see: a) T. Hayashi, N. Tokunaga, K. Yoshida, J. W. Han, *J. Am. Chem. Soc.* **2002**, *124*, 12102. b) J. W. Han, N. Tokunaga, T. Hayashi, *Synlett* **2002**, 871. c) H. W. Lee, F. L. Lam, C. M. So, C. P. Lau, A. S. C. Chan, F. Y. Kwong, *Angew. Chem., Int. Ed.* **2009**, *48*, 7436.
- For recent pertinent reviews on pinacol coupling, see: a) A. Chatterjee, N. N. Joshi, *Tetrahedron* **2006**, *62*, 12137. b) A. Gansäuer, H. Bluhm, *Chem. Rev.* **2000**, *100*, 2771.
- For early examples of titanium-mediated pinacol coupling, see: a) S. Tyrlik, I. Wolochowicz, *Bull. Soc. Chim. Fr.* **1973**, 2147. b) T. Mukaiyama, T. Sato, J. Hanna, *Chem. Lett.* **1973**, 1041. c) J. E. McMurry, M. P. Fleming, *J. Am. Chem. Soc.* **1974**, *96*, 4708. d) E. J. Corey, R. L. Danheiser, S. Chandrasekaran, *J. Org. Chem.* **1976**, *41*, 260. e) Y. Handa, J. Inanaga, *Tetrahedron Lett.* **1987**, *28*, 5717. For recent progress, see: f) M. Paradas, A. G. Campaña, R. E. Estévez, L. Á. de Cienfuegos, T. Jiménez, R. Robles, J. M. Cuerva, J. E. Oltra, *J. Org. Chem.* **2009**, *74*, 3616. g) M. C. Barden, J. Schwartz, *J. Am. Chem. Soc.* **1996**, *118*, 5484. h) T. Tsuritani, S. Ito, H. Shinokubo, K. Oshima, *J. Org. Chem.* **2000**, *65*, 5066. i) T. Li, W. Cui, J. Liu, J. Zhao, Z. Wang, *Chem. Commun.* **2000**, 139. j) T. Mukaiyama, N. Yoshimura, K. Igarashi, *Chem. Lett.* **2000**, 838. k) A. Bensari, J.-L. Renaud, O. Riant, *Org. Lett.* **2001**, *3*, 3863. l) A. Gansäuer, *Synlett* **1997**, 363. m) T. A. Lipski, M. A. Hilfiker, S. G. Nelson, *J. Org. Chem.* **1997**, *62*, 4566. n) A. Chatterjee, T. H. Bennur, N. N. Joshi, *J. Org. Chem.* **2003**, *68*, 5668. o) Y.-G. Li, Q.-S. Tian, J. Zhao, Y. Feng, M.-J. Li, T.-P. You, *Tetrahedron: Asymmetry* **2004**, *15*, 1707.
- E. A. Matiushenkov, N. A. Sokolov, O. G. Kulinkovich, *Synlett* **2004**, 77.
- The *dl*-selectivity of the titanium-mediated pinacol coupling of aryl alkyl ketones has been reported in refs. 7f and 8.
- CCDC-805917 contains the supplementary crystallographic data for *dl*-**21**. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre via http://www.ccdc.cam.ac.uk/data_request/cif. See also Supporting Information.
- Supporting Information is available electronically on the CSJ-Journal Web site, <http://www.csj.jp/journals/chem-lett/index.html>.