

powder and sodium iodide.¹ A compound was secured having a melting point of 255–258°. It was contaminated with amyl alcohol, which was removed in part by application of heat. The yield was small and complete purification was not possible. Analysis for sulfur gave 8.67%, while theory calls for 12.35%. The compound had a bitter taste. Repetitions of this experiment did not again yield this compound, and lack of starting materials forced the abandonment of the project. In view of the known instances in which seven- and eight-membered ring compounds have been made from diphenic acid, the failure of this synthesis is without explanation, and seems worthy of report.

CONTRIBUTION FROM THE
CHEMICAL LABORATORY OF
THE UNIVERSITY OF ARKANSAS
FAYETTEVILLE, ARKANSAS

E. WERTHEIM

RECEIVED DECEMBER 2, 1930
PUBLISHED MARCH 6, 1931

The *p*-Bromoanilides of Isobutyric and Isovaleric Acids.—For purposes of comparison in identification work in this Laboratory it was desired to have samples of the *p*-bromoanilides of a number of the paraffin acids. The most comprehensive list of such *p*-bromoanilides which appears in the literature is that reported by Robertson [Robertson, *J. Chem. Soc.*, 115, 1210 (1919)]. This list, however, does not contain the derivatives of isobutyric or isovaleric acids.

The procedure used for the preparation of the *p*-bromoanilides of the acids from acetic to stearic is a modification of Robertson's procedure and is as follows: One gram (1 mol) of the acid or its sodium salt was placed in a small flask or test-tube and treated with 1 mol of thionyl chloride and heated under a reflux condenser for thirty minutes. The reaction mixture was then cooled and treated with 2 mols of *p*-bromoaniline in 30 cc. of benzene. After warming on a steam-bath for a few minutes the benzene solution was cooled, transferred to a separatory funnel, washed twice with water, twice with 5% hydrochloric acid, once with 5% sodium hydroxide and finally with water. The benzene solution was then evaporated to dryness and the remaining bromoanilide recrystallized from ethyl alcohol. The melting points and analyses of these derivatives of isobutyric and isovaleric acids are

| | M. p., °C. | Bromine, % | |
|--------------------------------|------------|------------|----------------|
| | | Calcd. | Found (Carius) |
| <i>p</i> -Bromoisobutyranilide | 150–151 | 33.03 | 32.43 |
| <i>p</i> -Bromoisovaleranilide | 128–129 | 31.10 | 31.12 |

The *p*-bromoisobutyranilide has been reported by Norton [*Am. Chem. J.*, 7, 116 (1885)] as melting at 128°. Judging from the low melting point

¹ Finkelstein, *Ber.*, 43, 1530 (1910).

and satisfactory analysis reported for this *p*-bromoanilide it is quite probable that it was contaminated with some of the ortho isomer, as it was prepared by the bromination of isobutyranilide. *p*-Bromoisovaleranilide does not appear to have been described previously in the literature.

CONTRIBUTION FROM THE
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF WISCONSIN
MADISON, WISCONSIN

MARGUERITE KUEHN
S. M. McELVAIN

RECEIVED JANUARY 26, 1931
PUBLISHED MARCH 6, 1931

COMMUNICATIONS TO THE EDITOR

THE α,β,γ -TRIMETHYLGLUTARIC ACIDS

Sir:

In a recent paper Michael and Ross¹ reported the failure to obtain α,β,γ -trimethyl- α -carboxyglutaric acid in a crystalline form. I had obtained this acid² and reported it as melting at 144–145°. Michael and Ross note that 145° is also the melting point of α,β -dimethyl- γ -carboxyglutaric acid, which suggests that the two are identical. The fact is that I obtained α,β,γ -trimethylglutaric acid melting at 134° from the decomposition of my acid, whereas α,β -dimethyl- γ -carboxyglutaric acid upon decomposition gives α,β -dimethylglutaric acid melting at 87°.³

Michael and Ross obtained α,β,γ -trimethyl- α -carboxyglutaric acid as a sirup which on distillation gave an acid melting at 115–125°. In my work on these compounds I found that it was necessary to obtain the tricarboxylic acid in a pure state before decomposing it in order to effect the purification of the α,β,γ -trimethylglutaric acid. Even so it required ten recrystallizations to reach the melting point of 134°. The compound which Michael and Ross describe as *cis*- α,β,γ -trimethylglutaric acid, which according to their statement "softened at 115° and melted at 125°," was probably an impure acid identical with mine, which melted at 134°. The liquid which they call *trans*- α,β,γ -trimethylglutaric acid was, apparently, the same as the liquid remaining in the mother liquors from the crystallization of the acid melting at 134°. Neither their work nor mine is sufficiently complete to characterize this liquid as a definite chemical individual. Inasmuch as modern theory predicts the occurrence of *three* stereoisomeric forms (one racemic and two meso acids) it would, in my opinion, be unfortunate if it should be so described in handbooks.

Michael and Ross attempted to separate their acids into a "*cis*" and a "*trans*" form by converting the "*cis*" form into the anhydride and the imide. Granting free rotation about a single bond *all three* of the possible

¹ Michael and Ross, *THIS JOURNAL*, **52**, 4607 (1930).

² Ray, *ibid.*, **50**, 558 (1928).

³ Thorpe, *J. Chem. Soc.*, **83**, 358 (1903).