Carbon-rich organic molecules have been attracting much

interest regarding their novel structures and potential ap-

plications.¹ Dehydrobenzoannulenes (DBAs), which were

investigated extensively during the late 1950s through the

mid-1970s with regard to their aromaticity/antiaromaticity,²

constitute one of the central classes of carbon-rich mol-

ecules.³ Thus, multiple fused DBAs have been synthesized

as scaffolds of two-dimensional carbon networks such as

(1) For reviews, see: (a) Diederich, F.; Rubin, Y. Angew. Chem., Int.

Ed. Engl. 1992, 31, 1101. (b) Carbon Rich Compounds I; de Meijere, A.,

Ed.; Springer: Berlin, 1998. (c) Carbon Rich Compounds II; de Meijere,

A., Ed.; Springer: Berlin, 1999. (d) Bunz, U. H. F.; Rubin, Y.; Tobe, Y.

Synthesis and Properties of **Trefoil-Shaped** Tris(hexadehydrotribenzo[12]annulene) and Tris(tetradehydrotribenzo[12]annulene)

Takashi Yoshimura.[†] Akiko Inaba.[†] Motohiro Sonoda.[†] Kazukuni Tahara.[†] Yoshito Tobe,*,† and Richard Vaughan Williams[‡]

Division of Frontier Materials Science, Graduate School of Engineering Science, Osaka University, and CREST, Japan Science and Technology Agency (JST), Toyonaka, Osaka 560-8531, Japan, and Department of Chemistry, University of Idaho, P.O. Box 442343, Moscow, Idaho 83844-2343

tobe@chem.es.osaka-u.ac.jp

Received March 31, 2006

[†] Osaka University.

[‡] University of Idaho.

Chem. Soc. Rev. 1999, 28, 107.

ABSTRACT



⁽²⁾ For selected reviews on annulenes, see: (a) Nakagawa, M. In The Chemistry of the Carbon-Carbon Triple Bond; Patai, S., Ed.; John Wiley & Sons: Bristol, 1978; p 635. (b) Balaban, A. T.; Banciu, M.; Ciorba, V. Annulenes, Benzo-, Hetero-, Homo- Derivatives, and their Valence Isomers; CRC Press: Boca Raton, 1987; Vol. 1, p 67. (c) Huang, N. Z.; Sondheimer, F. Acc. Chem. Res. 1982, 15, 96.

Trefoil-shaped tris(hexadehydrotribenzo[12]annulene) possessing a substructure of the ultimate two-dimensional C(sp)–C(sp²) network, graphyne, and the related tris(tetradehydrotribenzo[12]annulene) were synthesized, and their ground- and excited-state properties were investigated.

ORGANIC LETTERS

⁽³⁾ For reviews on DBAs, see: (a) Wong, H. N. C. In Advances in Theoretically Interesting Molecules; Thummel, R. P., Ed.; JAI Press: Greenwich, 1995; p 109. (b) Youngs, W. J.; Tessier, C. A.; Bradshaw, J. D. Chem. Rev. 1999, 99, 3153. (c) Marsden, J. A.; Palmer, G. J.; Haley, M. M. Eur. J. Org. Chem. 2003, 2355. (d) Tobe, Y.; Sonoda, M. In Modern Cyclophane Chemistry; Gleiter, R., Hopf, H., Eds.; Wiley-VCH: Weinheim, 2004; p 1. (e) Jones, C. S.; O'Connor, M. J.; Haley, M. M. In Acetylene Chemistry: Chemistry, Biology and Material Science; Diederich, F., Stang, P. J., Tykwinski, R. R., Eds.; Wiley-VCH: Weinheim, 2005; p 303.

*t*Bu, *t*Βι ťBu tΒú `tBu ťΒu tΒu

properties. A number of such molecules consisting of aromatic [18]DBA⁶ or [14]DBA⁷ subunits have been reported. On the other hand, there are few examples of such systems constructed from hexadehydrotribenzo[12]annulene, [12]DBA (1a).⁸ This is presumably because of the difficulty associated with the construction of the [12]DBA framework. The weakly antiaromatic [12]DBA exhibits interesting properties such as complexation with transition metals9 and forbidden electronic transitions between S₀ and S₁ states.¹⁰ Only diamond-shaped¹¹ and bow-tie-shaped^{11a,12} bis[12]DBA derivatives have been prepared. We therefore undertook the synthesis of the trefoil-shaped tris[12]DBA 2b to investigate the effect of the mode of fusion of the [12]DBA rings on the ground- and excited-state properties of multiply fused [12]DBAs and to establish a synthetic route to the "full wheel-shaped" hexakis[12]DBA, a substructure motif of graphyne. In addition, the structure and magnetic properties of the parent compound 2a were investigated by DFT calculations. We also planned to prepare tris[12]DBA 3, having peripheral double bonds with internal hydrogens that should allow better assessment of the effect of the multiple fusion on the tropicity on the [12]DBA ring.

DFT calculations (B3LYP/6-31G* level) of 2a predict that it possesses a D_3 symmetric structure in which the three 12membered rings are twisted slightly (with a dihedral angle of 7.5° between the peripheral triple bond and the bond shared by the 12-membered ring and the central benzene ring) because of the steric repulsion between the hydrogen atoms of the peripheral benzene rings. The central benzene ring adopts a very shallow chair conformation with a dihedral angle of 0.8° (C(1)-C(2)-C(3)-C(4) = 0.838° , C(2)- $C(3)-C(4)-C(5) = -0.840^{\circ}$; see Figure 1 for numbering. In general, fusion of the antiaromatic [12]annulene ring to a benzene ring causes bond length alternation in the latter, in such a way that the endocyclic bond shared by the [12]annulene ring becomes longer than that of the exocyclic bond.¹³ It is expected that the fusion of three [12]annulene rings to the central benzene ring of 2a in an alternate fashion

(4) (a) Baughman, R. H.; Eckhardt, H.; Kerteszz, M. J. Chem. Phys. **1987**, 87, 6687. (b) Narita, N.; Nagai, S.; Suzuki, S.; Nakano, K. Phys. *Rev. B* **1998**, 58, 11009. (c) Narita, N.; Nagai, S.; Suzuki, S.; Nakano, K. *Phys. Rev. B* **2000**, 62, 11146.

(7) (a) Marsden, J. A.; O'Chilor, M. J.; Haley, M. M. *Org. Lett.* **2004**, 6, 2385. (b) Marsden, J. A.; Miller, J. J.; Haley, M. M. *Angew. Chem., Int. Ed.* **2004**, *43*, 1694.

(8) Staab, H. A.; Graf, F. Tetrahedron Lett. 1966, 7, 751.



Figure 1. Structures and numbering of the internal benzene ring of **2a,b** and the aromatic protons of tris[12]DBAs and their corresponding reference compounds.

would emphasize the bond length alternation. However, this is not the case because the calculated bond lengths of the internal benzene ring of 2a (C(2)-C(3), C(4)-C(5), C(1)-C(6)) are slightly longer than those of **1a** and the peripheral ring of 2a presumably for the above-mentioned steric repulsion.¹⁴ The other calculated bond lengths of 2a, however, are similar to those of 1a. The chemical shift of the aromatic protons and nucleus-independent chemical shifts (NICS)¹⁵ of **1a** and **2a** were calculated using the GIAO-HF/ 6-31G*//B3LYP/6-31G* methods.16 The NICS values of the 12-membered ring (4.41) and benzene rings (-6.79) of **1a** using the GIAO-B3LYP/6-31G*//B3LYP/6-31G* methods were reported by Vollhardt et al.¹⁷ Our results are summarized in Table 1 together with the experimental data for the *tert*-butyl derivatives **1b** and **2b** described later. Despite the smaller bond length alternation of the central benzene ring of 2a, the less negative NICS value (-7.83) indicates that this ring is substantially less aromatic than those of 1a (-10.35) and the peripheral benzene rings of **2a** (-10.25). Because of the slight twisting of the [12]annulene rings of 2a, they are slightly less paratropic than that of 1a, and the chemical shifts of peripheral benzene protons are predicted to resonate at a lower field than those of 1a.

As we wished to extend our synthetic strategy to the synthesis of the full wheel-shaped hexakis[12]DBA, we avoided using cross-coupling reactions, which are extensively employed to construct [12]DBA structures, to form the peripheral triple bonds. To introduce the peripheral triple

⁽⁵⁾ Zhou, Y.; Feng, S. Solid State Commun. 2002, 122, 307.

^{(6) (}a) Wan, W. B.; Brand, S. C.; Pak, J. J.; Haley, M. M. Chem.-Eur. J. 2000, 6, 2044. (b) Wan, W. B.; Haley, M. M. J. Org. Chem. 2001, 66,

^{3893. (}c) Marsden, J. A.; Haley, M. M. J. Org. Chem. 2005, 70, 10213. (7) (a) Marsden, J. A.; O'Cnnor, M. J.; Haley, M. M. Org. Lett. 2004,

⁽⁹⁾ For example, see: (a) Ferrara, J. D.; Tessier-Youngs, C. A.; Youngs, W. J. J. Am. Chem. Soc. **1985**, 107, 6719. (b) Ferrara, J. D.; Djebli, A.;

Tessier-Youngs, C. A.; Youngs, W. J. J. Am. Chem. Soc. 1988, 110, 647. (10) Wirz, J. In Excited States in Organic Chemistry and Biology; Pullman, B., Goldblum, N., Eds.; D. Reidal Publishing Company: Dordrecht, 1977; p 283.

^{(11) (}a) Koehoe, J. M.; Kiley, J. H.; English, J. J.; Johnson, C. A.; Petersen, R. C.; Haley, M. M. Org. Lett. **2000**, 2, 969. (b) Sonoda, M.; Sakai, Y.; Yoshimura, T.; Tobe, Y.; Kamada, K. Chem. Lett. **2004**, 33, 972.

^{(12) (}a) Iyoda, M.; Sirinintasak, S.; Nishiyama, Y.; Vorasingha, A.;
Sultana, F.; Nakao, K.; Kuwatani, Y.; Matsuyama, H.; Yoshida, M.; Miyake,
Y. Synthesis 2004, 1527. (b) Miljanić, O. Š.; Vollhardt, K. P. C.; Whitener,
G. D. Synlett 2003, 29.

⁽¹³⁾ Mitchell, R. H.; Zhang, R.; Fan, W.; Berg, D. J. J. Am. Chem. Soc. 2005, 127, 16251.

⁽¹⁴⁾ Calculated bond lengths of 1a and 2a are listed in Table S1 in the Supporting Information.

⁽¹⁵⁾ Schleyer, P. v. R.; Maerker, C.; Deansfeld, A.; Jial, H.; Hommes, N. J. R. v. E. J. Am. Chem. Soc. **1996**, *118*, 6317.

⁽¹⁶⁾ The NICS value of the 12-membered ring of 2a using GIAO-B3LYP/6-31G*// B3LYP/6-31G* methods (3.57) was also smaller than that of 1a (4.41).

⁽¹⁷⁾ Matzger, A. J.; Vollhardt, K. P. C. Tetrahedron Lett. 1998, 39, 6791.

| Table 1. | Calculated and Experimental ¹ H NMR Chemical |
|------------|---|
| Shifts and | NICS Values for 1a , b and 2a , b ^{<i>a</i>} |

| | 1a | 2a | | |
|-------------------------|------------|------------|--|--|
| ¹ H NMI | | | | |
| H^{1} | 7.56(7.26) | 7.62(7.38) | | |
| H^2 | 7.36(7.20) | 7.45(7.29) | | |
| H^3 | | 7.45 | | |
| H^4 | (7.39) | 7.90(7.41) | | |
| NICS^{c} | | | | |
| peripheral benzene ring | -10.35 | -10.25 | | |
| central benzene ring | | -7.83 | | |
| [12]annulene ring | 2.92 | 2.72 | | |
| | | | | |

^{*a*} Calculated values were obtained using the GIAO-HF/6-31G*//B3LYP/ 6-31G* method. ^{*b*} Experimental values observed for the *tert*-butyl derivatives **1b** and **2b** are listed in parentheses. ^{*c*} At the center of each ring.

bonds while avoiding severe steric crowding, we employed double elimination from an intermediate with peripheral single bonds and appropriate leaving groups. We viewed the coupling of a sulfone with an aldehyde¹⁸ and the pinacol coupling as reasonable routes to such an intermediate.

The syntheses of compounds **2b** and **3** are shown in Scheme 1. Triol **6** was prepared in 48% yield by the regioselective Sonogashira coupling of **4a** with hydroxyalkyne **5**.¹⁹ After conversion of **6** to sulfone **8**, the Sonogashira coupling with formylalkyne **9** gave **10** in 44% yield.²⁰ Treatment of **10** with lithium hexamethyldisilazide and diethyl chlorophosphonate followed by an excess amount of lithium diisopropylamide furnished trefoil-shaped tris[12]-DBA **2b** in 3% yield as a yellow solid. The low yield of **2b** is ascribed to intermolecular reactions because the nucleophilic C–C bond formation could take place randomly, leaving unreacted functional groups which are susceptible to further reactions.

Next, for the pinacol coupling approach, hexaaldehyde 11 was prepared in 72% yield by the Sonogashira coupling of hexabromobenzene (4b) with an excess of 9. Pinacol coupling of 11 using a low-valent vanadium reagent gave hexaol 12 as a mixture of isomers in 74% yield. The ¹H NMR spectrum of 12 indicates that the hydroxy groups are anti in each 12-membered ring. Chlorination of 12 using thionyl chloride gave hexachloride 13, in which the stereocenters possessed both syn and anti stereochemistry. Finally, double elimination using potassium tert-butoxide gave 2b in 10% yield. Compound 3 was prepared by the McMurry coupling of 11. Treatment of 11 with TiCl₄, Zn, and CuI in a mixture of DME and THF (1:2) gave 3 in 9% yield as a sparingly soluble orange solid. The low yield of 3 is again ascribed to the site-random coupling reaction as in the case of the cyclization of 10.



To assess the effect of the mode of fusion on the tropicity of the [12]DBA rings of **2b** and **3**, we compared their ¹H NMR chemical shifts with those of reference compounds 1b and 14.²¹ The chemical shifts of H¹, H², and H⁴ of 2b (7.38, 7.29, and 7.41 ppm, respectively) appeared at a slightly lower field than those of 1b (7.26, 7.20, and 7.39 ppm, respectively), indicating a small decrease of paratropicity in **2b** compared to **1b**. It seems likely that H¹ of **2b** suffers from the anisotropic effect of the neighboring benzene rings to show more pronounced downfield shifts than H² and H⁴. The observed chemical shifts agree qualitatively with the theoretical values obtained for the DFT optimized geometry of 2a (Table 1). It is, therefore, concluded that the observed downfield shifts are attributed to the slight twisting of the [12]annulene ring. Similarly, in the case of **3**, the chemical shifts of its aromatic protons (7.42, 7.19, and 7.31 ppm) shifted slightly downfield compared to the corresponding reference compound 14 (7.40, 7.13, and 7.29 ppm). Moreover, the vinyl protons of 3 and 14 serve as good probes for the assessment of their tropicity, which appear as sharp singlets owing to the fast rotation of the vinyl group. The

⁽¹⁸⁾ Orita, A.; Yoshioka, N.; Struwe, P.; Braier, A.; Beckmann, A.; Otera, J. *Chem.-Eur. J.* **1999**, *5*, 1355.

⁽¹⁹⁾ Sonoda, M.; Inaba, A.; Itahashi, K.; Tobe, Y. Org. Lett. 2001, 3, 2419.

⁽²⁰⁾ Compound **10** was synthesized only by this sequence of coupling reactions. Namely, the first coupling of **4a** with **9** followed by the second coupling with **5** or the phenyl sulfone derived from **5** did not give the desired hexaethynylbenzene derivatives.

⁽²¹⁾ Staab, H. A.; Bader, R. Chem. Ber. 1970, 103, 1157.

remarkably low field resonance of **14** (8.54 ppm) was ascribed to both the local anisotropic effect of the triple bonds to the inner hydrogen and the paratropicity of the 12-membered ring, either upfield or downfield shifting depending on the position (inner or outer) of the hydrogen.²¹ The observed upfield shift of the vinyl proton of **3** (8.42 ppm) relative to that of **14** indicates that the paratropicity of the former is smaller than that of the latter.

To investigate the excited-state properties of **2b** and **3**, UV-vis and fluorescence spectra were measured, as shown in Figures 2a and 2b, respectively, together with those of



Figure 2. (a) Electronic absorption spectra of **1a** (black), **2b** (red), **3** (blue), **14** (violet), and **15** (green) in CHCl₃ at 25 °C. (b) Normalized fluorescence spectra of **1a** (black), **2b** (red), **3** (blue), **14** (violet), and **15** (green) in CHCl₃ at 25 °C.

1a, **14**, and hexakis(phenylethynyl)benzene (**15**).²² It has been reported by Wirz that compound **1a** exhibits characteristic absorption/emission spectra owing to its unique electronic structure due to the D_{3h} symmetry of the molecule.¹⁰ There

is a strongly allowed absorption band at around 290 nm, a weakly allowed band at ca. 350 nm, and a strongly forbidden band at 400 nm. Even though **14** is not of D_{3h} symmetry, it also exhibits a similar absorption spectrum. On the other hand, the spectra of **2b** and **3** are different from those of **1a** and **14**, as shown in Figure 2a. Not only do **2b** and **3** exhibit bathochromic shifts compared to **1a** and **14** but also the intensities of the bands relative to those of **1a** and **14** are completely different. The difference between the absorption spectra can be ascribed to the loss of D_{3h} symmetry in **2b** and **3**. However, the profiles of the spectra of **2b** and **3** are different from that of a hexaethynylbenzene derivative **15**; the former exhibit absorptions at a longer wavelength owing to their extended conjugation.

The characteristic fluorescence spectrum of **1a** exhibits apparent vibrational splitting.¹⁰ The profile of the fluorescence spectrum of **2b** is different from that of **1a**, with the 0-0 transition band as the strongest band, a typical profile for phenylacetylene derivatives such as **15** (Figure 2b). The profiles of **3** and **14** are similar, though the fluorescence band of **3** exhibits a considerable red shift relative to that of **14**.²³

In conclusion, trefoil-shaped tris[12]DBA **2b** and **3** have been synthesized by constructing the 12-membered rings through peripheral single-bond formation using intramolecular coupling followed by double elimination. Their [12]annulene rings show slightly smaller paratropicities than that of the corresponding mono[12]DBAs **1a** and **14** presumably because of the decrease of planarity due to steric repulsion. Similarly, their excited-state properties are considerably different from those of **1a** and **14**.

Acknowledgment. Support to T.Y. from the 21st Century COE Program "Integrated Eco-Chemistry" is gratefully acknowledged. K.T. is grateful for a JSPS Research Fellowship for Young Scientists.

Supporting Information Available: Experimental procedures and spectral data of new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

OL060781U

⁽²²⁾ For the UV spectrum, see: Nierle, J.; Barth, D.; Kuck, D. Eur. J. Org. Chem. 2004, 867.

⁽²³⁾ Quantum yields of the fluorescence of **1a**, **2b**, **3**, **14**, and **15** were determined to be 0.10 (lit., 0.15 in methylcyclohexane at room temperature²⁴), 0.03, <0.01, 0.01, and 0.21 in CHCl₃ at 25 °C, respectively.

⁽²⁴⁾ Janecka-Styrcz, K.; Lipiński, J.; Ruziewicz, Z. J. Lumin. 1978, 17, 83.