

Preparation of Diaza Dioxo Dithia Crown *p*-*tert*-Butylcalix[4]arene by S-Alkylation Reactions: Use of Metallo-Calix[4]arene Complexes as Reaction Templates

Boosayarat Tomapatanaget, Buncha Pulpoka, and Thawatchai Tuntulani*
 Department of Chemistry, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand

(Received July 1, 1998; CL-980493)

New nickel-thiolato-calix[4]arene complexes were produced by mean of metal-template synthesis employing S-alkylation of a nickel thiolato complex with a dichloroethyl- or a dibromoethyl-calix[4]arene derivative. Demetallation of a nickel-thiolato-calix[4]arene complex resulted in a *p*-*tert*-butylcalix[4]arene derivative containing two N-, two O- and two S-donors.


Reports of closed-ring crown ether type derivatives of calix[4]arenes which possess sulfur donors are quite rare compared to those containing nitrogen and oxygen donors.¹ This scarcity may stem from the difficulty in handling the mercaptan ligands due to their facile oxidizability to disulfide or polymerization. One way to overcome this obstacle is to use a metal thiolate complex as template for synthesizing macrocyclic compounds containing sulfur donors usually by S-alkylation reaction. Classic examples of this method were described by Busch et al.²

A well known square planar nickel(II) thiolate, (bme-daco)Ni or **1**, containing two nitrogen and two sulfur donors synthesized by the Darensbourg group has been demonstrated its versatility as a bidentate metallo-thiolate ligand.³ The available lone pairs on the thiolate sulfurs of **1** can also act as nucleophiles towards alkyl halides to produce a number of macrocyclic nickel complexes which possess interesting Ni^{III/II} redox potential.⁴

We would like to prepare a calix[4]arene derivative containing a macrocyclic N-, O- and S- donors by modifying the lower rim OH moiety. On this account we employ **1** as a precursor for N- and S-atom donors and a nucleophile, and dichloroethyl-calix[4]arene (**4**) or dibromoethyl-calix[4]arene (**5**) as a precursor for an O-atom donor and alkylating agent. The synthetic methodology for **4** and **5** is represented by equation 1. Chloroethyl tosylate (**2**) and bromoethyl tosylate (**3**) were prepared from the reaction of 2-chloroethanol and 2-bromoethanol, respectively, with *p*-toluenesulfonylchloride in the


presence of triethylamine and a catalytic amount of 4-dimethylaminopyridine (DMAP). The nucleophilic substitution reaction between *p*-*tert*-butylcalix[4]arene and **2** (2 equivalence) in acetonitrile in the presence of K₂CO₃ then yielded the dichloroethyl-calix[4]arene precursor (**4**) which could be readily separated from the reaction mixture by precipitating with methanol (67%). However, the similar reaction of the bromo substituent **3** (2 equivalence) with *p*-*tert*-butylcalix[4]arene gave a mixture of products which were separated on a silica gel column. Using a 40% mixture of hexane/methylene chloride as eluant gave in order dibromoethyl-calix[4]arene (**5**) and calix[4]arene dimer (**6**) in 23% and 41% yield, respectively. ¹H NMR spectra and elemental analysis results of compounds **4**-**6** agree with the proposed structures.⁵

The alkylation reactions of the metallo nucleophile **1** with **4** and **5** are shown in equation 2. Under a nitrogen atmosphere, dropwise addition of purple acetonitrile solution of **1** into a mixture of **4** and NaI gradually shifted the color of the solution to dark red upon stirring at reflux for 48 h. The mixture was then separated on a silica gel column. Elution with a mixture of 15% MeOH/CH₂Cl₂ gave in order the yellowish green Ni-calix[4]arene complex (**8**) in 11% yield, an unidentified green complex⁶ and the bisNi-calix[4]arene complex (**9**) in 35% yield. The distribution of the products depends largely on the concentration of reagents. Increasing the concentration enhanced the yield of **9** and lowered that of **8**. On the other hand, a reaction of **1** with **5** resulted in higher yield (18%) of the green Ni-calix[4]arene complex (**7**). All reactions presented in equation 2 yielded the trimetallic complex which adhered on silica gel as a by-product.⁷



Complexes **8** and **9** were characterized by ^1H NMR and MALDI-TOF MS.⁸ Due to the paramagnetic contribution from the nickel centers in these complexes their ^1H NMR spectrum were broad in ~2-4 ppm region where signals of the protons from component **1** should appear. Nevertheless, some characteristic of the complexes can be deduced when compared the integral ratio of proton signals of calix[4]arene aromatic protons with overall proton signals appearing between 2-4 ppm which belong to the segment of **1**, the methylene bridges between the phenolic units and the ethano bridges on the lower rim of calix[4]arene. For **8**, the ^1H NMR spectrum shows two intense signals due to the protons of *tert*-butyl moieties and an AB system in the region of 3-4 ppm suggesting the cone conformation of the complex. The integral ratio of the signals of aromatic protons of the calix[4]arene component and the proton signals in 1.93-4.30 ppm region is 8:36 indicating the 1:1 proportion structure **8**. MALDI-TOF MS of **8** shows a molecular ion peak at 935.5 m/z which corresponds to its molecular mass when stripped off NiL_2 . This is not unusual since the thioether is a worse donor than the thiolate and suggests that the complex can readily lose the nickel ion to produce a free ligand. ^1H NMR spectrum of **9** shows that the integral ratio of the aromatic protons of calix[4]arene to the protons signals in 1.48-4.40 ppm region is 8:56 indicating the 1:2 ratio structure **9**. MALDI-TOF MS of **9** gives two molecular ion peaks at 1412.1 and 1227.2 m/z corresponding to the $[\text{C}_{68}\text{H}_{102}\text{O}_4\text{N}_4\text{S}_4\text{Ni}_2]^+$ ion and the $[\text{C}_{68}\text{H}_{103}\text{O}_4\text{N}_4\text{S}_4\text{Ni}]^+$ ion, respectively. These results thus agree with the proposed structures of **8** and **9**. We have tried to characterize **8** and **9** by elemental analysis, but the results are unacceptable probably due to mixed counter ions (Cl^- and I^-) in these two complexes (this stems from the preparation procedure which involved both Cl^- and I^- ions). However, the elemental analysis result of complex **7** that undoubtedly contains two Br^- counter ions agrees with the proposed structure accompanied with 2 water molecules.⁹ In addition, ^1H NMR spectrum of **7** is broader than analogue **8**. Although we cannot deduce information of the solution structure of **8**, the result suggests that the geometry around the nickel center may alter from a diamagnetic, relatively square planar, geometry with I^- ion to a paramagnetic, relatively distorted tetrahedral, geometry with Br^- ion.

The nickel ion can be readily removed from the Nicalix[4]arene complex (**8**). Upon addition of excess KCN into an acetonitrile solution of **8** resulted in immediate decolorization of the yellowish green solution. Separation of the free ligand from water-soluble inorganic products can be accomplished by $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$ extraction which yielded slightly yellow solid in the organic layer (92%). The solid was also very soluble in CH_3OH .



^1H NMR spectrum of the solid is consistent with the free ligand structure (**10**).¹⁰ The ligand contains two N-, two O- and two S-donors in a crown ether configuration suitable for further complexation studies.¹¹

Unfortunately, preparative procedure for **1** is quite complicated and expensive.¹² We are, therefore, currently trying a less

expensive nickel thiolates such as Busch's nickel Schiff base complexes to further investigate this reaction.

We appreciate Prof. Marcetta Y. Darensbourg for providing nickel complex **1**. The Bio-Technology Institute, Chulalongkorn University is gratefully acknowledged for MALDI-TOF mass spectrometry results. This work was supported by the National Science and Technology Development Agency (Grant No. CO-39-06-EMT-09-01).

References and Notes

- 1 S. Shinkai, *Tetrahedron*, **49**, 8933 (1993); V. Böhmer, *Angew. Chem. Int. Ed. Engl.*, **34**, 713 (1995); A. Pochini and R. Ungaro, in "Comprehensive Supramolecular Chemistry," ed by F. Vogtle, Pergamon Press, London (1996), Vol. 2, p. 103; A. Ikeda and S. Shinkai, *Chem. Rev.*, **97**, 1713 (1997).
- 2 M. C. Thompson and D. H. Busch, *J. Am. Chem. Soc.*, **86**, 2651 (1964); L. F. Lindoy and D. H. Busch, *Inorg. Chem.*, **13**, 2494 (1974).
- 3 D. K. Mills, Y. M. Hsiao, P. J. Farmer, E. V. Atnip, J. H. Reibenspies, and M. Y. Darensbourg, *J. Am. Chem. Soc.*, **113**, 1421 (1991); T. Tuntulani, J. H. Reibenspies, P. J. Farmer, and M. Y. Darensbourg, *Inorg. Chem.*, **31**, 3497 (1992); G. Musie, P. J. Farmer, T. Tuntulani, J. H. Reibenspies, and M. Y. Darensbourg, *Inorg. Chem.*, **35**, 2176 (1996).
- 4 M. Y. Darensbourg, I. Font, D. K. Mills, M. Pala, and J. H. Reibenspies, *Inorg. Chem.*, **31**, 4965 (1992); G. Musie, J. H. Reibenspies, and M. Y. Darensbourg, *Inorg. Chem.*, **37**, 302 (1998).
- 5 4: ^1H NMR (200 MHz, CDCl_3) δ 0.95 (s, 18H), 1.30 (s, 18H), 3.41 and 4.30 (dd, 8H, $J = 13$ Hz), 4.00 (t, 4H, $J = 6$ Hz), 4.30 (t, 4H, $J = 6$ Hz), 6.78 (s, 4H), 7.00 (s, 2H), 7.07 (s, 4H). Anal. Calcd. for $\text{C}_{44}\text{H}_{62}\text{O}_4\text{Cl}_2$: C, 74.49; H, 8.07%. Found: C, 74.57; H, 7.90%. 5: ^1H NMR (200 MHz, CDCl_3) δ 0.94 (s, 18H), 1.28 (s, 18H), 4.32 and 3.32 (dd, 8H, $J = 13$ Hz), 3.83 (t, 4H, $J = 6$ Hz), 4.27 (t, 4H, $J = 6$ Hz), 6.78 (s, 4H), 6.93 (s, 2H), 7.05 (s, 4H). Anal. Calcd. for $\text{C}_{44}\text{H}_{62}\text{O}_4\text{Br}_2$: C, 66.82; H, 7.24%. Found: C, 66.99; H, 7.19%.
- 6: ^1H NMR (200 MHz, CDCl_3) δ 0.98 (s, 36H), 1.25 (s, 36H), 3.36 and 4.51 (dd, 16H, $J = 13$ Hz), 4.60 (s, 8H), 6.85 (s, 8H), 7.00 (s, 8H), 7.63 (s, 4H). Anal. Calcd. for $\text{C}_{92}\text{H}_{116}\text{O}_8$: C, 81.86; H, 8.66%. Found: C, 81.86; H, 8.86%.
- 6 MALDI-TOF MS shows a molecular ion peak at 1026.0 m/z which is close to the molecular mass of the one arm substituted complex $[\text{C}_{58}\text{H}_{82}\text{O}_4\text{N}_2\text{S}_2\text{NiCl}]^+$ (MW = 1029.56 g/mol). However, we don't have other evidence to support the MS result since the ^1H NMR spectrum is too broad to deduce any information from it.
- 7 This nickel trimetallic complex composes of two units of **1** that acts as a bidentate ligand using S-donor to coordinate another Ni^{2+} ion in a square planar fashion. This complex is found in almost every reaction of **1**. See references 3 and 4.
- 8: ^1H NMR (200 MHz, CDCl_3) δ 0.87 (s, 18H), 1.29 (s, 18H), 1.93 (br m, 4H), 2.13 (br m, 4H), 3.20 and 4.30 (dd, 8H, $J = 13$ Hz), 3.54 (br, 8H), 3.63 (br, 2H), 3.80 (br, 2H), 4.05 (br m, 8H), 6.70 (s, 4H), 6.82 (s, 2H), 7.04 (s, 4H). MS (MALDI-TOF): 935.5 m/z ($[\text{M}-\text{NiL}_2]^+$). UV-vis (λ_{max}): 281 nm. 9: ^1H NMR (200 MHz, CDCl_3) δ 0.82 (s, 18H), 1.36 (s, 18H), 1.48-4.40 (br, 56H), 5.77 (s, 2H), 6.60 (s, 4H), 7.10 (s, 4H). MS (MALDI-TOF): 1412.1 m/z ($[\text{M}-\text{I}]^+$) and 1227.2 m/z ($[\text{M}-\text{NiL}_2]^+$). UV-vis (λ_{max}): 244 nm.
- 9 **7** can be crystallized by adding hexane into its CH_2Cl_2 solution. CH_2Cl_2 was then allowed to evaporate slowly to give green microcrystals of $\text{7} \cdot 2\text{H}_2\text{O}$. Anal. Calcd. for $\text{C}_{58}\text{H}_{82}\text{O}_4\text{S}_2\text{N}_2\text{Br}_2\text{Ni} \cdot 2\text{H}_2\text{O}$ ($\text{7} \cdot 2\text{H}_2\text{O}$): C, 58.54; H, 7.28; N, 2.35%. Found: C, 58.08; H, 7.05; N, 2.22%.
- 10 ^1H NMR (200 MHz, CD_3OD) δ 0.96 (s, 18H), 1.31 (s, 18H), 1.68 (m, 4H), 2.8 (m, 8H), 2.96 (m, 4H), 3.15 (m, 4H), 3.30 and 4.13 (dd, 8H, $J = 13$ Hz), 4.00 (m, 2H), 4.32 (m, 6H), 6.77 (s, 4H), 6.93 (s, 2H), 7.03 (s, 4H).
- 11 The N_2S_2 component can accommodate many transition metal ions such as Fe(II), Co(II), Ni(II), Cu(II), Zn(II), Pd(II), and Cd(II); T. Tuntulani, Ph.D. Thesis (1995), Texas A&M University, College Station, Texas.
- 12 D. K. Mills, I. Font, P. J. Farmer, Y.-M. Hsiao, T. Tuntulani, R. M. Buonomo, D. C. Goodman, G. Musie, C. A. Grapperhaus, M. J. Maguire, C.-H. Lai, M. L. Hatley, J. J. Smcc, J. A. Bellefeuille, and M. Y. Darensbourg, *Inorg. Synth.*, **32**, 89 (1998).