Platinum-catalyzed Aromatic C–H Silylation of Arenes with 1,1,1,3,5,5,5-Heptamethyltrisiloxane

Miki Murata,* Naoaki Fukuyama, Jun-ichi Wada, Shinji Watanabe, and Yuzuru Masuda Department of Materials Science, Kitami Institute of Technology, 165 Koencho, Kitami 090-8507

(Received May 16, 2007; CL-070535; E-mail: muratamk@mail.kitami-it.ac.jp)

The intermolecular dehydrogenative coupling of 1,1,1,3, 5,5,5-heptamethyltrisiloxane with arenes proceeded in the presence of a catalytic amount of platinum complexes prepared in situ from $PtCl_2$ and hydrotris(pyrazolyl)borate derivatives.

The versatility of arylsilanes in modern organic chemistry renders them attractive targets for synthesis. For example, the palladium-catalyzed Hiyama cross-coupling reaction of aryl-(alkoxy)silanes has become a powerful method for carbon-carbon bond formation.¹ Aryl(alkoxy)silanes can be easily prepared by the transition-metal-catalyzed silvlation of aryl halides with hydrosilanes, such as triethoxysilane² and 1,1,1,3,5,5,5heptamethyltrisiloxane (1).³ Clearly, from environmental and economical points of view, direct silvlation of aromatic C-H bonds of arenes 2 is a more attractive goal. Therefore, much attention has been focused on the C-H silvlation of aromatic hydrocarbons 2 using hydrosilanes as a silicon source, and numerous catalyst systems, such as $IrCl(CO)(PPh_3)_2$,⁴ $Pt(dba)_2$,⁵ $Tp^{Me2}Pt(Me)_2(H)$ ($Tp^{Me2} = hydrotris(3,5-dimethylpyrazolyl)$ borate),⁶ Cp*Rh(H)₂(SiEt₃)₂,⁷ and Ru₃(CO)₁₂,⁸ have been reported to affect this dehydrogenative coupling reaction. Above all, the platinum complexes are highly efficient catalysts for the hydrogen-acceptor-free processes; however, the silvlating reagents were restricted to triorganosilanes.^{5,6} Tetraorganosilicon reagents thus obtained are less reactive for the Hiyama crosscoupling reaction, mainly because electronegative groups, such as halogen and alkoxy group, on the silicon atom are required for the formation of penta-coordinated silicate intermediates.¹ Consequently, we have sought an alternative silvlating reagent which provided reactive arylsilanes;^{9,10} in this paper, wish to report a platinum-catalyzed C-H silvlation of arenes 2 with 1,1,1,3,5,5,5-heptamethyltrisiloxane (1; Eq 1).¹¹

$$Si^{-}H + H^{-}Ar \xrightarrow{PtCl_{2} / Tp^{Me2}K} Si^{-}Ar + H_{2}$$
(1)

$$1 \quad 2 (10 \text{ equiv}) \qquad 3$$

$$Me_{3}SiO_{Si^{-}} = Me_{3}SiO_{Si^{-}}$$

$$Me$$

An initial screen was performed using platinum precursors and additional (pyrazolyl)borate ligands for the dehydrogenative coupling of benzene (**2a**) and **1**. These salts are commercially available and can be easily handled in air. The results are summarized in Table 1. This coupling work was achieved with the aid of the Tp ligand; i.e., treatment of **1** (1 equiv.) with **2a** (10 equiv.) in the presence of 3 mol % of a metal complex prepared in situ from PtCl₂ and potassium hydrotris(pyrazolyl)borate (TpK) at 150 °C was found to lead to the corresponding phenyltrisiloxane **3a** in 80% yield (Entries 1 and 2). Under the conditions using an excess amount of **2a**, the formation of disilylated

Table 1. C–H silvlation of benzene (2a) with hydrosilanes^a

Entry	Hydrosilane	Ligand	Yield/% ^b
1	1	none	0
2	1	ТрК	80
3	1	BpK	35
4	1	Tp ^{Me2} K	93
5°	1	Tp ^{Me2} K	91
6	(Me ₃ SiO)Me ₂ SiH	Tp ^{Me2} K	46 ^d
7	Et ₃ SiH	Tp ^{Me2} K	37 ^e

^aReaction conditions: **1** (0.5 mmol), **2a** (5 mmol), PtCl₂ (0.015 mmol), ligand (0.015 mmol), 150 °C, 24 h. ^bGC yields of **3a** are based on **1**. ^cThe reaction was carried out at 200 °C using PtCl₂ (0.005 mmol) and Tp^{Me2}K (0.005 mmol). ^dGC yield of (Me₃SiO)Me₂SiPh. ^eGC yield of Et₃SiPh.

products was completely suppressed. Although dihydridobis(pyrazolyl)borate (Bp) was less efficient than Tp as supporting ligands under the same conditions (Entry 3), the use of $Tp^{Me2}K$ showed improved reactivity, providing a 93% yield of **3a** (Entry 4). The present reaction system is similar to but more mild than the previously reported $Tp^{Me2}Pt(Me)_2(H)$ -catalyzed C–H silylation.⁶ Additionally, it is noteworthy that this silylation was catalyzed by 1 mol % of the catalyst although the elevated temperature was required (Entry 5). Under our optimized conditions, the use of 1,1,1,3,3-pentamethyldisiloxane and triethylsilane induced a lowering of the reactivity (Entries 6 and 7). The present study demonstrates that **1** is an efficient silylating reagent for the aromatic C–H silylation.

The results obtained with representative arenes 2, giving aryltrisiloxanes 3 similarly as above, are listed in Table 2.12The yields and product ratios were determined by GC analysis of crude reaction mixtures. The reaction of mono-substituted arenes, such as trifluoromethylbenzene (2b), toluene (2c), and chlorobenzene (2d), resulted in a mixture of meta and para regioisomers and electronic characteristics of the substituent on 2 hardly affected the meta/para ratios (Entries 1-3). Electron-deficient arene 2b were more reactive than 2a.¹³ In these cases, the formation of ortho isomers was completely suppressed, whereas the reported platinum-catalyzed silvlation of 2c and 2d generated ortho isomers.^{5,6} In the reaction of anisole (2e), however, the product contained all regioisomers (Entry 4).⁵ Being consistent with these results, disubstituted arenes 2f-2h were silvlated regioselectively for steric reasons (Entries 5-7).^{6,7} In addition, the silvlation of 1,4-difluorobenzene (2i) occurred exclusively at the 2-position (Entry 8). Unfortunately, all attempts at the C-H silvlation of aniline and phenol were unsuccessful.

Finally, the potential versatility of the present silvlation was demonstrated by the one-pot synthesis of unsymmetrical biaryls, as we have reported that isolated 3-aryl-1,1,1,3,5,5,5-hepta-

Table 2. Isomer distributions for C-H silvlation^a



^aReaction conditions: **1** (0.5 mmol), **2** (5 mmol), catalyst (0.015 mmol), 200 °C, 24 h. ^bYield and isomer ratio were determined by GC analysis. ^cAt 150 °C. ^dA trace amount of **3a** (\approx 4%) was also obtained.

methyltrisiloxanes **3** participated in the palladium-catalyzed Hiyama cross-coupling.³ Thus, the silylation of benzene (**2a**) (20 equiv.) with **1** (2.0 equiv.) followed by removal of the volatile materials gave the crude product **3a**, which coupled with 4-bromo- α , α , α -trifluorotoluene (1.0 equiv.) in the presence of PdCl₂(PCy₃)₂ and TBAF to give the corresponding biaryl in 83% total yield (Eq 2).



In conclusion, 1,1,1,3,5,5,5-heptamethyltrisiloxane (1) was found to promote the platinum-catalyzed intermolecular dehydrogenative coupling with arenes 2. Since arylsiloxanes 3 thus obtained exhibited good reactivity for converting the silicon functionality, the present aromatic C–H silylation may be useful in the field of synthetic organic chemistry. Further investigations of this and related transition-metal-catalyzed C–H functionalizations are currently underway in our laboratory. This work was partially supported by Grant-in-Aid for Scientific Research on Priority Areas (No. 19027006, "Synergistic Effects for Creation of Functional Molecules") from MEXT.

References and Notes

- For recent reviews, see: a) T. Hiyama, E. Shirakawa, *Top. Curr. Chem.* 2002, 219, 61. b) C. J. Handy, A. S. Manoso, W. T. McElroy, W. M. Seganish, P. DeShong, *Tetrahedron* 2005, 61, 12201.
- 2 a) M. Murata, K. Suzuki, S. Watanabe, Y. Masuda, J. Org. Chem. 1997, 62, 8569. b) A. S. Manoso, P. DeShong, J. Org. Chem. 2001, 66, 7449. c) M. Murata, M. Ishikura, M. Nagata, S. Watanabe, Y. Masuda, Org. Lett. 2002, 4, 1843. d) M. Murata, H. Yamasaki, T. Ueta, M. Nagata, M. Ishikura, S. Watanabe, Y. Masuda, Tetrahedron 2007, 63, 4087.
- 3 M. Murata, K. Ota, H. Yamasaki, S. Watanabe, Y. Masuda, Synlett 2007, 1387.
- 4 W. A. Gustavson, P. S. Epstein, M. D. Curtis, Organometallics 1982, 1, 884.
- 5 Y. Uchimaru, A. M. M. El Sayed, M. Tanaka, Organometallics 1993, 12, 2065.
- 6 N. Tsukada, J. F. Hartwig, J. Am. Chem. Soc. 2005, 127, 5022.
- K. Ezbiansky, P. I. Djurovich, M. LaForest, D. J. Sinning, R. Zayes, D. H. Berry, *Organometallics* 1998, 17, 1455.
- 8 a) F. Kakiuchi, K. Igi, M. Matsumoto, N. Chatani, S. Murai, *Chem. Lett.* 2001, 422. b) F. Kakiuchi, K. Igi, M. Matsumoto, T. Hayamizu, N. Chatani, S. Murai, *Chem. Lett.* 2002, 396.
 c) F. Kakiuchi, M. Matsumoto, K. Tsuchiya, K. Igi, T. Hayamizu, N. Chatani, S. Murai, *J. Organomet. Chem.* 2003, 686, 134.
- 9 Quite recently, Miyaura et al. reported on an iridiumcatalyzed C-H silylation using 1-hydrosilatrane. See: E. Kishida, T. Saiki, T. Ishiyama, N. Miyaura, 53rd Symposium on Organometallic Chemistry, Osaka, Japan, September, 2006, Abstr. No., PB253; For an example of Hiyama coupling of 1-arylsilatranes, see: S. Riggleman, P. DeShong, J. Org. Chem. 2003, 68, 8106.
- 10 For examples of the aromatic C-H silylation of arenes with halodisilanes; see: a) T. Ishiyama, K. Sato, Y. Nishio, N. Miyaura, Angew. Chem., Int. Ed. 2003, 42, 5346. b) T. Saiki, Y. Nishio, T. Ishiyama, N. Miyaura, Organometallics 2006, 25, 6068.
- 11 1,1,1,3,5,5,5-Heptamethyltrisiloxane (1) is commercially available.
- 12 General procedure for the aromatic C–H silylation: $PtCl_2$ (15 µmol) and $Tp^{Me2}K$ (15 µmol) were placed in a resealable Schlenk tube. The tube was evacuated and backfilled with nitrogen and then charged with the arene 2 (5 mmol). After being stirred at room temperature for 1 h, 1,1,1,3,5,5,5-heptamethyltrisiloxane (1; 0.50 mmol) was added. The reaction mixture was then stirred at 200 °C for 24 h. After the reaction, the mixture was analyzed by GC and GC-MS. The volatile material was removed in vacuo, and the residue was purified by Kugelrohr distillation to afford the desired 3-aryl-1,1,1,3,5,5,5-heptamethyltrisiloxane **3**.
- 13 Relative product ratios from the reactions in equimolar mixtures of benzene (2a) and substituted arenes 2b, 2c, and 2e were 1:1.33, 1:0.61, and 1:0.40, respectively.