

# Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. Burugupalli, M. Richardson and S. J. Williams, *Org. Biomol. Chem.*, 2017, DOI: 10.1039/C7OB01786C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Organic and Biomolecular Chemistry

## ARTICLE

# Total synthesis and mass spectrometric analysis of a *Mycobacterium tuberculosis* phosphatidylglycerol featuring a two-step synthesis of (*R*)-tuberculostearic acid

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Satvika Burugupalli, Mark B. Richardson, Spencer J. Williams\*

We report the total synthesis of (*R*)-tuberculostearic acid-containing *Mycobacterium tuberculosis* phosphatidylglycerol (PG). The approach features a two-step synthesis of (*R*)-tuberculostearic acid, involving an (*S*)-citronellyl bromide linchpin, and the phosphoramidite-assisted assembly of the full PG structure. Collision-induced dissociation mass spectrometry of two chemically-synthesized PG acyl regioisomers revealed diagnostic product ions formed by preferential loss of carboxylate at the secondary (sn-2) position.


## Introduction

Phosphatidylglycerols are phospholipids produced by a wide range of organisms including bacteria, fungi, plants and animals.<sup>1</sup> Within animal cells PG is mainly found in the mitochondrial membranes,<sup>2</sup> and is believed to be released into the cellular milieu only upon cellular damage or stress. PGs are also produced by a wide range of bacteria, including important human pathogens such as *Listeria*, *Streptococcus*, and *Mycobacterium tuberculosis* (*Mtb*). PGs can be presented on several antigen presenting proteins of the cluster of differentiation 1 (CD1) class, specifically CD1b<sup>3</sup> and CD1d,<sup>4, 5</sup> where they are recognized by natural killer T cells and lipid-reactive T cells. Thus PGs appear to be a molecular signature allowing immune recognition of bacterial infection or damaged self.

In mycobacteria, PG is an intermediate in the biosynthesis of lysinylated PG,<sup>6</sup> phosphatidylinositol,<sup>7</sup> and cardiolipin.<sup>7, 8</sup> It has been proposed that PG is synthesized in two steps. Step 1 is catalyzed by phosphatidylglycerophosphate synthetase (PgsA3), which catalyzes the phosphatidylation of glycerol-3-phosphate by CDP-diacylglycerol to afford phosphatidylglycerophosphate,<sup>7</sup> and step 2 which involves the dephosphorylation of phosphatidylglycerophosphate to afford PG, by a phosphatidylglycerophosphatase. *Mtb* PG occurs as a range of lipoforms including an abundant species that bears (*R*)-tuberculostearic acid (*R*-TBSA) at the sn-1 position, and palmitic acid at the sn-2 position,<sup>4</sup> a decoration that matches that of other phosphatidyl-containing lipids<sup>9</sup> such as

phosphatidylethanolamine,<sup>10</sup> and phosphatidylinositol mannosides.<sup>11</sup> Possibly because of its role as a biosynthetic intermediate, *Mtb* PG occurs at low levels and the challenges of its isolation<sup>4</sup> from the natural source are compounded by the pathogenicity and low growth rate of *Mtb*. Recent work has shown that *Mtb* PG can be presented by the CD1d antigen presenting molecule to natural killer T cells that display 'type 2' T cell receptors.<sup>4</sup> Type 2 NKT cells are distinguished from their better studied type I variants, by lacking reactivity to the prototypical NKT ligand  $\alpha$ -galactosyl ceramide, and through possessing a diverse array of  $\alpha\beta$  chains that comprise the T cell receptor.<sup>12</sup> While most studies have focused on type I NKT cells, type II NKT cells are more abundant in humans, and are functionally distinct from type I NKT cells.<sup>13</sup> However, studies of type 2 NKT cells have been limited by a lack of well-defined lipid antigens.

Herein we report the total synthesis of the TBSA-containing *Mtb* PG (**1**) using phosphoramidite chemistry. As part of this effort we have devised the shortest synthesis yet reported of *R*-TBSA (**2**) in just two steps from commercially available (*S*)-citronellyl bromide, and its incorporation into *Mtb* PG. Finally, we demonstrate that collision-induced dissociation mass spectrometry of chemically-synthesized PG regioisomers results in the formation of diagnostic fragment ions that have utility in the structural characterization of PGs.



**Fig 1** Structure of *Mycobacterium tuberculosis* (*Mtb*) TBSA-containing phosphatidylglycerol (PG).

School of Chemistry and Bio21 Molecular Science and Biotechnology Institute, University of Melbourne, Parkville, Victoria, Australia 3010.

E-mail: sjwill@unimelb.edu.au Address here.











Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

## Organic and Biomolecular Chemistry

View Article Online  
DOI: 10.1039/C7OB01786C

## ARTICLE

Table 1. Previous synthetic approaches to (*R*)-tuberculostearic acid.

Starting material/key intermediate	Approach/steps	Ref.	Starting material/key intermediate	Approach/steps	Ref.
 ( <i>R</i> )-2-methyldecanol	Chiral resolution • 10 steps <sup>a</sup>	Prout, 1948 <sup>14</sup>	 bromocasylate	Chiral resolution • 7 steps	Hahn, 2015 <sup>15</sup>
 methyl ( <i>R</i> )-β-methylglutarate	Chiral resolution • 9 steps	Linstead, 1951 <sup>16</sup>	 isopropyl 10-undecenoate	Catalytic enantioselective induction • 6 steps <sup>b</sup>	Feringa, Minnaard, 2010 <sup>10</sup>
 ( <i>S</i> )-citronellol	Chiral pool • 8 steps	Larsen, 2007 <sup>17</sup>	 nonanal	Catalytic enantioselective induction • 5 steps	Minnaard, 2013 <sup>18</sup>
 Roche ester	Chiral pool • 7 steps	Seeberger, 2006 <sup>19</sup>	 Roche ester	Chiral pool • 5 steps	Hung, 2015 <sup>20</sup>
 ( <i>R</i> )-oxazolidinone	Chiral auxiliary • 7 steps	Williams, 2013 <sup>21</sup>	 ( <i>S</i> )-citronellyl bromide	Chiral pool • 4 steps	Baird, 2006 <sup>22</sup>


<sup>a</sup> (*R*)-2-Methyldecanol was synthesized from bromooctane in 5 steps by resolution of an intermediate brucine salt.<sup>b</sup> Isopropyl 10-undecenoate can be synthesized in one step from commercially-available undecylenic acid.

## Results and discussion

Synthesis of (*R*)-tuberculostearic acid

Numerous syntheses of *R*-TBSA have been reported, using chiral pool sources,<sup>17, 19, 20</sup> chiral auxiliaries,<sup>21</sup> catalytic enantioselective induction,<sup>10, 18</sup> and resolution methods<sup>14-16</sup> (Table 1). The most efficient synthesis reported to date is that of Roberts and Baird who prepared TBSA in four steps from (*S*)-citronellyl bromide, in a route involving stepwise elongations using copper(I)-catalyzed cross-coupling and Wittig extension.<sup>22</sup> We were inspired by this use of citronellyl bromide as a linchpin, and anticipated that diverting the cross-coupled intermediate into a Grubbs-cross-metathesis/hydrogenation process could provide further synthetic abbreviation. Thus, according to the reported method,<sup>22</sup> copper(I)-catalyzed cross-coupling of heptylmagnesium bromide with (*S*)-citronellyl bromide afforded the alkene **3** (Scheme 1). Initially, a 1:1 mixture of ethyl 6-heptenoate and **3** were reacted in the presence of Hoveyda-Grubbs II catalyst, affording a 58% yield of an *E/Z*


mixture of cross-coupled alkenes. We next attempted a direct tandem cross-metathesis/hydrogenation reaction. Use of a 1:1 mixture of alkenes afforded a 48% yield of ethyl *R*-TBSA, which could be improved to 76% by using a 3:1 ratio of ethyl heptenoate/**3**. Saponification of ethyl *R*-TBSA afforded *R*-TBSA (**2**) in three steps. Encouraged by these results, we next enacted a two-step synthesis of *R*-TBSA by employing a 3:1 ratio of 6-heptenoic acid/**3** under the same conditions, which directly afforded *R*-TBSA in 73% yield.

Scheme 1 Two-step synthesis of (*R*)-tuberculostearic acid.

## Synthesis of phosphatidylglycerols


Reported methods for the synthesis of PGs include chemoenzymatic methods,<sup>23</sup> and purely chemical approaches;<sup>24</sup> however, the reported chemical methods typically employ highly moisture sensitive POCl<sub>3</sub> as a phosphorylation reagent and do not allow regioselective introduction of two different acyl groups. In devising an approach to *Mtb* PG, we elected to utilize contemporary phosphoramidite chemistry, which has been widely applied in the synthesis of oligonucleotides,<sup>25</sup> phosphatidylinositols,<sup>26</sup> and cardiolipin.<sup>27</sup>

We chose to prepare **1**, and analogues lacking the 10-methyl group with alternate arrangements of the fatty acyl groups. For the construction of the requisite diacylglycerols, we chose to employ the approach of Minnaard, which proceeds with high regioselectivity and without acyl group migration.<sup>18</sup> Accordingly, opening of benzyl (*R*)-glycidyl ether (**5**) with palmitic acid, stearic acid, or *R*-TBSA using Jacobsen pre-catalyst **C1**,<sup>28</sup> followed by esterification with a second fatty acid and DCC/DMAP, gave the diesters **6a-c**. Hydrogenolysis of the benzyl ether afforded the diacylglycerols **7a-c**, which were used immediately without purification in the next step.



Scheme 2 Preparation of diacylglycerols.

Assembly of the PG structure relied upon the use of a phosphordiamidite linchpin, benzyl bis(*N,N*-diisopropyl)phosphordiamidite (Scheme 3).<sup>17, 29</sup> Reaction of the diacylglycerols **7a-c** with this phosphordiamidite, catalyzed by diisopropyl ammonium tetraazolid (DIPAT), a weak acid that does not activate the product,<sup>25</sup> afforded the unstable phosphoramidites **8a-c**. Compounds **8a-c** were purified by a quick passage over silica gel to remove any residual reactant or reagents, and then immediately subjected to a second phosphorylation reaction with dibenzylglycerol **9**, catalyzed by 1*H*-tetrazole. Upon consumption of the phosphoramidate, *t*-BuOOH was added to oxidize the intermediate phosphite to the protected PGs **10a-c**. Finally, hydrogenolysis of the benzyl groups provided *Mtb* PG **1**, and analogues **11a** and **11b**.

Scheme 3 Preparation of *Mtb* PG **1** and analogues.

## Collision-induced dissociation mass spectrometry of phosphatidylglycerols


Early studies reported that negative-ion fast-atom bombardment or electrospray-ionization with collision-induced dissociation/tandem mass spectrometry of phosphatidylethanolamine (PE) and PG result in fragmentation to provide high abundance fatty acid carboxylate (RCO<sub>2</sub><sup>-</sup>) fragment ions.<sup>30, 31</sup> These studies revealed similar fragmentation pathways for PE and PG, and showed that regioisomeric PEs fragment in a characteristic way to produce RCO<sub>2</sub><sup>-</sup> ions for which those derived from the sn-2 position are of greater intensity, thereby allowing structural characterization of fatty acyl regiochemistry.<sup>10, 31</sup> Subsequently, it was shown that CID/MS-MS of PG occurs with similar results, producing RCO<sub>2</sub><sup>-</sup> ions in which those from the sn-2 position are more abundant, in what has been described as a charge-driven process.<sup>32, 33</sup> In order to confirm these fragmentation patterns we studied the chemically synthesized regioisomers **11a** and **11b** negative ion CID/MS<sup>2</sup> experiments using electro-spray ionization. Fragmentation of the parent anions [M-H]<sup>-</sup> (*m/z* 749.5) formed from the regioisomers **11a** (C<sub>16</sub>/C<sub>18</sub>) and **11b** (C<sub>18</sub>/C<sub>16</sub>), yielded characteristic fatty acid carboxylate (RCO<sub>2</sub><sup>-</sup>) fragment ions at *m/z* 255.2 and 283.3 (Figure 2a,b). In each case the more abundant ion is that derived from the fatty acid located in the sn-2 position. This work confirms that CID-induced fragmentation patterns can be used to identify the structures of asymmetrical PG molecules. Figure 2c shows the CID-MS<sup>2</sup> spectrum of **1**, revealing that fragmentation of the parent ion [M-H]<sup>-</sup> (*m/z* 763.5) affords daughter ions at *m/z* 255.2 and 297.3. Here, the most abundant fragment (*m/z* 255.2) is that derived from the palmitic acid located at the sn-2 position. We note that these and earlier observations for CID-

## ARTICLE

## Organic Biomolecular Chemistry

MS of PGs differ from those for the structurally-related phosphatidylinositols<sup>32</sup> and phosphatidylinositolmannosides,<sup>11, 17</sup> which under similar conditions fragment by neutral loss of fatty acids and fatty acid ketenes in a diagnostic way. While these systems also yield  $\text{RCO}_2^-$  fragment ions, their relative intensities do not reflect the regiochemistry of their location on the phosphatidyl group because other pathways also yield these ions in abundances that are sensitive to the applied collision energy.<sup>32</sup>

presented lipid antigen, other closely related PGs from mammals, *Escherichia coli*, *Corynebacterium glutamicum* and *Listeria monocytogenes* have been identified as CD1b-presented lipid antigens.<sup>3</sup> Access to synthetic *Mtb* PG should support studies to understand its immunological properties, and investigation of the effect of the branched-methyl group in the fatty acyl chain. Collision-induced dissociation mass spectrometry analysis confirms the desired acylation patterns in our synthetically acquired materials, and provides a foundation for structural analyses of mycobacterial PGs.



**Fig 2** Multistage mass spectrometry ( $\text{MS}^2$ ) experiments involving the collision-induced dissociation of negative ions formed by electro-spray ionization.† (a)  $\text{MS}^2$  spectrum of deprotonated **11a** ( $\text{C}_{16}/\text{C}_{18}$ ). (b)  $\text{MS}^2$  spectrum of deprotonated **11b** ( $\text{C}_{18}/\text{C}_{16}$ ). (c)  $\text{MS}^2$  spectrum of deprotonated **1**. (d) Fragmentation of compound **1**. See Experimental for details.

## Conclusions

In this work we disclose the first total synthesis of *Mtb* PG **1**, and analogues **11a** and **11b**. Our approach includes the most concise preparation of *R*-TBSA reported to date. We note that while *Mtb* PG has only been studied in the context of a CD1d-

## Experimental

## General methods

Proton ( $^1\text{H}$  NMR, 400 or 500 MHz), proton-decoupled carbon-13 ( $^{13}\text{C}$  NMR, 100 or 125 MHz), and phosphorus-31 ( $^{31}\text{P}$  NMR, 202 MHz) nuclear magnetic resonance spectra were obtained in deuteriochloroform, methanol- $\text{d}_4$  ( $\text{CD}_3\text{OD}$ ) or  $\text{DMSO}-\text{d}_6$  with residual protonated solvent or solvent carbon signals as internal standards. Abbreviations for multiplicity are s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet. Flash chromatography was carried out on silica gel 60 according to the procedure of Still *et al.*<sup>34</sup> Analytical thin layer chromatography (t.l.c.) was conducted on aluminium-backed 2 mm thick silica gel 60 F<sub>254</sub> and chromatograms were visualized with ceric ammonium molybdate (Hanessian's stain), potassium permanganate or 5%  $\text{H}_2\text{SO}_4/\text{MeOH}$ , with charring as necessary. High resolution mass spectra (HRMS) were obtained using an ESI-TOF-MS; all samples were run using 0.1% formic acid. Dry  $\text{CH}_2\text{Cl}_2$ , THF, and  $\text{Et}_2\text{O}$  were obtained from a dry solvent apparatus (Glass Contour of SG Water, Nashua, U.S.A.) as per the procedure of Pangborn *et al.*<sup>35</sup> Dry DMF was dried over 4 Å molecular sieves. Pet. spirits refers to petroleum ether, boiling range 40–60 °C.

1. Synthesis of (*R*)-tuberculostearic acid (**2**)(*R*)-2,6-Dimethyltetradecene (**4**)

See: Roberts and Baird.<sup>22</sup> Hexylmagnesium bromide (3.14 ml, 4.56 mmol) (2.0 M in  $\text{Et}_2\text{O}$ ) was added to a solution of (*S*)-citronellyl bromide (0.23 ml, 1.14 mmol) in dry  $\text{Et}_2\text{O}$  (2.3 ml) at  $-78^\circ\text{C}$  followed by dilithium tetrachlorocuprate ( $\text{Li}_2\text{CuCl}_4$ ) (0.1 M in THF) (1.94 ml, 0.12 mmol). The solution was allowed to warm to room temperature and stirred for 36 h. The reaction mixture was quenched with sat aq.  $\text{NH}_4\text{Cl}$ , extracted with  $\text{Et}_2\text{O}$ , and the organic phase washed with brine, dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure. Flash chromatography (0.5%  $\text{Et}_2\text{O}$ :petroleum ether) of the residue afforded **4** as a colorless oil (240 mg, 95%);  $[\alpha]_{\text{D}}^{22} +0.75$  (c 0.5 in  $\text{CHCl}_3$ ) (lit.<sup>22</sup>  $[\alpha]_{\text{D}}^{25} +1.5$  in  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  0.88–0.83 (6 H, m,  $2 \times \text{CH}_3$ ), 1.15–1.06 (2 H, m,  $\text{CH}_2$ ), 1.31–1.22 (20 H, m,  $10 \times \text{CH}_2$ ), 1.60 (3 H, s,  $\text{CH}_3$ ), 1.68 (3 H, s,  $\text{CH}_3$ ), 1.95 (2 H, dd,  $J = 6.8, 13.7$  Hz,  $\text{CHCH}_2$ ), 5.10 (1 H, t,  $J = 7.0$  Hz,  $\text{CH}=\text{C}$ );  $^{13}\text{C}$  NMR (101 MHz,  $\text{CDCl}_3$ )  $\delta$  14.2, 17.7, 19.2, 22.8, 25.7, 27.1, 29.5, 29.8, 30.1, 32.0, 32.5, 37.1, 37.3, 125.2, 131.0; HRMS calcd for  $\text{C}_{16}\text{H}_{32}$   $[\text{M}+\text{H}]^+$  225.2538, found 225.2534.



**Ethyl (*R*)-10-methyloctadecanoate**

Hoveyda-Grubbs 2<sup>nd</sup> generation catalyst (HG-II) (2.8 mg, 4 mol%) was added to a solution of **4** (250 mg, 0.11 mmol) and ethyl heptenoate (59 mg, 0.33 mmol) in dry dichloromethane (1.14 ml) and stirred for 24 h under reflux. Upon consumption of starting material, the reaction mixture was concentrated under reduced pressure. EtOAc (5.0 ml) was added to the residue and the solution was subjected to hydrogenation under high pressure (37 bar) in a Buchi® hydrogenator overnight at room temp after which the solvent was removed under reduced pressure. Flash chromatography (2% Et<sub>2</sub>O/petroleum spirits) of the crude mixture afforded the title compound as a colourless oil (28 mg, 76%);  $[\alpha]_D^{25} -0.14$  (*c* 0.5 in CHCl<sub>3</sub>) (lit.<sup>14</sup>  $[\alpha]_D^{25} -0.16$  in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.82 (3 H, d, *J* = 6.5 Hz, CHCH<sub>3</sub>), 0.87 (3 H, t, *J* = 6.8 Hz, CH<sub>3</sub>), 1.10–1.03 (2 H, m, CH<sub>2</sub>), 1.31–1.20 (28 H, m, 12  $\times$  CH<sub>2</sub>, 1  $\times$  CH, COOCH<sub>2</sub>CH<sub>3</sub>), 1.61 (2 H, d, *J* = 7.1 Hz, COCH<sub>2</sub>CH<sub>2</sub>), 2.27 (2 H, t, *J* = 7.6 Hz, COCH<sub>2</sub>), 4.11 (2 H, d, *J* = 7.1 Hz, C=OCH<sub>2</sub>); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 14.4, 19.8, 22.8, 25.1, 27.1, 27.2, 29.3, 29.4, 29.5, 29.6, 29.8, 30.0, 30.1, 32.0, 32.8, 32.8, 34.5, 37.2, 60.2, 174.0 (C=O); HRMS calcd for C<sub>21</sub>H<sub>42</sub>O<sub>2</sub> [M+H]<sup>+</sup> 327.3218, found 327.3220.

**(*R*)-Tuberculostearic acid ((*R*)-10-methyloctadecanoic acid) (**2**)**

**(a) From ethyl (*R*)-10-methyloctadecanoate:** 2 M KOH (0.76 ml, 1.5 mmol) was added to a solution of the ethyl ester (50 mg, 0.153 mmol) in MeOH/THF (1:1, 0.8 ml) and stirred under reflux overnight. The solution was acidified with aq citric acid (1 M), extracted with EtOAc, and the organic phase washed with brine, dried (MgSO<sub>4</sub>) and concentrated under reduced pressure to afford **2** as a colorless oil (44 mg, 97%);  $[\alpha]_D^{25} -0.9$  (*c* 1.0 in CHCl<sub>3</sub>) (lit.<sup>22</sup>  $[\alpha]_D^{25} -0.34$  in CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.83 (3 H, d, *J* = 6.5 Hz, CHCH<sub>3</sub>), 0.88 (3 H, t, *J* = 6.8 Hz, CH<sub>2</sub>CH<sub>3</sub>), 1.12–1.04 (2 H, m, CH<sub>2</sub>), 1.36–1.21 (25 H, m, 12  $\times$  CH<sub>2</sub>, 1  $\times$  CH), 1.67–1.59 (2 H, m, OCOCH<sub>2</sub>CH<sub>2</sub>), 2.35 (2 H, t, *J* = 7.5 Hz, OCOCH<sub>2</sub>); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 19.8, 22.8, 24.8, 27.1, 27.2, 29.2, 29.4, 29.5, 29.6, 29.8, 30.0, 30.1, 32.0, 32.8, 34.0, 37.2, 179.5; HRMS calcd for C<sub>18</sub>H<sub>36</sub>O<sub>2</sub> [M+H]<sup>+</sup> 285.2749, found 285.2747.

**(b) From alkene **4**:** HG-II (2.80 mg, 4 mol%) was added to a solution of **4** (25 mg, 0.114 mmol) and heptenoic acid (59 mg, 0.334 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (1.14 ml) and stirred for 24 h under reflux. The reaction mixture was concentrated under reduced pressure. EtOAc (5.0 ml) was added to the residue and the solution was subjected to hydrogenation under high pressure (37.5 bar) in a Buchi® hydrogenator overnight at room temperature after which solvent was removed under reduced pressure. Flash chromatography (2% Et<sub>2</sub>O/petroleum spirits) of the crude mixture afforded **2** as a colorless oil (25 mg, 73%).

**2. Synthesis of diacylglycerols****3-*O*-Benzyl-1-*O*-((*R*)-10-methyloctadecanoyl)-2-*O*-palmitoyl-sn-glycerol (**6c**)**

A solution of (*R*)-TBSA **2** (100 mg, 335  $\mu$ mol) and Co[salen] **C1** (4.0 mg, 2 mol%) in Et<sub>2</sub>O (1 ml) was stirred under an

oxygen balloon for 10 min. The solvent was evaporated under vacuum to afford a brown residue. THF (0.2 ml) and Hunig's base (58  $\mu$ l, 372  $\mu$ mol) was added to the residue, and the mixture was stirred for 5 min. (*R*)-Glycidyl benzyl ether (56.7  $\mu$ l, 372  $\mu$ mol) was added and the resulting mixture was stirred overnight at room temperature followed by removal of solvents in vacuo. Heptane (800  $\mu$ l), palmitic acid (114 mg, 446  $\mu$ mol) and DMAP (2.27 mg, 18.6  $\mu$ mol, 5 mol%) were added to the resulting residue. The mixture was cooled to 0 °C, followed by addition of DCC (91.9 mg, 446  $\mu$ mol) and stirred overnight. The solvent was removed in vacuo and the residue was subjected to flash chromatography (10% Et<sub>2</sub>O:pentane) to afford the title compound as a colorless thick liquid (178 mg, 77%);  $[\alpha]_D^{25} +6.2$  (*c* = 0.5, CHCl<sub>3</sub>),  $[\alpha]_D^{23}$  lit.<sup>19</sup> +5.45 (*c* = 3.9, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.83 (3 H, d, *J* = 6.5 Hz, CCH<sub>3</sub> (TBSA)), 0.88 (6 H, t, *J* = 6.8 Hz, 2  $\times$  CH<sub>3</sub>), 1.07 (2 H, m, CH<sub>2</sub> (TBSA)), 1.25 (48 H, m, 24  $\times$  CH<sub>2</sub>), 1.60 (4 H, m, 2  $\times$  COCH<sub>2</sub>CH<sub>2</sub>), 2.25–2.34 (4 H, m, 2  $\times$  COCH<sub>2</sub>), 3.59 (2 H, d, *J* = 5.2 Hz, H<sub>3a,3b</sub>), 4.19 (1 H, dd, *J* = 11.9, 6.4 Hz, H<sub>1a</sub>), 4.34 (1 H, dd, *J* = 11.9, 3.8 Hz, H<sub>1b</sub>), 4.50–4.58 (2 H, m, CH<sub>2</sub>Ph), 5.24 (1 H, m, H<sub>2</sub>), 7.27–7.37 (5 H, m, Ph); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 19.8, 22.8, 25.0, 25.1, 27.2, 29.2, 29.2, 29.4, 29.5, 29.6, 29.6, 29.8, 30.1, 30.1, 32.0, 32.9, 34.2, 34.4, 37.2, 62.8, 68.4, 70.1, 73.4, 127.7, 127.9, 128.5, 137.8 (Ph), 173.2, 173.5 (C=O); HRMS: calcd for C<sub>45</sub>H<sub>80</sub>O<sub>5</sub> [M + H]<sup>+</sup> 701.6039, found 701.6038.

**3-*O*-Benzyl-1-*O*-palmitoyl-2-*O*-stearoyl-sn-glycerol (**6a**)**

Synthesized as for **6c**, 250 mg (0.36 mmol, 66%).  $[\alpha]_D^{25} +5.6$  (*c* = 0.5, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (6 H, t, *J* = 6.8 Hz, 2  $\times$  CH<sub>3</sub>), 1.27 (48 H, m, 24  $\times$  CH<sub>2</sub>), 1.60 (4 H, m, 2  $\times$  COCH<sub>2</sub>CH<sub>2</sub>), 2.25–2.34 (4 H, m, 2  $\times$  COCH<sub>2</sub>), 3.59 (2 H, d, *J* = 5.2 Hz, H<sub>3a,3b</sub>), 4.19 (1 H, dd, *J* = 11.9, 6.4 Hz, H<sub>1a</sub>), 4.34 (1 H, dd, *J* = 11.9, 3.8 Hz, H<sub>1b</sub>), 4.50–4.58 (2 H, m, CH<sub>2</sub>Ph), 5.24 (1 H, m, H<sub>2</sub>), 7.28–7.37 (5 H, m, Ph); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 22.8, 25.0, 25.1, 29.2, 29.2, 29.4, 29.5, 29.6, 29.8, 29.8, 32.0, 34.2, 34.4, 62.8, 68.4, 70.1, 73.4, 127.7, 127.9, 128.5, 137.8, 173.2, 173.5; HRMS: calcd for C<sub>44</sub>H<sub>78</sub>O<sub>5</sub> [M + H]<sup>+</sup> 687.5883, found 687.5884.

**3-*O*-Benzyl-1-*O*-stearoyl-2-*O*-palmitoyl-sn-glycerol (**6b**)**

Synthesized as for **6c**, 62 mg (0.093 mmol, 64%).  $[\alpha]_D^{25} +5.8$  (*c* = 0.5, CHCl<sub>3</sub>), lit.<sup>36</sup>  $[\alpha]_D^{23} +5.4$  (*c* = 1.63, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (6 H, t, *J* = 6.8 Hz, 2  $\times$  CH<sub>3</sub>), 1.27 (48 H, m, 24  $\times$  CH<sub>2</sub>), 1.60 (4 H, m, 2  $\times$  COCH<sub>2</sub>CH<sub>2</sub>), 2.25–2.34 (4 H, m, 2  $\times$  COCH<sub>2</sub>), 3.59 (2 H, d, *J* = 5.2 Hz, H<sub>3a,3b</sub>), 4.19 (1 H, dd, *J* = 11.9, 6.4 Hz, H<sub>1a</sub>), 4.34 (1 H, dd, *J* = 11.9, 3.8 Hz, H<sub>1b</sub>), 4.50–4.58 (2 H, m, CH<sub>2</sub>Ph), 5.24 (1 H, m, H<sub>2</sub>), 7.28–7.37 (5 H, m, Ph); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 22.8, 25.0, 25.1, 29.2, 29.2, 29.4, 29.5, 29.6, 29.8, 29.8, 32.0, 34.2, 34.4, 62.8, 68.4, 70.1, 73.4, 127.7, 127.9, 128.5, 137.8 (Ph), 173.2, 173.5 (C=O); HRMS: calcd for C<sub>44</sub>H<sub>78</sub>O<sub>5</sub> [M + H]<sup>+</sup> 687.5883, found 687.5884.

**1-*O*-((*R*)-10-Methyloctadecanoyl)-2-*O*-palmitoyl-sn-glycerol (**7c**)**

A mixture of **6c** (65 mg, 0.089 mmol) and Pd(OH)<sub>2</sub> on carbon (20%, 40 mg, 0.285 mmol) in ethanol (3 ml) was stirred under

## ARTICLE

## Organic Biomolecular Chemistry

hydrogen at 10 bar for 2 h. The reaction mixture was filtered through Celite and the solvent removed in vacuo to afford the product as a colorless liquid (61 mg, 99%). The compound was used directly without delay or purification.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  0.83 (3 H, d,  $J$  = 6.5 Hz,  $\text{CCH}_3$  (TBSA)), 0.88 (6 H, t,  $J$  = 6.8 Hz,  $2 \times \text{CH}_3$ ), 1.08 (2 H, m,  $\text{CH}_2$  (TBSA)), 1.27 (48 H, m,  $24 \times \text{CH}_2$ ), 1.62 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.30–2.37 (4 H, m,  $2 \times \text{COCH}_2$ ), 3.73 (2 H, d,  $J$  = 5.2 Hz, H3a,3b), 4.24 (1 H, dd,  $J$  = 11.9, 5.6 Hz, H1a), 4.32 (1 H, dd,  $J$  = 11.9, 4.6 Hz, H1b), 5.06–5.11 (1 H, m, H2).

**1-O-Palmitoyl-2-O-stearoyl-sn-glycerol (7a)**

Synthesized as for **7c**, 30.4 mg (98%).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.88 (6 H, t,  $J$  = 7.0 Hz,  $2 \times \text{CH}_3$ ), 1.27–1.34 (48 H, m,  $24 \times \text{CH}_2$ ), 1.62 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.30–2.37 (4 H, m,  $2 \times \text{COCH}_2$ ), 3.73 (2 H, dd,  $J$  = 5.0, 1.4 Hz, H3a,3b), 4.23 (1 H, dd,  $J$  = 11.9, 5.7 Hz, H1a), 4.32 (1 H, dd,  $J$  = 11.9, 4.5 Hz, H1b), 5.06–5.11 (1 H, m, H2).

**1-O-Stearoyl-2-O-palmitoyl-sn-glycerol (7b)**

Synthesized as for **7c**, 40.2 mg (96%).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.88 (6 H, t,  $J$  = 7.0 Hz,  $2 \times \text{CH}_3$ ), 1.27–1.34 (48 H, m,  $24 \times \text{CH}_2$ ), 1.62 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.30–2.37 (4 H, m,  $2 \times \text{COCH}_2$ ), 3.73 (2 H, dd,  $J$  = 5.0, 1.4 Hz, H3a,3b), 4.23 (1 H, dd,  $J$  = 11.9, 5.7 Hz, H1a), 4.32 (1 H, dd,  $J$  = 11.9, 4.5 Hz, H1b), 5.06–5.11 (1 H, m, H2).

**3. Synthesis of phosphatidylglycerols****O-Benzyl-O-(1-O-((R)-10-methyloctadecanoyl)-2-O-palmitoyl-sn-glycer-3-yl) N,N-diisopropylphosphoramidite (8c)**

Diisopropylammonium tetrazolide (34.1 mg, 0.199 mmol) was added to a solution of alcohol **7c** (58.0 mg, 0.094 mmol) and benzyl bis(*N,N*-diisopropyl)phosphoramidite<sup>37</sup> (62.3 mg, 0.189 mmol) in dry  $\text{CH}_2\text{Cl}_2$  at 0 °C under nitrogen. The reaction mixture was stirred for 30 min at 0 °C and 25 min at room temperature. The solvent was removed in vacuo and the residue was purified by column chromatography on alumina (5% EtOAc/petroleum ether) to afford the title compound as a colorless oil (50 mg, 85%);  $[\alpha]_{\text{D}}^{23} +5.4$  ( $c$  = 0.5,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.83 (3 H, d,  $J$  = 6.6 Hz,  $\text{CCH}_3$  (TBSA)), 0.88 (6 H, t,  $J$  = 6.9 Hz,  $2 \times \text{CH}_3$ ), 1.18 (2 H, m,  $\text{CH}_2$ ), 1.17–1.20 (12 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ ), 1.23–1.35 (49 H, m,  $24 \times \text{CH}_2$ ,  $1 \times \text{CH}$ ), 1.57–1.62 (4 H, m,  $\text{COCH}_2\text{CH}_2$ ), 2.21–2.31 (4 H, m,  $\text{COCH}_2$ ), 3.60–3.81 (4 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ , H3a,3b), 4.13–4.20 (1 H, m, H1a), 4.31–4.37 (1 H, m, H1b), 4.64–4.77 (2 H, m,  $\text{CH}_2\text{Ph}$ ), 5.15–5.22 (1 H, m, H2), 7.30–7.41 (5 H, m, Ph);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  14.2, 19.8, 22.8, 24.6, 24.7, 24.8, 25.0, 29.2, 29.4, 29.5, 29.6, 29.8, 29.8, 32.0, 34.3, 43.1, 43.2, 61.6, 61.7, 61.9, 62.6, 63.2, 65.4, 65.6, 70.9 (N-CH), 127.0, 127.4, 128.3, 139.4 (Ph), 173.1, 173.5 (C=O);  $^{31}\text{P}$  NMR (202 MHz,  $\text{CDCl}_3$ )  $\delta$  148.8; HRMS-ESI:  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{51}\text{H}_{95}\text{O}_6\text{NP}$  848.6897; found 848.6885.

**O-Benzyl-O-(1-O-((R)-10-methyloctadecanoyl)-2-O-palmitoyl-sn-glycer-1-yl) O-(2,3-di-O-benzyl-sn-glycer-1-yl) phosphate (10c)**

1*H*-Tetrazole (4.8 mg, 0.068 mmol) was added to a solution of phosphoramidite (42.0 mg, 0.049 mmol) and 2,3-di-O-benzyl-sn-glycerol **9**<sup>38</sup> (10.7 mg, 0.0392 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (2 ml) at 0 °C under nitrogen. The reaction mixture was stirred for 15 min at 0 °C and 2 hours at room temperature. The reaction mixture was cooled to 0 °C and *t*-BuOOH (5 M in decane) (5  $\mu\text{l}$ , 0.0582 mmol) was added and stirred for 2 h warming to room temperature. Excess oxidant was quenched with sodium metabisulfite (1M, 0.1 ml), the reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , extracted, dried ( $\text{MgSO}_4$ ), filtered and concentrated. The residue was purified by column chromatography on silica gel (35% EtOAc/ petroleum ether, 0.5% Et<sub>3</sub>N) to afford the title compound as an oil (25 mg, 55%);  $[\alpha]_{\text{D}}^{24} 6.3$  ( $c$  0.25,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ )  $\delta$  0.83 (3 H, d,  $J$  = 6.6 Hz,  $\text{CCH}_3$  (TBSA)), 0.88 (6 H, t,  $J$  = 6.9 Hz,  $2 \times \text{CH}_3$  (TBSA, palmitoyl)), 1.06–1.07 (2 H, m,  $\text{CH}_2$ ), 1.26 (49 H, m,  $24 \times \text{CH}_2$ ,  $1 \times \text{CH}$ ), 1.57 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.26 (4 H, m,  $2 \times \text{COCH}_2$ ), 3.56 (2 H, t,  $J$  = 5.6 Hz, H3'a,3'b), 3.77 (1 H, m, H2'), 4.03–4.16 (4 H, m, H3a,3b,1'a,1'b), 4.23 (2 H, m, H1a,1b), 4.51 (2 H, d,  $J$  = 4.8 Hz,  $\text{CH}_2\text{Ph}$ ), 4.60–4.67 (2 H, m,  $\text{CH}_2\text{Ph}$ ), 5.02–5.05 (2 H, m,  $\text{CH}_2\text{Ph}$ ), 5.16 (1 H, m, H2'), 7.32 (15 H, m, Ph);  $^{13}\text{C}$  NMR (126 MHz,  $\text{CDCl}_3$ )  $\delta$  14.2, 19.8, 22.8, 24.9, 27.2, 29.2, 29.2, 29.4, 29.5, 29.6, 29.7, 29.8, 30.1, 30.2, 32.0, 32.9, 34.1, 34.2, 37.2, 61.8, 65.5, 67.3, 67.3, 69.1, 69.4, 69.4, 69.5, 69.6, 72.3, 73.6, 127.7, 127.8, 127.9, 128.0, 128.5, 128.5, 128.7 (Ph), 172.9, 173.3 (C=O);  $^{31}\text{P}$  NMR (202 MHz,  $\text{CDCl}_3$ )  $\delta$  -1.01, -0.98; HRMS:  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{62}\text{H}_{99}\text{O}_{10}\text{P}$ , 1035.7009; found 1035.7006.

**Triethylammonium O-(1-O-(R)-10-methyloctadecanoyl-2-O-palmitoyl-sn-glycer-1-yl) O-(sn-glycer-1-yl) phosphate (1)**

A mixture of  $\text{Pd}(\text{OH})_2$  on carbon (20%, 5 mg, 0.035 mmol), **10c** (10 mg, 9.65  $\mu\text{mol}$ ) in EtOAc:methanol (5 ml, 5:1) was stirred under hydrogen (20 bar) for 4 h at room temperature. The reaction mixture is filtered through a short plug of Celite to remove the catalyst. Triethylamine (5  $\mu\text{l}$ ) was added to the filtrate and solvent removed under reduced pressure. The resulting residue was purified by flash chromatography (20% methanol:chloroform) to afford **1** as a colorless oil (9 mg, 98%);  $[\alpha]_{\text{D}}^{24} +5.8$  ( $c$  0.25,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR (500 MHz,  $\text{CD}_3\text{OD}:\text{CDCl}_3$  (1:2))  $\delta$  0.80 (3 H, d,  $J$  = 6.6 Hz,  $\text{CCH}_3$  (TBSA)), 0.85 (6 H, t,  $J$  = 6.8 Hz,  $2 \times \text{CH}_3$  (TBSA, palmitoyl)), 1.03–1.06 (2 H, m,  $\text{CH}_2$ ), 1.17–1.36 (62 H, m,  $26 \times \text{CH}_2$ ,  $1 \times \text{CH}$ ,  $\text{HN}(\text{CH}_2\text{CH}_3)_3$ ), 1.57 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.29 (4 H, dd,  $J$  = 15.6, 8.0 Hz,  $2 \times \text{COCH}_2$ ), 3.11 (4 H, q,  $J$  = 7.3 Hz,  $\text{NH}(\text{CH}_2\text{CH}_3)_3$ ), 3.58 (2 H, s, H3'a,3'b), 3.76 (1 H, s, H2'), 3.90 (2 H, s, H1'a,1'b), 3.98 (2 H, s, H3a,3b), 4.15 (1 H, dd,  $J$  = 11.8, 6.5 Hz, H1b), 4.38 (1 H, d,  $J$  = 12.0 Hz, H1a), 5.20 (1 H, s, H2);  $^{13}\text{C}$  NMR (126 MHz,  $\text{CD}_3\text{OD}:\text{CDCl}_3$  (1:2))  $\delta$  14.2, 19.9, 23.0, 25.2, 25.2, 27.4, 29.5, 29.5, 29.7, 29.9, 29.9, 30.0, 30.3, 30.4, 32.3, 33.1, 34.4, 34.6, 37.4, 46.8, 62.9, 64.2, 67.1, 70.7, 71.4 (glycerol C), 174.0, 174.3 (C=O);  $^{31}\text{P}$  NMR (202 MHz,  $\text{CD}_3\text{OD}:\text{CDCl}_3$  (1:2))  $\delta$  4.78; HRMS:  $[\text{M} + \text{H}]^+$  calcd for  $\text{C}_{47}\text{H}_{96}\text{NO}_{10}\text{P}$  866.6805; found 866.6806.

**O-Benzyl-O-(1-O-(palmitoyl)-2-O-stearoyl-sn-glycer-3-yl) N,N-diisopropylphosphoramidite (8a)**

Alcohol **7a** (30 mg, 0.050 mmol) was treated as for the preparation of **8c** to afford **8a** (30.4 mg, 72%).  $[\alpha]_D^{24} +5.8$  ( $c = 0.5$ , CHCl<sub>3</sub>); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (6 H, t,  $J = 6.8$  Hz,  $2 \times \text{CH}_3$ ), 1.17–1.20 (12 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ ), 1.23–1.35 (52 H, m,  $26 \times \text{CH}_2$ ), 1.57–1.62 (4 H, m, COCH<sub>2</sub>CH<sub>2</sub>), 2.21–2.31 (4 H, m, COCH<sub>2</sub>), 3.60–3.67 (2 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ ), 3.69–3.75 (1 H, m, H3b), 3.76–3.81 (1 H, m, H3a), 4.13–4.20 (1 H, m, H1a), 4.34 (1 H, td,  $J = 12.2, 3.8$  Hz, H1b), 4.63–4.79 (2 H, m, CH<sub>2</sub>Ph), 5.18–5.19 (1 H, m, H2), 7.27–7.35 (5 H, m, Ph); <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 22.8, 24.6, 24.7, 24.8, 25.0, 29.2, 29.4, 29.5, 29.6, 29.8, 29.8, 32.0, 34.3, 43.1, 43.2, 61.6, 61.7, 61.9, 62.6, 63.2, 65.4, 65.6, 70.9 (N-CH), 127.0, 127.4, 128.3, 139.4 (Ph), 173.1, 173.5 (C=O); <sup>31</sup>P NMR (202 MHz, CDCl<sub>3</sub>)  $\delta$  148.8; HRMS-ESI:  $[\text{M} + \text{H}]^+$  calcd for C<sub>50</sub>H<sub>92</sub>NO<sub>6</sub>P 834.6696; found 834.6696.

**O-Benzyl-O-(1-O-(stearoyl)-2-O-palmitoyl-sn-glycer-3-yl) N,N-diisopropylphosphoramidite (8b)**

Alcohol **8a** (36.0 mg, 0.60 mmol) was treated as for the preparation of **8c** to afford **8b** (29.8 mg, 58%).  $[\alpha]_D^{24} +5.5$  ( $c = 0.5$ , CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (6 H, t,  $J = 6.8$  Hz,  $2 \times \text{CH}_3$ ), 1.17–1.20 (12 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ ), 1.23–1.35 (52 H, m,  $26 \times \text{CH}_2$ ), 1.57–1.62 (4 H, m, COCH<sub>2</sub>CH<sub>2</sub>), 2.21–2.31 (4 H, m, COCH<sub>2</sub>), 3.58–3.83 (4 H, m,  $2 \times \text{CH}(\text{CH}_3)_2$ , H3a,3b), 4.12–4.19 (1 H, m, H1a), 4.30–4.38 (1 H, m, H1b), 4.62–4.80 (2 H, m, CH<sub>2</sub>Ph), 5.17–5.24 (1 H, m, H2), 7.27–7.35 (5 H, m, Ph); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 22.8, 24.6, 24.7, 24.8, 25.0, 29.2, 29.4, 29.5, 29.6, 29.8, 29.8, 32.0, 34.3, 43.1, 43.2, 61.6, 61.7, 61.9, 62.6, 63.2, 65.4, 65.6, 70.9 (N-CH), 127.0, 127.4, 128.3, 139.4 (Ph), 173.1, 173.5 (C=O); <sup>31</sup>P NMR (202 MHz, CDCl<sub>3</sub>)  $\delta$  148.0; HRMS-ESI:  $[\text{M} + \text{H}]^+$  calcd for C<sub>50</sub>H<sub>92</sub>NO<sub>6</sub>P 834.6696; found 834.6696.

**O-Benzyl-O-(1-O-(palmitoyl)-2-O-stearoyl-sn-glycer-3-yl)-O-(2,3-di-O-benzyl-sn-glycer-1-yl) phosphate (10a)**

1H-Tetrazole (2.7 mg, 0.039 mmol) was added to a solution of phosphoramidite (24.0 mg, 0.028 mmol) and 2,3-di-O-benzyl-sn-glycerol **9**<sup>38</sup> (9.4 mg, 0.034 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (2 ml) at 0 °C under nitrogen. The reaction mixture was stirred for 15 min at 0 °C and 2 h at room temperature. The reaction mixture was cooled to 0 °C and t-BuOOH (5 M in decane) (4.1  $\mu$ l, 0.042 mmol) was added to the reaction mixture and stirred for 2 h with warming to room temperature. Excess oxidant was quenched with sodium metabisulfite (1 M, 0.1 ml), the reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, extracted, dried (MgSO<sub>4</sub>), filtered and concentrated. The solvent was removed in vacuo and the residue was purified by column chromatography on silica gel (35% EtOAc/ petroleum ether, 0.5% Et<sub>3</sub>N) which resulted in a 1:1 mixture the product and unreacted dibenzyl glycerol, which was directly subjected to hydrogenolysis.

**O-Benzyl-O-(1-O-(stearoyl)-2-O-palmitoyl-sn-glycer-3-yl)-O-(2,3-di-O-benzyl-sn-glycer-1-yl) phosphate (10b)**

Synthesized as for **10c**, 10.2 mg (34%).  $[\alpha]_D^{25} +6.2$  ( $c = 0.25$ , CHCl<sub>3</sub>); <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (6 H, t,  $J = 6.8$  Hz,  $2 \times \text{CH}_3$ ), 1.27 (56 H, m,  $28 \times \text{CH}_2$ ), 1.57 (4 H, m,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.26 (4 H, t,  $J = 7.6$  Hz,  $2 \times \text{COCH}_2$ ), 3.56 (2 H, t,  $J = 5.6$  Hz, H3'a,3'b), 3.75–3.79 (1 H, m, H2'), 4.03–4.29 (6 H, m, H1'a,1'b,3a,3b,1a,1b), 4.50–4.54 (2 H, m, CH<sub>2</sub>Ph), 4.60–4.68 (2 H, m, CH<sub>2</sub>Ph), 5.04 (2 H, m, CH<sub>2</sub>Ph), 5.13–5.20 (1 H, m, H2), 7.27–7.35 (15 H, m, Ph); <sup>13</sup>C NMR (126 MHz, CDCl<sub>3</sub>)  $\delta$  14.2, 22.8, 24.9, 27.2, 29.2, 29.2, 29.4, 29.5, 29.6, 29.7, 29.8, 30.1, 30.2, 32.0, 32.9, 34.1, 34.2, 37.2, 61.8, 65.5, 67.3, 67.3, 69.1, 69.4, 69.4, 69.5, 69.6, 72.3, 73.6, 127.7, 127.8, 127.9, 128.0, 128.5, 128.5, 128.7 (Ph), 172.9, 173.3 (C=O); <sup>31</sup>P NMR (202 MHz, CDCl<sub>3</sub>)  $\delta$  -1.01, -0.98; HRMS:  $[\text{M} + \text{H}]^+$  calcd for C<sub>61</sub>H<sub>97</sub>O<sub>10</sub>P 1021.6853; found 1021.6852.

**Triethylammonium O-(1-O-palmitoyl-2-O-stearoyl-sn-glycer-1-yl) phosphate (11a)**

Synthesized as for **1**, 6.2 mg (27% over 2 steps).  $[\alpha]_D^{25} +5.6$  ( $c = 0.25$ , CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD:CDCl<sub>3</sub> (1:2))  $\delta$  0.84 (6 H, t,  $J = 6.4$  Hz,  $2 \times \text{CH}_3$ ), 1.10–1.14 (9 H, m, HN(CH<sub>2</sub>CH<sub>3</sub>)<sub>3</sub>), 1.17–1.36 (56 H, m,  $28 \times \text{CH}_2$ ), 1.52 (4 H, s,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.23 (4 H, dd,  $J = 14.5, 7.2$  Hz,  $2 \times \text{COCH}_2$ ), 3.46–3.61 (6 H, s, H3'a,3'b, HN(CH<sub>2</sub>CH<sub>3</sub>)<sub>3</sub>), 3.65–3.73 (1 H, m, H2'), 3.83–3.90 (4 H, m, H1'a,1'b,3a,3b), 3.95 (2 H, m, H3a,3b), 4.11 (1 H, dd,  $J = 12.0, 6.6$  Hz, H1b), 5.16 (1 H, s, H2); <sup>13</sup>C NMR (126 MHz, CD<sub>3</sub>OD:CDCl<sub>3</sub> (1:2))  $\delta$  14.2, 23.0, 25.2, 25.2, 29.5, 29.5, 29.7, 29.7, 29.9, 30.0, 30.0, 32.3, 34.4, 34.6, 46.5, 62.8, 63.0, 63.7, 66.8, 70.8, 70.9, 71.5, 72.7, 174.0, 174.4; <sup>31</sup>P NMR (202 MHz, CDCl<sub>3</sub>)  $\delta$  4.78; HRMS:  $[\text{M} + \text{H}]^+$  calcd for C<sub>40</sub>H<sub>79</sub>O<sub>10</sub>P 751.5444; found 751.5444.

**O-(1-O-Stearoyl-2-O-palmitoyl-sn-glycer-1-yl)-O-(sn-glycer-1-yl) phosphate (11b)**

Synthesized as for **1**, 9.1 mg (90%).  $[\alpha]_D^{25} +4.8$  ( $c = 0.25$ , CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD:CDCl<sub>3</sub> (1:2))  $\delta$  0.84 (6 H, t,  $J = 6.4$  Hz,  $2 \times \text{CH}_3$ ), 1.17–1.36 (56 H, m,  $28 \times \text{CH}_2$ ), 1.57 (4 H, s,  $2 \times \text{COCH}_2\text{CH}_2$ ), 2.29 (4 H, dd,  $J = 14.5, 7.2$  Hz,  $2 \times \text{COCH}_2$ ), 3.57 (2 H, s, H3'a,3'b), 3.70–3.78 (1 H, m, H2'), 3.84–3.90 (2 H, s, H1'a, H1'b), 3.95 (2 H, m, H3a,3b), 4.15 (1 H, dd,  $J = 11.9, 6.7$  Hz, H1b), 4.37 (1 H, m, H1a), 5.20 (1 H, s, H2); <sup>13</sup>C NMR (126 MHz, CD<sub>3</sub>OD:CDCl<sub>3</sub> (1:2))  $\delta$  14.2, 23.0, 25.2, 25.2, 29.5, 29.5, 29.7, 29.7, 29.9, 30.0, 30.0, 32.3, 34.4, 34.6, 46.5, 62.8, 63.0, 63.7, 66.8, 70.8, 70.9, 71.5, 72.7, 174.0, 174.4; <sup>31</sup>P NMR (202 MHz, CDCl<sub>3</sub>)  $\delta$  5.25; HRMS:  $[\text{M} + \text{H}]^+$  calcd for C<sub>40</sub>H<sub>79</sub>O<sub>10</sub>P 751.5444; found 751.5444.

**Collision-induced dissociation mass spectrometry of phosphatidylglycerols**

Characterization of lipid ions, introduced to a Q Exactive Plus Orbitrap mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) via nanoESI (nESI) using an Advion Triversa Nanomate (Advion, Ithaca, NY), was performed in negative ionization mode by higher-energy collision induced dissociation (HCD)-tandem mass spectrometry (MS/MS). Deprotonated precursor ions were mono-isotopically isolated using a window of  $\pm 0.5$   $m/z$ . Product ion spectra were acquired



## ARTICLE

## Organic Biomolecular Chemistry

using a mass resolving power of 70,000. Spectra shown are the average of 100 scans.

## Acknowledgements

We acknowledge the Australian Research Council for financial support (DP160100597). Prof Gavin Reid and Mengxuan Fang are thanked for assistance with the mass spectrometric analysis of PGs.

## References

1. S. Furse, *J. Chem. Biol.*, 2017, **10**, 1-9.
2. S. E. Horvath and G. Daum, *Prog. Lipid Res.*, 2013, **52**, 590-614.
3. I. Van Rhijn, T. van Berlo, T. Hilmenyuk, T. Y. Cheng, B. J. Wolf, R. V. Tatituri, A. P. Uldrich, G. Napolitani, V. Cerundolo, J. D. Altman, P. Willemsen, S. Huang, J. Rossjohn, G. S. Besra, M. B. Brenner, D. I. Godfrey and D. B. Moody, *Proc. Natl. Acad. Sci. USA*, 2016, **113**, 380-385.
4. R. V. Tatituri, G. F. Watts, V. Bhowruth, N. Barton, A. Rothchild, F. F. Hsu, C. F. Almeida, L. R. Cox, L. Eggeling, S. Cardell, J. Rossjohn, D. I. Godfrey, S. M. Behar, G. S. Besra, M. B. Brenner and M. Brigl, *Proc. Natl. Acad. Sci. USA*, 2013, **110**, 1827-1832.
5. B. J. Wolf, R. V. Tatituri, C. F. Almeida, J. Le Nours, V. Bhowruth, D. Johnson, A. P. Uldrich, F. F. Hsu, M. Brigl, G. S. Besra, J. Rossjohn, D. I. Godfrey and M. B. Brenner, *J. Immunol.*, 2015, **195**, 2540-2551.
6. E. Maloney, D. Stankowska, J. Zhang, M. Fol, Q. J. Cheng, S. Lun, W. R. Bishai, M. Rajagopalan, D. Chatterjee and M. V. Madiraju, *PLoS Pathog.*, 2009, **5**, e1000534.
7. M. Jackson, D. C. Crick and P. J. Brennan, *J. Biol. Chem.*, 2000, **275**, 30092-30099.
8. P. V. Sarma, L. Srikanth, K. Venkatesh, P. S. Murthy and P. U. Sarma, *Bioinformation*, 2013, **9**, 690-695.
9. H. Okuyama, T. Kankura and S. Nojima, *J. Biochem.*, 1967, **61**, 732-737.
10. B. ter Horst, C. Seshadri, L. Sweet, D. C. Young, B. L. Feringa, D. B. Moody and A. J. Minnaard, *J. Lipid Res.*, 2010, **51**, 1017-1022.
11. F. F. Hsu, J. Turk, R. M. Owens, E. R. Rhoades and D. G. Russell, *J. Am. Soc. Mass Spectrom.*, 2007, **18**, 479-492.
12. J. Rossjohn, S. Gras, J. J. Miles, S. J. Turner, D. I. Godfrey and J. McCluskey, *Annu. Rev. Immunol.*, 2015, **33**, 169-200.
13. M. V. Dhodapkar and V. Kumar, *J. Immunol.*, 2017, **198**, 1015-1021.
14. F. S. Prout, J. Cason and A. W. Ingersoll, *J. Am. Chem. Soc.*, 1948, **70**, 298-305.
15. S. Mekala and R. C. Hahn, *J. Org. Chem.*, 2015, **80**, 1610-1617.
16. R. P. Linstead, J. C. Lunt and B. C. L. Weedon, *J. Chem. Soc.*, 1951, 1130-1132.
17. B. S. Dyer, J. D. Jones, G. D. Ainge, M. Denis, D. S. Larsen and G. F. Painter, *J. Org. Chem.*, 2007, **72**, 3282-3288.
18. P. Fodran and A. J. Minnaard, *Org. Biomol. Chem.*, 2013, **11**, 6919-6928.
19. X. Liu, B. L. Stocker and P. H. Seeberger, *J. Am. Chem. Soc.*, 2006, **128**, 3638-3648.
20. P. S. Patil, T.-J. R. Cheng, M. M. L. Zulueta, S.-T. Yang, L. S. Lico and S.-C. Hung, *Nat. Commun.*, 2015, **6**, 7239.
21. B. Cao, X. Chen, Y. Yamaro-Botte, M. B. Richardson, K. L. Martin, G. N. Khairallah, T. W. Rupasinghe, R. M. O'Flaherty, R. A. O'Hair, J. E. Ralton, P. K. Crellin, R. L. Coppel, M. J. McConville and S. J. Williams, *J. Org. Chem.*, 2013, **78**, 2175-2190.
22. I. O. Roberts and M. S. Baird, *Chem. Phys. Lipids*, 2006, **142**, 111-117.
23. P. D'Arrigo, L. de Ferra, G. Pedrocchi-Fantoni, D. Scarcelli, S. Servi and A. Strini, *J. Chem. Soc. Perkin Trans. 1*, 1996, 2657-2660; R. Sato, Y. Itabashi, H. Fujishima, H. Okuyama and A. Kuksis, *Lipids*, 2004, **39**, 1025-1030.
24. P. Woolley and H. Eibl, *Chem. Phys. Lipids*, 1988, **47**, 55-62; R. Wohlgemuth, N. Waespe-Sarcevic and J. Seelig, *Biochemistry*, 1980, **19**, 3315-3321; E. Baer and D. Buchnea, *J. Biol. Chem.*, 1958, **232**, 895-902; R. M. Saunders and H. P. Schwarz, *J. Am. Chem. Soc.*, 1966, **88**, 3844-3847.
25. M. H. Caruthers, A. D. Barone, S. L. Beaucage, D. R. Dodds, E. F. Fisher, L. J. McBride, M. Matteucci, Z. Stabinsky and J.-Y. Tang, *Methods Enzymol.*, 1987, **154**, 287-313.
26. G. D. Prestwich, *Acc. Chem. Res.*, 1996, **29**, 503-513.
27. S. M. Ali, M. U. Ahmad, P. Koslosky, K. Kasireddy, U. Murali Krishna and I. Ahmad, *Tetrahedron*, 2006, **62**, 6990-6997.
28. E. N. Jacobsen, F. Kakiuchi, R. G. Konsler, J. F. Larow and M. Tokunaga, *Tetrahedron Lett.*, 1997, **38**, 773-776.
29. C. E. Dreef, C. J. J. Elie, P. Hoogerhout, G. A. van der Marel and J. H. van Boom, *Tetrahedron Letters*, 1988, **29**, 6513-6515.
30. N. J. Jensen, K. B. Tomer and M. L. Gross, *Lipids*, 1987, **22**, 480-489.
31. P. B. Smith, A. P. Snyder and C. S. Harden, *Anal. Chem.*, 1995, **67**, 1824-1830.
32. F.-F. Hsu and J. Turk, *J. Am. Soc. Mass Spectrom.*, 2000, **11**, 986-999.
33. R. V. Tatituri, B. J. Wolf, M. B. Brenner, J. Turk and F. F. Hsu, *Anal. Bioanal. Chem.*, 2015, **407**, 2519-2528.
34. W. C. Still, M. Kahn and A. M. Mitra, *J. Org. Chem.*, 1978, **43**, 2923-2925.
35. A. B. Pangborn, M. A. Giardello, R. H. Grubbs, R. K. Rosen and F. J. Timmers, *Organometallics*, 1996, **15**, 1518-1520.
36. S. Front, N. Court, M.-L. Bourigault, S. Rose, B. Ryffel, F. Erard, V. F. J. Quesniaux and O. R. Martin, *ChemMedChem*, 2011, **6**, 2081-2093.
37. G. M. Rankin, B. J. Compton, K. A. Johnston, C. M. Hayman, G. F. Painter and D. S. Larsen, *J. Org. Chem.*, 2012, **77**, 6743-6759.
38. N. Chen and J. Xie, *J. Org. Chem.*, 2014, **79**, 10716-10721; C. A. A. van Boeckel, G. M. Visser and J. H. van Boom, *Tetrahedron*, 1985, **41**, 4557-4565.