Organic Cite This: Org. Lett. XXXX, XXX, XXX–XXX

Loïc R. E. Pantaine,[†] John A. Milligan,[†] Jennifer K. Matsui,[†][©] Christopher B. Kelly,^{*,‡,§}[©] and Gary A. Molander*^{,†}

[†]Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, 231 South 34th Street, Philadelphia, Pennsylvania 19104-6323, United States

[‡]Department of Chemistry, Virginia Commonwealth University, 1001 West Main Street, P.O. Box 842006, Richmond, Virginia 23284-9069, United States

[§]Medicines for All Institute, Virginia Commonwealth University, Biotech 8 737 North Fifth Street, Richmond, Virginia 23219-1441, United States

S Supporting Information

ABSTRACT: Photoredox-mediated radical/polar crossover (RPC) processes offer new avenues for the synthesis of cyclic molecules. This process has been realized for the construction of medium-sized saturated nitrogen heterocycles. Photocatalytically generated alkyl radicals possessing pendant leaving groups engage imines in C-C bond formation, and subsequent reduction of the intermediate nitrogen-centered radical triggers anionic ring closure. With the aid of visible light irradiation, substituted pyrrolidines, piperidines, and azepanes can be prepared under mild, redox-neutral conditions.

eterocyclic compounds comprise 87% of known small molecule therapeutics.¹ Nitrogen-based heterocycles in particular are among the most common subunit in drug molecules.² These structures also serve as the backbone of ligands and appear in numerous natural and designed molecules.^{1,2} In the context of drug development, the improved pharmacological properties of compounds with higher fractions of sp³-hybridized centers³ has sparked an increasing interest in incorporating saturated nitrogen heterocycles in bioactive molecules.⁴ Piperidines are the most prominent saturated nitrogen heterocycles in therapeutic compounds, followed closely by piperazines and pyrrolidines.² The popularity and utility of these heterocycles motivate the pursuit of novel synthetic methods to prepare these cyclic systems in a mild and modular fashion.

The most common approaches to introduce saturated nitrogen heterocycles are C-N bond-forming processes through S_NAr chemistry, cross-coupling strategies such as Buchwald-Hartwig amination, or peptide coupling.^{1,5} However, many medicinally relevant compounds possess α substitution (often aryl, Figure 1), which poses challenges to these methods (Figure 2). A classical method for the synthesis of α -arylated nitrogen heterocycles is a lithiation followed by treatment with an electrophile or conversion into a suitable nucleophile (i.e., cuprate or zincate) for cross-coupling. Another class of increasingly popular reactions are singleelectron variants of the lithiation approach, initiated through either hydrogen-atom transfer (HAT)⁷ or photoredox-induced

Mild Conditions Redox-Neutral 35 Examples Up to 84% yield



Figure 1. Bioactive nitrogen-containing heterocycles.

oxidation of radical precursors (such as a carboxylic acids or trifluoroborates).⁸ Although powerful, these strategies are contingent on having a preconstructed saturated N-heterocycle that is subsequently functionalized with the group of interest.

A less common approach to α -arylated nitrogen heterocycles is the construction of the ring from acyclic precursors through an annulation process. The most obvious way to accomplish such a process, reductive amination, is notoriously challenging,

Received: February 17, 2019

Letter

pubs.acs.org/OrgLett



considering that the cyclic imine intermediate is prone to polymerization/dimerization.⁹ Recently, a resurgence in methods based on the Hofmann–Löffler–Freytag (HLF) reaction have yielded elegant approaches to pyrrolidines from amines with distal olefins.¹⁰ Likewise, PCET has allowed lactams to be prepared from cyclization of an amidyl radical containing a pendant olefin.¹¹

Recently, Bode and co-workers have developed a coppermediated annulation process using "SnAP" reagents.¹² This strategy involves condensation of an organotin-tethered amine with an aldehyde to generate an aldimine. Subsequent oxidative generation of a heteroatom-stabilized primary carbon-centered radical via destannylative fragmentation triggers *6-endo-trig* cyclization. Reduction of the aminyl radical provides the desired nitrogen heterocycle (e.g., thiomorpholine, oxazepane, 3alkoxypyrrolidine). A silicon variant of the annulation paradigm has also been developed using "SLAP" reagents and transitioned to continuous flow.¹³

The identification of various precursors that furnish alkyl radicals under photoredox conditions, such as Bode's SLAP reagents, has led to the discovery of previously nonviable reactions and allowed new synthetic disconnections to be realized.¹⁴ Visible-light photoredox catalysis provides a means to tap the synthetic potential of radicals in a way not possible when using stoichiometric radical generation techniques, while still retaining the benefits associated with odd-electron reactivity. These hallmarks of photoredox processes have precipitated their rapid uptake in both academia and industry, especially for late-stage functionalization.¹⁵ Visible-light-mediated approaches have revolutionized an array of different reaction classes, including cross-coupling,¹⁶ cycloaddition,¹⁷ and, recently, radical/polar crossover (RPC)¹⁸ processes.

In the context of the latter, our group has initiated a program to apply various radical precursors in RPC processes that lead to annulation reactions, including the cyclopropanation of alkenes (Figure 3).¹⁹ Inspired by the successes of these annulation processes and the SLAP/SnAP paradigm,^{13,14} a novel approach to the synthesis of saturated nitrogen-based heterocycles from acyclic species was considered. The crux of this approach is the



Lette

Figure 3. Comparison of related developments to this work.

radical alkylation of an imine using an oxidizable, bifunctional radical precursor. Subsequent reduction of the resulting nitrogen-centered radical would furnish an amide anion that could engage a distal leaving group in an intramolecular displacement. A series of annulation processes (5-, 6-, or even 7-*exo-tet* ring closures) can be envisioned using this improved radical/polar crossover approach. Indeed, typical methods to accomplish this type of process require a combination of stoichiometric $Mn_2(CO)_{10}$, $InCl_3$, and UV irradiation.²⁰

The proposed annulation makes use of a small set of easily accessible bis(catecholato)silicate radical precursors.²¹ These reagents have previously been effective for the radical alkylation of imines²² and the cyclopropanation of olefins^{19a} (Figure 3). Motivated by these previous successes, a preliminary screen was conducted wherein imine 1 was irradiated with a series of 3-halopropylsilicates (Cl, Br, I) in the presence of a photocatalyst [Ru(bpy)₃(PF₆)₂] in DMF (Table 1, entries 1–3). Although



	1				
cı (Ph X H photocatal solvent (0		► _{CI}	Ph.N 2
entry ^a	Х	photocat. ^b	solvent	time (h)	yield $(\%)^c$
1	Cl	[Ru]	DMF	16	8
2	Br	[Ru]	DMF	16	20
3	Ι	[Ru]	DMF	16	10
4 ^{<i>d</i>}	Br	[Ru]	DMF	48	46
5 ^e	Br	[Ru]	DMSO	16	46
6 ^e	Br	[Ir]	DMSO	16	54
7 ^e	Br	4CzIPN	DMSO	16	35
8 ^f	Br	[Ir]	DMSO	16	60
9 ^f	Br	[Ir]	DMSO	0.5	59
$10^{f,g}$	Br	[Ir]	DMSO	16	-
11 ^f	Br	-	DMSO	16	17 ^h

^{*a*}Conditions unless otherwise noted: silicate (5 equiv, 1.5 mmol), imine (1.0 equiv, 0.3 mmol), photocatalyst (2 mol %), DMSO (0.1 M), irradiating with 21 W CFL. ^{*b*}Photocatalyst: [Ru] = [Ru(bpy)₃]-(PF₆)₂ and [Ir] = [Ir{dF(CF₃)₂ppy}₂(bpy)]PF₆. ^{*c*}Isolated yield after purification. ^{*d*}2 equiv of silicate used. ^{*e*}A 1:2 silicate/imine ratio used along with 4 W blue LEDs. ^{*f*}A 1:2 silicate/imine ratio used along with 30 W blue LEDs. ^{*g*}Reaction conducted in the dark. ^{*h*}NMR yield of crude reaction mixture. Scheme 1. Scope of the Radical Alkylation/Cyclization Processes with Imines^{*a,b*}



^aReaction conditions unless otherwise noted: alkylsilicate (1 equiv, 0.3 mmol), imine (2.0 equiv, 0.6 mmol), photocatalyst (2 mol %), DMSO (0.1 M), irradiating with 30 W blue LEDs at 35 °C for 30 min; all yields are isolated yields after purification. ^bReaction run for 16 h. ^cReaction conducted on 1 mmol scale with 1 mol % photocatalyst. ^dReaction conducted on 2 mmol scale with 1 mol % of photocatalyst.

annulation was observed in all cases, the reaction generally proceeded slowly and gave poor conversion to the desired pyrrolidine 2. Interestingly, of this series, the 3-bromopropylsilicate proved best. Speculatively, the 3-bromopropyl radical provides a good compromise of stability (3-iodopropyl radical is prone to $S_{\rm H}2$ 3-exo-tet cyclization²³) and reactivity (some addition without cyclization was observed when using 3chloropropylsilicate). Reducing the loading of the 3-bromopropylsilicate and extending the reaction time led to more than doubling the yield of 2 (entry 4). High throughput experimentation (HTE) techniques revealed that using a 2fold excess of the imine reagent provided a higher yield (see Supporting Information). However, excess imine could be recovered upon chromatographic purification in cases where the imine is valuable (such as late-stage derivatization scenarios). Scale up of these conditions gave the identical yield in a much shorter period (entry 5). Evaluation of alternative photocatalysts revealed that $Ir[dF(CF_3)ppy]_2(bpy)PF_6$ was superior (entries 6–7). Using this photocatalyst and a more powerful light source, the reaction time was reduced to a mere 30 min (entry 9). Various additives were screened, but gave either similar or

diminished yields (see Supporting Information for details). Control studies confirmed irradiation was necessary for reactivity (entry 10), but the reaction could proceed (albeit in poor yield) in the absence of a photocatalyst (entry 11). Given that alkylsilicates are known to fragment when irradiated with UV-A/UV-B light, an electron donor–acceptor complex²⁴ between the highly conjugated imine and the bis(catecholato) moiety may form, facilitating light-mediated radical generation.

With suitable conditions established, the scope of the annulation process was examined (Scheme 1). In general, when using 3-bromopropylsilicate, a variety of N-phenyl aldimines were amenable to the RPC cyclization process to provide the corresponding pyrrolidines. In some cases, extending the reaction time to 16 h was necessary to give improved yields. The reaction tolerated imines derived from both electron-rich and electron-poor benzaldehydes, including a free phenol (5). The cyclization of benzothiophene, isoxazole, and pyridine-containing imines successfully provided pyrrolidines 10-12. Substituents with a variety of steric and electronic characteristics were tolerated on either of the aryl rings on the imine substrate. In particular, replacement of the N-phenyl

group with a *p*-methoxyphenyl (PMP) group gave a comparable yield of pyrrolidine product (2 vs 19). Given the ability to oxidatively cleave PMP groups, these species would be suitable for further derivatization. Additionally, other cleavable groups were compatible with the RPC annulation process. Both ptoluenesulfonyl and N,N-dimethylsulfamidyl aldimines underwent the cyclization process to provide 22 and 23, respectively, but in diminished yields. Unfortunately, the process could not be extended to unprotected (NH) imines, N-Boc-imines, or hydrazones. Using a 4-bromobutylsilicate and identical reaction conditions, piperidines were accessible from aldimines in comparable yields (24-27). Attempts using either a 5bromopentylsilicate or 5-iodopentylsilicate reagent did provide azepane 28, although in low yield. The major product observed was the uncyclized alkyl halide. In these cases, the rate of protonation by the acidic ammonium counterion of the alkylsilicate likely exceeds the rate of 7-exo-tet cyclization. Smaller rings could not be obtained through this method; only traces of the corresponding aziridine were observed. The inability to access the requisite 2-bromoethylsilicate (likely because of facile β -silyl elimination) prevented any attempts at preparing azetidines.

Ketimines were also amenable to the reaction, allowing pyrrolidines with α -quaternary centers to be assembled rapidly. In general, ketimines derived from benzophenones fared best. Various aryl ring substituents had minimal effect on yield (29-32), an observation consistent with that seen in aldimines. Mixed aryl-alkyl ketimines could be used, albeit with diminished yields. Subjecting a ketimine derived from an α -CF₃ ketone enabled synthesis of α -CF₃-substituted pyrrolidine 33. Similarly, an acetophenone-derived ketimine was competent in the annulation process (34). Ketimines also provide opportunities for bicyclic ring construction, as demonstrated by the preparation of benzopyrrolizidine 35. This bicyclic structure bears resemblance to the core of recently reported anticancer candidates.²⁵ Piperidines bearing α -quaternary centers are also accessible using this RPC approach. Aryl ketiminoacetates and ketimidates, however, were not amenable to this process. The formation of piperidines from ketimines using the analogous 4bromobutylsilicate reagent was not as effective as piperidine formation, and in some cases unexpected side products were observed.

Based on the prior art^{18–21} and our own observations during the course of this study, the following sequence of events constitute a plausible mechanistic path for the annulation process: (1) Visible-light-mediated photoexcitation of I to its excited state II; (2) Reductive quenching of II ($E_{1/2}$ [P*/P]: 1.35 V vs SCE) by the bifunctional alkylsilicate ($E_{1/2}$ = +0.4–0.7 vs SCE for most alkylsilicates); (3) Homolytic fragmentation of oxidized IV to furnish alkyl radical V, which readily adds to imine VI, giving the *N*-centered radical VII; (4) SET reduction of VII by the reduced state of III to provide amide anion VIII; (5) C–N bond formation and ring closure by way of anionic cyclization, giving IX (Figure 4).

The radical/polar annulation process described herein enables the rapid construction of saturated nitrogen heterocycles from acyclic precursors. This RPC reaction accommodates numerous functional groups, occurs under mild conditions, and can be completed in as little as 30 min. Furthermore, its modular nature allows various medium-sized rings to be assembled from a single imine by adjusting the alkylsilicate used, and it also permits quaternary centers to be established α to nitrogen with ease. The inherent flexibility and



Figure 4. Plausible mechanism for the annulation process.

benign conditions of this strategy make it ideal for the late-stage introduction of saturated *N*-containing heterocycles.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.9b00602.

Experimental procedures, details of high-throughput screening, and ¹H, ¹³C, and ¹⁹F NMR spectra for all compounds (PDF)

AUTHOR INFORMATION

Corresponding Authors

*E-mail: gmolandr@sas.upenn.edu.

*E-mail: cbkelly@vcu.edu.

ORCID ®

Jennifer K. Matsui: 0000-0003-0693-0404 Christopher B. Kelly: 0000-0002-5530-8606 Gary A. Molander: 0000-0002-9114-5584

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors are grateful for the financial support provided by NIGMS (R01 GM 113878 to G.M.). C.B.K. is grateful for an NIH NRSA fellowship (F32 GM117634) and acknowledges start-up funds from Virginia Commonwealth University. J.K.M. is grateful for a BMS graduate Fellowship in Synthetic Organic Chemistry. We thank Dr. Charles W. Ross, III (University of Pennsylvania) for obtaining HRMS data. We acknowledge Johnson Matthey for fine chemicals donations and Kessil Lighting for donation of a prototype PR160 Rig.

REFERENCES

(1) (a) Zhang, T. Y. The Evolving Landscape of Heterocycles in Drugs and Drug Candidates. Adv. Heterocycl. Chem. 2017, 121, 1–12.
(b) Taylor, R. D.; MacCoss, M.; Lawson, A. D. G. Rings in Drugs. J. Med. Chem. 2014, 57, 5845–5859. (c) Baumann, M.; Baxendale, I. R. An overview of the synthetic routes to the best-selling drugs containing 6-membered heterocycles. Beilstein J. Org. Chem. 2013, 9, 2265–2319.
(d) Baumann, M.; Baxendale, I. R.; Ley, S. V.; Nikbin, N. An overview of the key routes to the best-selling 5-membered ring heterocyclic pharmaceuticals. Beilstein J. Org. Chem. 2011, 7, 442–495.

(2) Vitaku, E.; Smith, D. T.; Njardarson, J. T. Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals. *J. Med. Chem.* **2014**, *57*, 10257–10274.

(3) (a) Lovering, F.; Bikker, J.; Humblet, C. Escape from Flatland: Increasing Saturation as an Approach to Improving Clinical Success. *J. Med. Chem.* **2009**, *52*, 6752–6756. (b) Dandapani, S.; Marcaurelle, L. A. Accessing New Chemical Space for 'Undruggable' Targets. *Nat. Chem. Biol.* **2010**, *6*, 861–863. (c) López-Vallejo, F.; Giulianotti, M. A.; Houghten, R. A.; Medina-Franco, J. L. Expanding the Medicinally Relevant Chemical Space with Compound Libraries. *Drug Discovery Today* **2012**, *17*, 718–726.

(4) Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. Organic synthesis provides opportunities to transform drug discovery. *Nat. Chem.* **2018**, *10*, 383–394.

(5) Schneider, N.; Lowe, D. M.; Sayle, R. A.; Tarselli, M. A.; Landrum, G. A. Big Data from Pharmaceutical Patents: A Computational Analysis of Medicinal Chemists' Bread and Butter. *J. Med. Chem.* **2016**, *59*, 4385–4402.

(6) Vo, C.-V. T.; Bode, J. W. Synthesis of Saturated N-Heterocycles. *J. Org. Chem.* **2014**, *79*, 2809–2815.

 $(\overline{7})$ (a) Capaldo, L.; Ravelli, D. Hydrogen Atom Transfer (HAT): A Versatile Strategy for Substrate Activation in Photocatalyzed Organic Synthesis. *Eur. J. Org. Chem.* **2017**, 2017, 2056–2071. (b) Ahneman, D. T.; Doyle, A. G. C-H functionalization of amines with aryl halides by nickel-photoredox catalysis. *Chem. Sci.* **2016**, 7, 7002–7006. (c) Heitz, D. R.; Tellis, J. C.; Molander, G. A. Photochemical Nickel-Catalyzed C-H Arylation: Synthetic Scope and Mechanistic Investigations. *J. Am. Chem. Soc.* **2016**, 138, 12715–12718. (d) Vega, J. A.; Alonso, J. M.; Mendez, G.; Ciordia, M.; Delgado, F.; Trabanco, A. A. Continuous Flow α -Arylation of *N,N*-Dialkylhydrazones under Visible-Light Photoredox Catalysis. *Org. Lett.* **2017**, *19*, 938–941.

(8) Zuo, Z.; Ahneman, D. T.; Chu, L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. Merging photoredox with nickel catalysis: Coupling α -carboxyl sp³ carbons with aryl halides. *Science* **2014**, *345*, 437–440.

(9) Iwanejko, J.; Wojaczyńska, E. Cyclic imines – preparation and application in synthesis. *Org. Biomol. Chem.* **2018**, *16*, 7296–7314.

(10) (a) Musacchio, A. J.; Nguyen, L. Q.; Beard, H.; Knowles, R. R. Catalytic Olefin Hydroamination with Aminium Radical Cations: A Photoredox Method for Direct C-N Bond Formation. J. Am. Chem. Soc. **2014**, 136, 12217–12220. (b) Wappes, A.; Fosu, S. C.; Chopko, T. C.; Nagib, D. A. Triiodide-Mediated δ -Amination of Secondary C–H Bonds. Angew. Chem., Int. Ed. **2016**, 55, 9974–9978. (c) Martinez, C.; Muniz, K. An Iodine-Catalyzed Hofmann-Loffler Reaction. Angew. Chem., Int. Ed. **2015**, 54, 8287–8291.

(11) (a) Miller, D. C.; Choi, G. J.; Orbe, H. S.; Knowles, R. R. Catalytic Olefin Hydroamidation Enabled by Proton-Coupled Electron Transfer. *J. Am. Chem. Soc.* **2015**, *137*, 13492–13495. (b) Zheng, S.; Gutiérrez-Bonet, A.; Molander, G. A. Merging Photoredox PCET with Ni-Catalyzed Cross-Coupling: Cascade Amidoarylation of Unactivated Olefins. *Chem.* **2019**, *5*, 339–352.

(12) (a) Luescher, M. U.; Vo, C.-V. T.; Bode, J. W. SnAP Reagents for the Synthesis of Piperazines and Morpholines. *Org. Lett.* **2014**, *16*, 1236–1239. (b) Luescher, M. U.; Bode, J. W. SnAP-eX Reagents for the Synthesis of Exocyclic 3-Amino- and 3-Alkoxypyrrolidines and Piperidines from Aldehydes. *Org. Lett.* **2016**, *18*, 2652–2655.

(13) (a) Hsieh, S.-Y.; Bode, J. W. Silicon Amine Reagents for the Photocatalytic Synthesis of Piperazines from Aldehydes and Ketones. *Org. Lett.* **2016**, *18*, 2098–2101. (b) Hsieh, S.-Y.; Bode, J. W. Lewis Acid Induced Toggle from Ir(II) to Ir(IV) Pathways in Photocatalytic Reactions: Synthesis of Thiomorpholines and Thiazepanes from Aldehydes and SLAP Reagents. *ACS Cent. Sci.* **2017**, *3*, 66–72. (c) Jackl, M. K.; Legnani, L.; Morandi, B.; Bode, J. W. Org. Lett. **2017**, *19*, 4696–4699.

(14) For reviews on photoredox catalysis, see: (a) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322–5363. (b) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116*, 10075–10166. (c) Matsui, J. K.; Lang, S. B.; Heitz, D. R.; Molander, G. A. Photoredox Mediated Routes to Radicals: The Value of Catalytic Radical Generation in Synthetic Methods Development. *ACS Catal.* **2017**, *7*,

2563–2575. (d) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. J. Org. Chem. 2016, 81, 6898–6926.

(15) (a) Zhang, R.; Li, G.; Wismer, M.; Vachal, P.; Colletti, S. L.; Shi, Z.-C. Profiling and Application of Photoredox C(sp3)–C(sp2) Cross-Coupling in Medicinal Chemistry. *ACS Med. Chem. Lett.* **2018**, *9*, 773– 777. (b) Douglas, J. J.; Sevrin, M. J.; Stephenson, C. R. J. Visible Light Photocatalysis: Applications and New Disconnections in the Synthesis of Pharmaceutical Agents. *Org. Process Res. Dev.* **2016**, *20*, 1134–1147. (16) (a) Tellis, J. C.; Kelly, C. B.; Primer, D. N.; Jouffroy, M.; Patel, N. R.; Molander, G. A. Single-Electron Transmetalation via Photoredox/ Nickel Dual Catalysis: Unlocking a New Paradigm for sp³–sp² Cross-Coupling. *Acc. Chem. Res.* **2016**, *49*, 1429–1439. (b) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116*, 10035–10074. (c) Gui, Y.-Y.; Sun, L.; Lu, Z.-P.; Yu, D.-G. Photoredox sheds new light on nickel catalysis: from carbon– carbon to carbon–heteroatom bond formation. *Org. Chem. Front.* **2016**, 3, 522–526.

(17) Yoon, T. P. Visible Light Photocatalysis: The Development of Photocatalytic Radical Ion Cycloadditions. *ACS Catal.* **2013**, *3*, 895–902.

(18) For examples, see: (a) Xiao, T.; Li, L.; Zhou, L. Synthesis of Functionalized gem-Difluoroalkenes via a Photocatalytic Decarboxylative/Defluorinative Reaction. J. Org. Chem. 2016, 81, 7908–7916. (b) Lang, S. B.; Wiles, R. J.; Kelly, C. B.; Molander, G. A. Photoredox Generation of Carbon-Centered Radicals Enables the Construction of 1,1-Difluoroalkene Carbonyl Mimics. Angew. Chem., Int. Ed. 2017, 56, 15073–15077. (c) Grandjean, J.-M. M.; Nicewicz, D. A. Synthesis of Highly Substituted Tetrahydrofurans by Catalytic Polar-Radical-Crossover Cycloadditions of Alkenes and Alkenols. Angew. Chem., Int. Ed. 2013, 52, 3967–3971. (d) Perkowski, A. J.; Nicewicz, D. A. Direct Catalytic Anti-Markovnikov Addition of Carboxylic Acids to Alkenes. J. Am. Chem. Soc. 2013, 135, 10334–10337.

(19) (a) Phelan, J. P.; Lang, S. B.; Compton, J. S.; Kelly, C. B.; Dykstra, R.; Gutierrez, O.; Molander, G. A. Redox-Neutral Photocatalytic Cyclopropanation via Radical/Polar Crossover. *J. Am. Chem. Soc.* **2018**, *140*, 8037–8047. (b) Milligan, J. A.; Phelan, J. P.; Polites, V. C.; Kelly, C. B.; Molander, G. A. Radical/Polar Annulation Reactions (RPARs) Enable the Modular Construction of Cyclopropanes. *Org. Lett.* **2018**, *20*, 6840–6844.

(20) Slater, K. A.; Friestad, G. K. Mn-Mediated Radical-Ionic Annulations of Chiral N-Acylhydrazones. J. Org. Chem. 2015, 80, 6432–6440.

(21) (a) Jouffroy, M.; Primer, D. N.; Molander, G. A. Base-Free Photoredox/Nickel Dual-Catalytic Cross-Coupling of Ammonium Alkylsilicates. J. Am. Chem. Soc. **2016**, 138, 475–478. (b) Lin, K.; Kelly, C. B.; Jouffroy, M.; Molander, G. A. Preparation of Diisopropylammonium Bis(catecholato) cyclohexylsilicate. Org. Synth. **2017**, 94, 16–33.

(22) Patel, N. R.; Kelly, C. B.; Siegenfeld, A. P.; Molander, G. A. Mild, Redox-Neutral Alkylation of Imines Enabled by an Organic Photocatalyst. *ACS Catal.* **2017**, *7*, 1766–1770.

(23) (a) Otteson, D.; Michl, J. A Procedure fo Gas-Phas Dehalogenation of Organic Dihalides with Alkali Metal Vapors using Microwave and/or Ultrsound Excitation and Matrix Isolation of Products. *J. Org. Chem.* **1984**, *49*, 866–873. (b) Ohkita, T.; Tsuchiya, Y.; Togo, H. Radical 3-exo-tet Cyclization of 1,3-Dihalopropanes with SmI₂ to form Cyclopropanes. *Tetrahedron* **2008**, *64*, 7247–7251.

(24) Kandukuri, S. R.; Bahamonde, A.; Chatterjee, I.; Jurberg, I. D.; Escudero-Adán, E. C.; Melchiorre, P. X-Ray Characterization of an Electron Donor–Acceptor Complex that Drives the Photochemical Alkylation of Indoles. *Angew. Chem., Int. Ed.* **2015**, *54*, 1485–1489.

(25) Yunlin, Y.; Xiangyu, M.; Xiaotong, L.; Jinxi, X.; Guoshu, C. A sixring spiro porphyrin compound and preparation method thereof. Patent CN 108329324 A, July 27, 2018.