A pseudorotaxane-based molecular machine controlled by light and thermal stimuli[†]

Kyu-Sung Jeong,* Kyoung-Jin Chang and Young-Jae An

Department of Chemistry, Yonsei University, Seoul 120-749, S. Korea. E-mail: ksjeong@yonsei.ac.kr; Fax: (+)82-2-364-7050; Tel: (+)82-2-2123-2643

Received (in Cambridge, UK) 25th March 2003, Accepted 1st May 2003 First published as an Advance Article on the web 19th May 2003

A new pseudorotaxane-based molecular machine exhibits extremely efficient switching between assembly and disassembly mode, controlled by the combination of light and thermal stimuli.

A large variety of molecular-level machines have been developed over the last decade,¹ accompanying a remarkable advance of supramolecular chemistry. A common and fundamental feature of these machines is the reversible switching of their physicochemical properties or states according to the applied stimulation. Pseudorotaxanes have been proven to be highly useful entities for the construction of artificial molecular machines.^{1,2} This is because their assembly and disassembly can be conveniently and reversibly controlled by chemical, electrochemical and photochemical methods. Herein we report a new pseudorotaxane-based machine that displays nearly complete, light-induced interconversion between assembly and disassembly states.

As depicted in Scheme 1, the assembly–disassembly switching of the molecular machine designed here stems from the *cis*– *trans* isomerisation of the N=N bond in the threading molecule.³ In the *trans* isomer (left), the diaryldiazenyl moiety sterically blocks the above binding site and the macrocycle cannot encircle it. In the *cis* isomer (right), however, the binding site is opened to allow for the formation of the pseudorotaxane complex between two molecular components with the aid of hydrogen bonding interactions. To maintain these structural features of the *cis* and *trans* isomers, the binding site should be in parallel with the light-responsive diaryldiazenyl moiety. For this purpose, the xanthene skeleton was found to be an ideal one owing to its conformational rigidity.

Synthesis of the threading molecule (thread) **1** is outlined in Scheme 2. Diaminoxanthene **4** was prepared by nitration of 2,7-di-*tert*-butyl-9,9-dimethyl-9*H*-xanthene (**2**),⁴ followed by reduction. Reaction of **4** with 1 equivalent of nitrosobenzene provided a diazo compound **5**.⁵ Terephthaloyl dichloride was sequentially coupled in one pot with **5** and diethylamine to give the thread **1**. As the binding partner of **1**,⁶ a tetralactam macrocycle **6** has been prepared because the nitro substituents at the *para* position greatly increase the strength of hydrogen



Scheme 1 Schematic representation of working principle of a new pseudorotaxane-based machine.

[†] Electronic supplementary information (ESI) available: synthesis of thread **1** and macrocycle **6**; ¹H–¹H NOESY spectra of *trans*-**1** and *cis*-**1**; model structures of *trans*-**1** and *cis*-**1**; UV–visible spectral change of thread **1** upon irradiation; ¹H NMR spectral changes of thread **1** induced by irradiation and heating. See http://www.rsc.org/suppdata/cc/b3/b303269h/ bonding interactions.7 The ¹H NMR spectroscopic analysis indicates that the thread 1 mainly exists as the thermally more stable trans isomer with a ratio of approximately 94 : 6 (trans-1/cis-1) at 23 °C in CDCl₃, based on the ¹H NMR integration. The population of *cis*-1 was increased to approximately 78% upon irradiation with UV light (365 nm).8 The ¹H NMR spectra of cis-1 and trans-1 are noticeably different from each other. Especially, the aromatic signals, H^d, H^e and H^f, of the diaryldiazenyl moiety in cis-1 were considerably upfield shifted $(\Delta \delta = \sim 1 \text{ ppm})$ relative to those in *trans*-1. These spectral behaviors are consistent with model structures of the thread 1; upon the isomerisation of *trans*-1 into *cis*-1, two aryl rings of the diaryldiazenyl moiety become stacked together (see Electronic Supplementary Information[†]), which results in large upfield shifts of the aromatic signals due to the shielding by the adjacent ring current.

The relative binding strengths of *cis*-1 and *trans*-1 to the macrocycle 6 can be clearly noticed in Fig. 1. Addition of the thermally stable *trans*-1 (2 equiv.) to a CDCl₃ solution containing 6 (2×10^{-3} mol L⁻¹) induced only a small downfield shift ($\Delta \delta = 0.1$ ppm) of the lactam NH signal of 6 (Fig. 1b), indicating that the hydrogen bonding interaction between *trans*-1 and 6 is negligible. However, the signal was strongly downfield shifted from 8.9 to 10.0 ppm when the solution was simply irradiated with UV light⁸ and consequently the relative amount of *cis*-1 was raised to 77% (Fig. 1c). This experiment provides direct evidence for the macrocycle 6 binding much more strongly to cis-1 than to trans-1. Moreover, the signals for aromatic hydrogens, Ha and Hb, of cis-1 were significantly upfield shifted ($\Delta \delta > 2$ ppm) on complex formation. This strongly supports the proposal that the terephthalamido plane of *cis*-1 is threaded inside the cavity of 6, surrounded by aryl surfaces, to form a pseudorotaxane-type complex. The NH signal moved back to the original position on



Scheme 2 Reagents and conditions: i, HNO_3 , acetic acid, room temp. (49%); ii, Raney Ni, H_2 gas, THF, room temp. (93%); iii, nitrosobenzene, acetic acid, $CHCl_3$ (61%); iv, terephthaloyl dichloride, diethylamine, Et_3N , $CHCl_3$ (26%).

10.1039/b303269h



Fig. 1 Partial ¹H NMR spectra (left) in CDCl₃ at 5 °C of (a) macrocycle **6** (2 mM); (b) macrocycle **6** (2 mM) + *trans*-**1** (4 mM); (c) after irradiation for 3 h with UV light (365 nm) ⁸; (d) after heating (50–55 °C) for 12 h, and ¹H NMR titration curves (right) plotting NH chemical shift of macrocycle **6** against equivalent of **1** (*trans*-**1** and *cis*-**1**).

heating the solution that causes the isomerisation of *cis*-1 into *trans*-1 (Fig. 1d). This cycle can be repeated by continuously alternating irradiation and heating, which results in reversible switching between assembly and disassembly modes.

The association constants between the thread 1 (cis-1 and trans-1) and the macrocycle 6 were determined and compared by ¹H NMR titration experiments. The experiments were performed in CDCl₃ at a low temperature, 5 °C, to prevent possible thermal isomerisation of the thread 1. Under these titration conditions, time-averaged resonances for the free and the complexed species were always observed. As demonstrated in Fig. 1 (right), the NH chemical shift changes of 6 induced by *trans*-1 are very small compared to those induced by *cis*-1. Even these small changes may be attributed to a small amount (~ 6%) of cis-1 existing in the stock solution of trans-1. The association constants (K_a) were calculated by nonlinear least squares fitting methods⁹ and found to be $\leq 1 \text{ M}^{-1}$ for *trans*-1 and $5200 \pm 100 \text{ M}^{-1}$ for *cis*-1,¹⁰ respectively. The difference in binding affinities is nearly four orders of magnitude, one of the highest exhibited by light-driven molecular machines developed to date.

In conclusion, we have described a new molecular machine based on a pseudorotaxane whose assembly occurs only with the *cis* isomer of the threading molecule. On the other hand, the pseudorotaxane is completely disassembled into its molecular components on isomerisation to the *trans* isomer caused by an external stimulus.

This work was financially supported by the Korea Science and Engineering Foundation (R02-2002-000-00115-0).

Notes and references

 For reviews, see: (a) V. Balzani, M. Gómez-López and J. F. Stoddart, Acc. Chem. Res., 1998, 31, 405; (b) V. Balzani, A. Credi, F. M. Raymo and J. F. Stoddart, Angew. Chem., Int. Ed., 2000, 39, 3348; (c) Acc. Chem. Res., 2001, 34, (6)) (Special Issue on Molecular Machines).

- 2 (a) P. R. Ashton, V. Balzani, O. Kocian, L. Prodi, N. Spencer and J. F. Stoddart, J. Am. Chem. Soc., 1998, **120**, 11190; (b) M. Asakawa, P. R. Ashton, V. Balzani, C. L. Brown, A. Credi, O. A. Matthews, S. P. Newton, F. M. Raymo, A. N. Shipway, N. Spencer, A. Quick, J. F. Stoddart, A. J. P. White and D. J. Williams, Chem. Eur. J., 1999, **5**, 860; (c) E. Ishow, A. Credi, V. Balzani, F. Spadola and L. Mandolini, Chem. Eur. J., 1999, **5**, 984; (d) J. W. Lee, K. Kim and K. Kim, Chem. Commun., 2001, 1042; (e) V. Balzani, A. Credi, F. Marchioni and J. F. Stoddart, Chem. Commun., 2001, 1860; (f) R. Ballardini, V. Balzani, M. Clemente-León, A. Credi, M. T. Gandolfi, E. Ishow, J. Perkins, J. F. Stoddart, H.-R. Tseng and S. Wenger, J. Am. Chem. Soc., 2002, **124**, 12786.
- 3 For some examples of azobenzene-containing molecular machines, see: (*a*) H. Murakami, A. Kawabuchi, K. Kotoo, M. Kunitake and N. Nakashima, *J. Am. Chem. Soc.*, 1997, **119**, 7605; (*b*) C. A. Stanier, S. J. Alderman, T. D. W. Claridge and H. L. Anderson, *Angew. Chem., Int. Ed.*, 2002, **41**, 1769 and see also the references 2*c* and 2*e*.
- 4 J. S. Nowick, P. Ballester, F. Ebmeyer and J. Rebek, Jr., J. Am. Chem. Soc., 1990, 112, 8902.
- 5 (a) W. Shuangxi and C. A. Rigiberto, *Org. Lett.*, 2001, 3, 3831; (b) M.
 H. Davey, V. Y. Lee, R. D. Miller and T. J. Marks, *J. Org. Chem.*, 1999, 64, 4976.
- 6 For studies of binding between terephthalamides and tetralactam metallomacrocycles, see: (a) C. A. Hunter and L. D. Sarson, Angew. Chem., Int. Ed. Engl., 1994, 33, 2313; (b) K.-S. Jeong, Y. L. Cho, J. U. Song, H.-Y. Chang and M. G. Choi, J. Am. Chem. Soc., 1998, 120, 10982; (c) K.-S. Jeong, J. W. Lee, T.-Y. Park and S.-Y. Chang, Chem. Commun., 1999, 2069. For synthesis and spectroscopic properties of 6, see Electronic Supplementary Information[†].
- 7 H. S. Kim and K.-S. Jeong, unpublished results. For metallomacrocycles showing similar substituent effects, see: K.-S. Jeong, Y. L. Cho, S.-Y. Chang, T.-Y. Park and J. U. Song, *J. Org. Chem.*, 1999, **64**, 9459.
- 8 A common UV lamp for TLC spot detection was used as the UV light source (365 nm, long-wave). The ¹H NMR spectral change was negligible after 3 h irradiation.
- 9 R. S. Macomber, J. Chem. Educ., 1992, 69, 375; L. Fielding, Tetrahedron, 2000, 56, 6151.
- 10 In the titration of **6** with *cis*-**1**, the stock solution of *cis*-**1** contained 30% of *trans*-**1** based on the ¹H NMR integrations. Therefore, the concentrations of *cis*-**1** have been corrected for the calculation of the association constants (K_a).