

ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: B. Zhang and X. Zhang, *Chem. Commun.*, 2015, DOI: 10.1039/C5CC08394J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

Pd-Catalyzed *gem*-Difluoroallylation of Arylboronic Acids with γ , γ -Difluoroallylic Acetates

Bo Zhang^a and Xingang Zhang^{*a}

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

A highly regio- and stereo-selective palladium-catalyzed *gem*-difluoroallylation of arylboronic acids with γ , γ -difluoroallylic acetates has been described. The method allows the synthesis of a variety of *gem*-difluoroallylated arenes with a tosyloxy group on the C=C double bond, thus providing a good opportunity for the down-stream transformations.

As a distinct class of organic compounds in pharmaceuticals and agrochemicals, difluoroalkylated arenes have been received great attentions recently.^{1,2} This is because the introduction of difluoromethylene (CF₂) group at benzylic position can dramatically improve the metabolic stability and change the physiological activity of the bioactive molecules.³ Among the numerous difluoroalkylated arenes, *gem*-difluoroallylic arenes represent one of the extremely appealing compounds owing to their important applications in medicinal chemistry⁴ and versatile synthetic utility of the C=C double bond. The most common approach to prepare such a valuable structural motif relies on the Wittig reaction or HWE reaction from aryl difluoromethylated aldehydes or ketones.⁵ Despite the utility of this approach, multiple steps were required to prepare the difluoroalkylated starting materials, which restricts its widespread synthetic applications. In particular, this approach is not suitable for the late stage synthesis of difluoroalkylated complexes due to the requirement of difluoroalkylated precursors. From the point of view of synthetic simplicity and generality, a direct introduction of *gem*-difluoroallyl group onto aromatic rings would be an attractive alternative (Scheme 1a).^{6,7} However, the regiochemical selectivity of this synthetic strategy and suppressing undesired defluorination side reaction

resulted from *gem*-difluoroallylated products remain the synthetic challenges thus far.⁷

Several years ago, a palladium catalyzed reaction of 3,3-difluoro-2-propen-1-ol acetate with phenylzinc chloride was reported by Shi.^{6a} Although a high γ -substitution regioselectivity was observed, an undesired defluorination of resulting *gem*-difluoroallylic product occurred (Scheme 1b). In this context, a sterically hindered 1-substituted 3-bromo-3,3-difluoropropene (BDFP) was used as a substrate, leading to *gem*-difluoroallyl product in high regioselectivity without observation of defluorination side reaction.⁷ However, the reaction of BDFP with phenylzinc chloride or tributylphenyltin still resulted in defluorinated product. To address this crucial issue, very recently, a highly efficient method to prepare *gem*-difluoroallylated arenes through palladium-catalyzed reaction of organoborons with bromodifluoromethylated alkenes has been developed by us,⁸ which represents a first example to prepare such a kind of structure catalyzed by transition-metal (Scheme 1c). Inspired by this preliminary study, we questioned that whether γ , γ -difluoroallylic electrophiles can also serve as suitable coupling partner to prepare *gem*-difluoroallylated arenes (Scheme 1d). Herein, we described a Pd-catalyzed *gem*-difluoroallylation of arylboronic acids with γ , γ -difluoroallylic acetates. The method provides a facile access to a variety of *gem*-difluoroallylated arenes with high *regio*- and *stereo*-selectivity.

Scheme 1. Strategies for regio-selective synthesis of *gem*-difluoroallylated arenes

^aKey Laboratory of Organofluorine Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 345 Lingling Road, Shanghai 200032, China. E-mail: xqzhang@mail.sioc.ac.cn

Fax: +86 (21) 64166128; Tel: +86 (21) 54925333

[†]Electronic Supplementary Information (ESI) available: Detailed experimental procedures, and characterization data for new compounds. See DOI: 10.1039/x0xx00000x

We began our studies by choosing 3,3-difluoro-1-phenyl-2-(tosyloxy) allyl acetate **1a** and phenyl boronic acid **2a** as model substrates (Table 1). The use of **1a** is because it can be easily prepared from low-cost and widely available 2,2,2-trifluoroethanol.⁹ In addition, the transformations of tosyloxy group on the C=C double bond can easily lead to a variety of difluoroalkylated arenes.¹⁰ To our delight, a 37% yield of the desired product **3a** with high stereoselectivity (Z/E > 20:1, determined by ¹⁹F NMR) was obtained when the reaction was carried out with **1a** (1.0 equiv), **2a** (1.5 equiv), Pd(PPh₃)₄ (10 mol %), K₂CO₃ (2.0 equiv) in dioxane at 80 °C (entry 1). Notably, no branched γ,γ-difluoroallylic isomer and defluorinated side product were observed in the reaction process. We reasoned that this is probably because of the steric effect of the substituent. Further optimization of the reaction conditions by examining the phosphine ligands (entries 2-8) revealed that the bidentate ligand Xantphos¹¹ was the best ligand, providing **3a** in 57% yield (entry 8). The dppf and the monodentate ligand Me-Phos also provided moderate yields, but other ligands showed less activity (entries 3 and 6).

Table 1 Representative results for optimization of Pd-catalyzed *gem*-difluoroalkylation of **2a** with **1a**^a

entry	[Pd]	Ligand	Additive	yield (%) ^b
1	Pd(PPh ₃) ₄	--	--	37
2	Pd(PPh ₃) ₄	PPh ₃	--	42
3	Pd(PPh ₃) ₄	Me-Phos	--	53
4	Pd(PPh ₃) ₄	S-Phos	--	45
5	Pd(PPh ₃) ₄	dppe	--	18
6	Pd(PPh ₃) ₄	dppf	--	50
7	Pd(PPh ₃) ₄	BINAP	--	43
8	Pd(PPh ₃) ₄	Xantphos	--	57
9	Pd ₂ (dba) ₃	Xantphos	--	49
10	Pd(OAc) ₂	Xantphos	--	ND
11 ^c	Pd(PPh ₃) ₄	Xantphos	--	64
12 ^d	Pd(PPh ₃) ₄	Xantphos	--	72 (68)
13 ^e	Pd(PPh ₃) ₄	Xantphos	--	30
14 ^d	Pd(PPh ₃) ₄	Xantphos	H ₂ O (1.25)	71 (72)
15 ^d	Pd(PPh ₃) ₄	Xantphos	CuI (0.06)	89 (84)


^aReaction conditions (unless otherwise specified): **1a** (0.2 mmol, 1.0 equiv), **2a** (0.3 mmol, 1.5 equiv), dioxane (1.5 mL), K₂CO₃ (2.0 equiv), 5 h under N₂. ^bDetermined by ¹⁹F NMR using fluorobenzene as an internal standard and number in parenthesis is isolated yield. Z/E > 20:1, determined by ¹⁹F NMR. ^cPd(PPh₃)₄ (5 mol %), Xantphos (5 mol %). ^dPd(PPh₃)₄ (5 mol %), Xantphos (6 mol %). ^ePd(PPh₃)₄ (1 mol %), Xantphos (1.2 mol %).

Encouraged by these results, a survey of the reaction parameters, such as palladium sources, bases, and solvents were conducted (entries 8-10, for details, see the Supporting Information). The combination of Pd(PPh₃)₄, dioxane, and K₂CO₃ remained the optimal choice (entry 8). Decreasing the loading amount of Pd(PPh₃)₄ from 10 mol % to 5 mol % with a ratio of

Xantphos/Pd(PPh₃)₄ = 1.2 benefited the reaction efficiency, providing **3a** in 68% yield upon isolation (entry 12). However, the meanwhile, the homocoupling of **2a** and defluorination of **3a** were also observed. Taking into account the fact that the addition of H₂O or CuI can facilitate the transmetalation step of the Suzuki reaction,¹² H₂O and CuI were employed as additives to improve the reaction efficiency further. A best yield of **3a** (84% upon isolation) was obtained when CuI (6 mol %) was used (entry 15). H₂O also showed beneficial effect, but led to slightly lower yield (72% upon isolation, entry 14).

To demonstrate the substrate scope of this method, a variety of arylboronic acids were examined to react with **1a** (Table 2).

Table 2 Pd-catalyzed *gem*-difluoroalkylation of arylboronic acids **2** with **1a**^a




^aReaction conditions (unless otherwise specified): **1a** (0.4 mmol, 1.0 equiv), **2** (0.6 mmol, 1.5 equiv), K₂CO₃ (2.0 equiv) and dioxane (3 mL). All reported yields are isolated yields and the stereoselectivity of **3** is Z/E > 20:1. ^bCuI (6 mol %) instead of H₂O was used. ^cReaction ran for 15 h. ^dReaction run on a gram-scale.

Generally, moderate to high yields of **3** were obtained without observation of branched isomers and electron-rich arylboronic acids provided higher yields than that from electron-deficient ones. Most importantly, the tosyloxy group was compatible with the reaction without formation of diarylated byproducts, thus providing a good opportunity for downstream

transformations. However, it should be mentioned that in most of the cases, H₂O was proved to be an optimal additive. Versatile functional groups, such as vinyl, silyl, ester, enonizable ketone, and nitro, showed good tolerance (**3e**, **3f**, **3i**, and **3m-o**). The sterically hindered substrates were also applicable to the reaction, providing the corresponding products in moderate to good yields (**3k** and **3l**). The reaction can also be extended to other functionalized γ,γ -difluoroallylic acetates **1**, in which the substituents R did not interfere with the reaction efficiency. Notably, high yields of **3r** and **3s** were also obtained when R is an alkyl group, thus highlighting the generality of this method. It was also possible to synthesize compounds **3** on a gram-scale. A good yield of **3a** (76%) was obtained on a gram-scale synthesis, thus demonstrating the reliability of current reaction.

To demonstrate the utility of this method, the transformations of *gem*-difluoroallylated arene **3a** were performed.



Scheme 2 Transformations of compound **3a**

As shown in Scheme 2a, hydrogenation of **3a** provided difluoroalkylated arene **4** in excellent yield. Suzuki cross-coupling of **3a** with phenylboronic acid **2a** also underwent smoothly and provided trisubstituted **5** in 93% yield, which was further hydrogenated to produce compound **6** with high efficiency (Scheme 2b). This is noteworthy as difluoroalkylated arenes have important applications in medicinal chemistry.¹³ Compound **3a** can also be converted into difluoroalkylated ketone **7** smoothly, thus providing an alternative approach to prepare such a kind of fluorinated compound (Scheme 2c). Interestingly, the tosyloxy group can be easily substituted by methoxy group when compound **3a** was treated with MeOH

under basic condition at room temperature (Scheme 2d). Unexpectedly, extension of this strategy to replace tosyloxy group with amide group produce an unstable enamine intermediate, which subsequently eliminated under basic condition to afford an useful *gem*-difluoroallylated arene **3** (Scheme 2e), thus demonstrating the versatility of the compounds **3**.

In conclusion, we have developed an efficient and alternative strategy to prepare *gem*-difluoroallylated arenes through palladium-catalyzed reaction between arylboronic acids and γ,γ -difluoroallylic acetates with high *regio*- and *stereo*-selectivity. Application of the method led to different difluoroalkylated arenes efficiently, which may have potential applications in medicinal chemistry. The reason for high regiochemical selectivity of current reaction is probably because the reductive elimination of aryl(*gem*-difluoroallyl)palladium complex prefers generating of linear product due to the steric effect of the substituents.¹⁴ However, for the detailed reaction mechanism, it remains a point for discussion.

This work was financially supported by the National Basic Research Program of China (Nos. 2012CB821600 and 2015CB931900), the National Natural Science Foundation of China (21425208, 2141002, 21172242 and 21332010), and SIC.

Notes and references

- For selected reviews, see: (a) B. E. Smart, *J. Fluorine Chem.*, 2001, **109**, 3; (b) P. Maienfisch and R. G. Hall, *Chimia*, 2004, **58**, 93; (c) N. A. Meanwell, *J. Med. Chem.*, 2011, **54**, 2529; (d) Ni, L. Zhu and J. Hu, *Acta Chim. Sinica*, 2015, **73**, 90; (e) M.-C. Belhomme, T. Besset, T. Poisson and X. Pannecoucke, *Chem. Eur. J.*, 2015, **21**, 12836.
- For selected transition-metal-catalyzed difluoroalkylation, see: (a) Z. Feng, F. Chen and X. Zhang, *Org. Lett.*, 2012, **14**, 193; (b) Z. Feng, Q.-Q. Min, Y.-L. Xiao, B. Zhang and X. Zhang, *Angew. Chem. Int. Ed.*, 2014, **53**, 1669; (c) S. Ge, W. Chalovich and J. F. Hartwig, *J. Am. Chem. Soc.*, 2014, **136**, 4149; (d) Y.-L. Xiao, W.-H. Guo, G.-Z. He, Q. Pan and X. Zhang, *Angew. Chem. Int. Ed.*, 2014, **53**, 9909.
- For selected examples of bioactive molecules, see: (a) K. Shen, Y.-F. Keng, L. Wu, X.-L. Guo, D. S. Lawrence and Z.-Y. Zhang, *Biol. Chem.*, 2001, **276**, 47311; (b) C. S. Burgey, K. A. Robinson, T. A. Lyle, P. E. J. Sanderson, S. D. Lewis, B. J. Lucas, J. A. Krueger, R. Singh, C. Miller-Stein, R. B. White, B. Wong, E. Lyle, P. D. Williams, C. A. Coburn, B. D. Dorsey, J. C. Barrow, M. T. Stranieri, M. A. Molahan, G. R. Sitko, J. J. Cook, D. H. McMasters, C. M. McDonough, W. M. Sanders, A. A. Wallace, F. C. Clayton, D. Bohn, Y. M. Leonard, T. J. Detwiler Jr., J. Lynch Jr., Y. Yan, Z. Chen, L. Kuo, S. J. Gardell, J. A. Shafer and J. P. Vacca, *J. Med. Chem.*, 2003, **46**, 461; (c) M. O. Anderson, J. Zhang, Y. Liu, C. Yao, P.-W. Phuan and A. S. Verkman, *J. Med. Chem.*, 2012, **55**, 5942; (d) J. O. Link, J. G. Taylor, L. Xu, A. Mitchell, H. Guo, H. Liu, D. Kato, T. Kirschberg, J. Sun, N. Squires, J. Parrish, T. Keller, Z.-Y. Yang, C. Yang, M. Matles, Wang, K. Wang, G. Cheng, Y. Tian, E. Mogalian, E. Mondou, M. Cornpropt, J. Perry and M. C. Desai, *J. Med. Chem.*, 2014, **57**, 2033.
- (a) Y. Gai, Y. S. Or and G. Wang, *From PCT Int. Appl.*, WO/2010132163 A1 20101118, 2010; (b) C. W. Amdnde, B. A. Fish, M. E. Green, D. S. Jhonson, P. B. Mullins, C. J. O'Donnell, M. Pettersson, C. M. Stiff, C. Subramanyam, T. P. Tran and

- Navaratnam, From PCT Int. Appl., WO 2012131539 A1 20121004, 2012.
- 5 (a) J. Oldendorf and G. Haufe, *J. Prakt. Chem.*, 2000, **342**, 52; (b) S. Pikul, G. E. Mieling, K. K. Mieling, K. M. Solinsky, B. De, N. G. Almstead and N. M. George, From U.S. Pat. Appl. Publ., 20030139453 A1 20030724, 2003; (c) M. Bergeron, T. Johnson and J.-F. Paquin, *Angew. Chem. Int. Ed.*, 2011, **50**, 11112.
 - 6 (a) G.-q. Shi, X.-h. Huang and F.-J. Zhang, *Tetrahedron Lett.*, 1995, **36**, 6305; (b) G.-q. Shi and W.-I. Cai, *Synlett.*, 1996, **4**, 371.
 - 7 M. Kiriwara, T. Takuwa, M. Okumura, T. Wakikawa, H. Takahata, T. Momose, Y. Takeuchi and H. Nemoto, *Chem. Pharm. Bull.*, 2000, **48**, 885 and ref 5a.
 - 8 Q.-Q. Min, Z. Yin, Z. Feng, W.-H. Guo and X. Zhang, *J. Am. Soc. Chem.*, 2014, **136**, 1230.
 - 9 (a) G. H. Posner, B. T. Woodard, K. R. Crawford, S. Peleg, A. J. Brown, P. Dolan and T. W. Kensler, *Bioorg. Med. Chem.*, 2002, **10**, 2353; (b) P. V. Ramachandran and A. Chatterjee, *Org. Lett.*, 2008, **10**, 1195.
 - 10 For selected examples of transition-metal-catalyzed cross-coupling using alkenyl tosylates as a coupling partner, see: (a) M. Fujiwara, J. Ichikawa, T. Okauchi and T. Minami, *Tetrahedron Lett.*, 1999, **40**, 7261; (b) T. M. Gøgsig, L. S. Søbjerger, A. T. Lindhardt, K. L. Jensen and T. Skrydstrup, *J. Org. Chem.*, 2008, **73**, 3404; (c) J. Pschierer, N. Peschek and H. Plenio, *Green Chem.*, 2010, **12**, 636; (d) H. Zhang, C.-B. Zhou, Q.-Y. Chen, J.-C. Xiao and R. Hong, *Org. Lett.*, 2011, **13**, 560; (e) J. Bucher, T. Wurm, K. S. Nalivela, M. Rudolph, F. Rominger and A. S. K. Hashmi, *Angew. Chem. Int. Ed.*, 2014, **53**, 3854.
 - 11 (a) V. V. Grushin and W. J. Marshall, *J. Am. Soc. Chem.*, 2006, **128**, 12644; (b) V. I. Bakhmutov, F. Bozoglian, K. Gómez, G. González, V. V. Grushin, S. A. Macgregor, E. Martin, F. M. Miloserdov, M. A. Novikov, J. A. Panetier and L. V. Romashov, *Organometallics*, 2012, **31**, 1315.
 - 12 (a) G. B. Smith, G. C. Dezeny, D. L. Hughes, A. O. King and T. R. Verhoeven, *J. Org. Chem.*, 1994, **59**, 8151; (b) M. B. Thathagar, J. Beckers and G. Rothenberg, *J. Am. Soc. Chem.*, 2002, **124**, 11858.
 - 13 For selected examples, see: (a) H. Eto, Y. Kaneko and T. Sakamoto, *Chem. Pharm. Bull.*, 2000, **48**, 982; (b) F. Xue, H. Li, S. L. Delker, J. Fang, P. Martasek, L. J. Roman, T. L. Poulos and R. B. Silverman, *J. Am. Chem. Soc.* 2010, **132**, 14229 and ref 3c.
 - 14 (a) T. Hayashi, A. Yamamoto and T. Hagihara, *J. Org. Chem.*, 1986, **51**, 723; (b) B. Aakermark, S. Hansson and A. Vitagliano, *J. Am. Soc. Chem.*, 1990, **112**, 4587; (c) A. Togni, U. Burckhardt, V. Gramlich, P. S. Pregosin and R. Salzmänn, *J. Am. Soc. Chem.*, 1996, **118**, 1031.