

PII: S0040-4020(96)00752-1

# 1,2-Dithiines and Precursors, XVI<sup>1</sup>: Synthesis, Structure, and Reactivity of Non-Anellated 1,2-Dithiines\*

## Werner Schroth<sup>\*</sup>, Simona Dunger<sup>2</sup>, Frank Billig, Roland Spitzner, Rainer Herzschuh, Almut Vogt, Thomas Jende, Gunter Israel, Jens Barche, Dieter Ströhl

Institut für Organische Chemie der Martin-Luther-Universität Halle-Wittenberg, Kurt-Mothes-Straße 2, D-06099 Halle (Saale)

#### Joachim Sieler<sup>3</sup>

Institut für Anorganische Chemie der Universität Leipzig, Linnéstraße 3, D-04103 Leipzig

Abstract: Various monocyclic 1,2-dithiines 6a,b,d-t were prepared via (Z,Z)-1,4-difunctionalized butadienes (4-11,19,20). A twisted cyclic structure A is unequivocally proved rather than of the ringopened valence isomer B. The reactivity of these 1,2-dithiines is described. Thermal as well as day-light induced sulfur extrusion is an important feature of their chemistry. The latter mode of sulfur extrusion depends to a significant extent on absorption in the visible region. Copyright © 1996 Elsevier Science Ltd

### Introduction

Concurrent with our initial synthetic work<sup>1a-c</sup> on the title compounds A (*Scheme 1*) various 3,6-di(polyynyl) substituted  $C_{13}$ -representatives were discovered as plant constituents.<sup>4</sup> As subsequently reported,<sup>5</sup> these compounds ("thiarubrines") show promising biological activities. In view of these properties, 1,2-dithiines are becoming important compounds with respect to their synthesis, structure elucidation and reactivity.<sup>6</sup> From the beginning of our programme, our studies have focused on some other outstanding aspects i.e. a) the controversy of a cyclic disulfide structure versus a ring-opened valence isomer **B** (as disclosed in the related 1,2-



Scheme 1

dithiete series)<sup>6i,7</sup>, b) the presence of an unusual red colour, and c) the sulfur extrusion under mild conditions. The cyclic title system A represents formally an  $8\pi$ -electron system, a butadiene-bridged disulfide, and a sulfur-expanded thiophene, respectively. Theoretical treatment of these molecules has also been well documented.<sup>6g-j</sup>

Significant differences can exist both in the synthesis as well as the behaviour of non-anellated and anellated 1,2-dithiines. Thus we have carefully differentiated between both classes in our considerations. In general however, the synthesis of both types can be achieved from (Z,Z)-buta-1,3-dien-1,4-dithiols or their salts which suffer oxidation in the final step.<sup>8</sup>

The synthesis of non-anellated 1,2dithilnes starts from C<sub>4</sub>-precursors, especially butadiynes (C, D) as illustrated in Scheme 2. S-Deblocking of the intermediates E, R = protective group, affords the above mentioned (Z,Z)-butadiene-1,4-dithiols E (R = H) leading finally to maximally (in 3- and 6-posi-



tions) disubstituted 1,2-dithines  $\mathbf{F}$ . Access to  $\mathbf{E}$  is differentiated by the origin of the RS group. In our initial synthesis (path *a*), <sup>1a-c</sup> a two-fold nucleophilic thiol addition at C produces  $\mathbf{E}$ .<sup>9</sup> The whole sequence represents formally the addition of hydrogen sulfide to a butadiyne in an appropriate protected form. This method has been modified recently by *M. Koreeda* and *W. Yang*<sup>6b</sup> and by *D. E. Bierer* et al.<sup>6d</sup> In the other alternative (path *b*), recently reported by *E. Block* et al.,<sup>6c</sup> the RS-group is already installed in the butadiyne precursor  $\mathbf{D}$ , and the substituents R' and R'' are introduced subsequently by C,C-coupling via organotin functionalization. Both strategies complement each other.

The synthesis of anellated 1,2-dithilines is based on the successive assembly of the (Z,Z)-1,4-dithiobuta-1,3-diene unit starting from appropriate functionalized cyclic precursors (cf. references <sup>lh-o</sup>).

In view of the latest developments in the area of 1,2-dithiines a detailed presentation of various results of our current work now seems appropriate.

### **Results and Discussion**

#### 1. Synthesis

As outlined in *Scheme 3*, the synthesis can proceed via two sequences depending upon the choice of the Sprotective group (i.e. according to R in Scheme 2, path a). Its potential is demonstrated by the listed 1,2dithines 6 (cf. our initial reports in ref.<sup>1a,c,d</sup>). These include the parent compound (**6a**) as well as alkyl (**6b,d**g), aryl (**6h-s**) and hetaryl substituted 1,2-dithines (**6t**). The products 6 form, at room temperature, mainly yellow to red crystals, with exception of the oily parent compound **6a** and the 3,6-di-*n*-butyl-1,2-dithine (**6b**). The latter compound is *not* identical with a compound previously claimed to possess this structure and obtained by a different procedure.<sup>10</sup>

In sequence I the butadiynes 1 undergo a regio- and stereoselective nucleophilic bis-addition of benzylmercaptan (2) under basic conditions leading to the (Z,Z)-1,4-dithiobutadienes 4 (step [i]). If required the mono-adducts 3 can be isolated (steps [ia]/[ib]).<sup>11</sup> In the case of aryl substituents in 1 (h-s) the addition at the terminal  $\alpha$ - and  $\alpha$ '-positions of the butadiyne chain is strictly observed even although arylacetylenes normally undergo thiol addition at the  $\beta$ -position relative to the aryl group (cf. also example 15 in Scheme 4),<sup>12</sup> thus indicating the dominating influence of an adjacent triple bond compared with an aryl substituent. The reductive debenzylation of 4 was performed by means of sodium in liquid ammonia at -70°C (step ii). The resulting dithiolate salts 5 (M = Na) are immediately oxidized to the 1,2-dithiines 6, either by potassium cyanoferrate(III), ferric chloride, iodine or even air (step iii).

The thiol-addition to 1, the most crucial part of the whole sequence, is favoured by acceptor and disadvantaged by donor effects in the substituents  $R^1$  and  $R^2$ . For example, the *t*-butyl-diyne 1c yields only the monoadduct 3c in spite of an excess of 2, whilst 1d smoothly affords the bis-adduct 4d. Various procedures are possible. Thus 4h is obtainable from 1h and 2 by reaction in EtOH/NaOH at elevated temperature (sometimes more profitable under in situ conditions using benzylisothiouronium chloride) in a yield of about 55%, in benzene/KOH/18-crown-6 at 20°C, in 68% yield, in diglyme/KOH with (58%) or KO-*t*-Bu[catalytic] at room temperature, in 61% yield, or in sulfolane/KOH or KO-*t*-Bu[catalytic] at 20°C, in 62% yield. In DMSO/KOH cyclization of the mono-adducts 2 takes place forming thiophene formation due to the addition of CH<sub>2</sub> of the benzyl group to the triple bond.<sup>13a</sup> In general the twofold thiol addition in dimethylformamide with catalytic amounts of KOH, initially proposed by *M. Koreeda* and *W. Yang*,<sup>6a</sup> seems to be the most suitable technique (as previously stated,<sup>2a</sup> NaOH in place of KOH is not as efficiently).

The debenzylation of 4 (step [*ii*]) by alkali metal in liquid ammonia is disadvantaged by substituents which can suffer additional reductive attack, such as halogen (e.g.  $4i \rightarrow 6h$ ) and unsaturated groups (as in the synthesis of thiarubrines, cf. ref.<sup>6b,c</sup>)<sup>13b</sup>. A *selective* reductive debenzylation despite the presence of such substituents, however, is achievable with the aid of lithium 1-(N,N-dimethylamino)naphthalenide as described in ref.<sup>6c</sup>. An advantageous alternative consists of the use of S-protective groups other than CH<sub>2</sub>Ph as demonstrated recently with CH<sub>2</sub>CH<sub>2</sub>SiMe<sub>3</sub> <sup>6b</sup> and CH<sub>2</sub>CH<sub>2</sub>CN <sup>6d</sup> (R in Scheme 2) which allow deprotection by fragmentation or  $\beta$ -elimination, respectively, in the subsequent step. Another non-reductive technique is described in the following.

In sequence II t-butylmercaptan (7) can be used as protected thiol supplier (cf. ref.<sup>14</sup>). After uncomplicated addition of 7 to 1, the resulting (Z,Z)-1,4-di(t-butylthio)butadienes 8 are deprotected by reaction with (2-nitrophenyl)sulfenyl chloride affording the poorly soluble mixed bis-disulfides 9 (steps iv, v). From the latter the 1,2-dithiines 6 are smoothly produced by reduction with sodium borohydride and subsequent oxidation, or better by treatment with catalytic amounts of 2-mercaptoethanol, with facile removal of the co-formed bis(2-nitrophenyl) disulfide (step vi). Even attempts to recrystallize the 9, e.g. from DMF, symmetrization to the 1,2-dithiines 6 takes place spontaneously and the colour of the solution becomes red. This method is especially useful in the aromatic series with good yields throughout (examplified by 6 h.j.l).



From 5 the S-acetylderivatives 11 are obtained (by treatment with acetic anhydride or acetyl chloride) and since these can readily be saponified back to the 5 (e.g. with methanolic KOH) they can therefore be used as storable sources for the latter compounds (steps vii, viii). Furthermore, careful acidification of 5 enables the liberation of the (Z,Z)-buta-1,3-diene-1,4-dithiols  $10^{15}$  which can also be be prepared by reduction of the 1,2-dithines 6 with sodium borohydride. The latter compounds can be transformed into the thiophenes 12 by heating thus illustrating an intermediate step within the classical Paal-Knorr synthesis of thiophenes from 1,4-diketones (step x). Thiophenes 12 also result from 6 by extrusion of sulfur (step ix, see also part 2). The favoured formation of 12 is likewise indicated in the mass spectroscopic fragmentation of 6 or 10 and even of the precursors 4 and 8 (base peaks).

The intermediates 5 undergo other characteristic substitution reactions at the thiolate group which can occasionally be performed by use of the equivalents 10 or 11. As outlined in Scheme 4 the reaction with benzyl chloride leads back to the precursors 4, and with methyl iodide to the methylthio derivatives 13. The analogous ethyl thioethers 14 are identical with those produced by the addition of ethylmercaptan to the diacetylenes 1. Finally, vinylation of 5h with phenylacetylene results in the formation of the (Z)-styryl sulfide 15h. By contrast, alkylating ring closure reactions succeed only to rather limited extent. For example, the reaction of 5h with o-di(bromomethyl)benzene gives the ten-membered cyclic thioether 16 in low yield with thiophene 12h and 1.3-dihydro-benzo[c]thiophene (intrachain cyclization). Moreover, it should be emphasized that all attempts to obtain 1,4-dithiocines by ring closure reactions of 5a or 5h with both (Z)-1,2- and 1,1dichloroethene failed.<sup>16a</sup>

#### Sch

Various efforts to produce 1,2-dithiines 6 starting from 1,4-dicarbonyl compounds 17 instead of the diacetylenes 1, as illustrated in Scheme 5, were disappointing.<sup>16b</sup> Thus, the acid-assisted reaction of aromatic 1,4diketones with ethanethiol, indeed, affords the dimercaptals G and subsequently the 1,4-di(ethylthio)butadienes 14.<sup>1e</sup> But in attempts for their reductive deprotection with the aid of sodium in liquid ammonia the Birch reduction competes successfully, e.g. from 14h (Ar =  $C_6H_5$ ) (E)-1,4-diphenylbutene (18) is formed, and ethylmercaptan is liberated during work-up. The intermediate **H** of the Birch reduction offers two new benzylthio subunits which subsequently suffer a reductive fission at the indicated positions.



#### Scheme 5

By alkylation of the 14 with dimethyl sulfate (to disulfonium salts) and subsequent elimination by means of sodium hydroxide, the diacetylenes 1 can be obtained.<sup>1e</sup> Hence, the 14 also offer access to 1,2-dithiines 6 starting from 1,4-dicarbonyl compounds 17, albeit in a circuitous manner.

On the other hand, the acid-catalyzed reaction of the 17 with benzylmercaptan did not yield the required 1,4-di(benzylthio)-1,3-butadienes 4 and only the thiophenes 12,  $R^1 = R^2 = Ar$ , could be isolated. Here the initially produced 4 suffer cationic debenzylation. This was readily confirmed by the formation of 12,  $R^1$  =  $R^2 = Ar$ , from an authentic samples of 4 as well as from the mono-addition products 3,  $R^1 = R^2 = Ar$ , under the above noted conditions.

An alternative approach to 3,6-disubstituted 1,2-dithiines 6 involves the successive functional conversion of a suitable butadiene precursor as previously reported for the di-anellated series:<sup>1k</sup> This is illustrated in Scheme 6 where (Z,Z)-1,4-dilithio-1,4-diphenyl-butadiene (20), obtained from the corresponding dibromocompound 19 by halogen-metal exchange with *n*-butyllithium,<sup>17</sup> is treated (in situ) with elemental sulfur giving the 1,2-dithine **6h** directly in a good over-all yield. In this case no final oxidation step is necessary, possibly due to an intramolecular nucleophilic attack on the sulfur chain by the remaining anionic centre as shown in I. This method is, of course, limited by the availability of the dibromo-precursors 19.



#### 2. Structure and Reactivity

The surprising colour of the 1,2-dithines **6** with absorptions up to nearly 500 nm<sup>18</sup> should be due to a  $n \rightarrow \sigma^*$  transition.<sup>6g,i</sup> This is undoubtedly due to the disulfide group.<sup>19</sup> By contrast the isomeric 1,4-dithiines<sup>20</sup> show no discrete absorption maximum in the visible region. A clear dependence of the wave length in the visible region of the electronic spectra of **6** on the subsituents is not recognizable at this time (see *Table 1*). The absorptiom maximum in the visible range appears to be bathochromicly shifted with

Table 1. Absorption maxima in the visible region

Entry	$\lambda_{max} [nm] (lg \epsilon)$	Entry	$\lambda_{max} [nm] (lg \epsilon)$	Entry	$\lambda_{max} [nm] (lg \epsilon)$
6a	457(1.98) <sup>a</sup>	6h	$468(3.51)^a$	6n	460(3.50) <sup>b</sup>
6b	$440^{b}$		$464(3.72)^{b}$ $461(3.55)^{d}$	6р	465(3.26) <sup>b</sup>
6d	412(2.60) <sup>b</sup>	6i	$464(3 42)^{b}$	6q	465(3.47) <sup>b</sup>
6e	423(2.19) <sup>c</sup>	6k	$464(3.64)^{b}$	6r	460 <sup>b,e</sup>
6f	425(2.31) <sup>c</sup>	61	$464(3.60)^{b}$	6s	453(3.27) <sup>b</sup>
6g	423(2.53) <sup>c</sup>	6m	463(3.65) <sup>b</sup>	6t	478(3.79) <sup>b</sup>
<i>a</i> in c	yclohexane; <sup>b</sup> in N	leCN; '	in <i>n</i> -hexane; <sup>d</sup> in	EtOH	<sup>e</sup> poorly soluble

increasing conjugative linking of the butadiene bridge. The UV absorption of the 6 (with two maxima at 230 - 250 and 310 - 350 nm) differs only slightly from those of the corresponding thiophenes 12.

Moreover, the (red) colour repeatedly caused doubts as to a cyclic disulfide structure corresponding to A and led us to re-examine the question of whether these molecules might exist as the open chain thioxo valence isomer form **B**, possibly in an equilibrium or in a no-bond resonance relationship (cf. ref.<sup>4a,9c,18a</sup>; concerning the influence of substituents on the stability of A and B, respectively: cf. ref.<sup>6h,j</sup>). It should be noted, however, that conjugated thiocarbonyl compounds, as in the case of **B**, usually exhibit their  $n \rightarrow \pi^*$  transitions at wave lengths substantially above 500 nm.

The cyclic disulfide structure **A** is unequivocally proved confirming our initial assertion (cf. ref.<sup>1c,d,21a</sup>). *Figure 1* illustrates the <sup>1</sup>H NMR 500 MHz spectrum of the parent compound **6a**, it is characterized by an AA'XX' type spectrum with chemical shifts exclusively in the olefinic region (cf. also ref.<sup>1f,h-i,k,l,n</sup>). In accord with this observation is the absence of any <sup>13</sup>C NMR indications for a thiocarbonyl group ( $\delta > 200$  ppm), as also stated in the spectra of the 3,6-disubstituted 1,2-dithiines. It should be emphasized that the ring-opened butadiene precursors **4** (as well as **10** and **11**) exist in the s-transoid conformation in solution, as concluded from a thorough NMR study.<sup>1f</sup> This conformation has been confirmed by X-ray crystallography.<sup>21b</sup>



The X-ray analysis of 3,6-di(morpholinomethyl)-1,2-dithiine  $(\mathbf{6g})^{22}$  and of 3,6-di(4-cyclohexylphenyl)-1,2dithiine  $(\mathbf{6p})^{23}$ , shown in *Figures 2* and 3, respectively, reveals a remarkable distortion of the 6-membered ring, as indicated by the C-S-S-C torsion angles of 52.6° (**6g**) and of 54.6° (**6p**), respectively. Further, by the twist of the C,C-double bonds against each other of 26.5° for **6g** and of 31.5° for **6p**, the repulsion of the adjacent S-p orbitals is minimized. The S-S distance of about 2 Å agrees with that of "normal" disulfides and excludes any consideration of a no-bond resonance with the ring-opened valence isomer of type **B**. These findings are consistent with that from our X-ray elucidations in the di-anellated 1,2-dithiine series<sup>24a</sup> and of a tetrasubstituted 1,2-dithiine<sup>24c</sup> (see finally informations in ref.<sup>24c</sup>). The ring inversion barrier of the cyclic disulfide should be very low, as previously demonstrated by the fact that the <sup>1</sup>H NMR spectrum is unaltered up to -95°C in the case of the di-(*R*)borneno anellated 1,2-dithiine.<sup>1i,n</sup> Finally it should be noted that the geometry of the cyclic system agrees well with the results of *ab-initio* calculations.<sup>6i,j</sup>



of 3,6-di(morpholinomethyl)-1,2-dithiine (8g). Selected data – *bond lengths*: S1-S1a = 2.0657(11), C1-S1 = 1.771, C1-C2 = 1.336(3), C2-C2a = 1.455(4) Å; *torsion angles*: C1a-S1a-S1-C1 = 52.6, S1a-S1-C1-C2 = 40.3, S1-C1-C2-C2a = 1.6, C1-C2-C2a-C1a = 26.5, C2-C2a-C1a-S1a = 1.6, C2a-C1a-S1a-S1 =  $40.3^{\circ}$ . Figure 3. X-ray crystal structure<sup>2,3</sup> of 3,6-di(4-cyclohexylphenyl)-1,2-dithine (**8p**). Selected data – *bond lengths*: S1-S2 = 2.503(5), C1-S1 = 1.780(10), C3-C4 = 1.337(15), C2-C3 = 1.420(20) Å; *torsion angles*: C1-S1-S2-C4 = -54.6), S2-S1-C1-C2 = 37.2, S1-S2-C4-C3 = 42.2, S1-C1-C2-C3 = 4.4, C1-C2-C3-C4 = -31.5, C2-C3-C4-S2  $\approx$  -1.3°.

Thus the predominance of the cyclic disulfide structure  $\mathbf{A} = \mathbf{6}$  versus its ring-opened valence isomer **B** seems comparable with the preference of the enthiols **10** versus the butane-1,4-dithiones for the same reasons. <sup>15b</sup> Alterations can be induced by  $\pi$ -donor groups in 3- and 6-position.<sup>6h,j,25</sup> A quite contrary situation exists in the oxygen series (cf. ref.<sup>10</sup>, and literature cited therein).

In accord with the rod-like shape of the 3,6-disubstituted 1,2-dithiines (cf. Figure 3), 3,6-di(4-*n*-butyl-phenyl)-1,2-dithiine (**6n**) displays liquid crystalline behaviour (nematic phase) on melting at 106°C for a short time prior to sulfur extrusion. By contrast the corresponding thiophene **12n** is inactive.

$E_{1/2}[V]$ -0.67 <sup>a</sup> -0.70 <sup>a</sup> -0.68 <sup>c</sup> -0.66 <sup>a</sup> -0.65 <sup>c</sup>	-0.63 <sup>a</sup>	-0.57 <sup>c</sup>	-0.73 <sup>a</sup>
-0.74 <sup>b</sup> -0.71 <sup>c</sup> -0.67 <sup>c</sup>	-0.62 <sup>c</sup>		-0.72 <sup>c</sup>

Table 2. Polarographic half wave potentials<sup>26a</sup>

The facile reduction of the 1,2-dithines 6 to the dimercapto-precursors 5 or 10 (see above) is in line with polarographic studies.<sup>26a</sup> The halfwave potentials appear, as listed in *Table 2*, in the range of about -0.5 to -0.7 V. These values accord with those of diaryl disulfides (-1.4 to -1.8 V)<sup>26b</sup> rather

than with those of dialkyl disulfides  $(-0.5 \text{ to } -0.9 \text{ V})^{26c}$ . The polarographic half wave potential of the known dibenzo-1,2-dithiine ("*o*-diphenylene disulfide")<sup>26d</sup> was found to be  $E_{1/2} = -0.56 \text{ V}$  (SCE, DMF, NEt<sub>4</sub>ClO<sub>4</sub>).

The olefinic character is demonstrated by the ready reaction of **6h** with bromine (*Scheme 7*). A tetrabromo adduct, stable only at low temperature and which decomposed to a compound of uncertain structure at room temperature, was produced. The structure assignment of the product as **21** is mainly based on the NMR indication of new sp<sup>3</sup>-C atoms [<sup>1</sup>H: CHBr 5.40 ppm; <sup>13</sup>C: CHBr 65.52, C(Br)Ph 81.77 ppm]. Moreover, no diastereomeric mixture could be observed. At -30°C the <sup>1</sup>H NMR spectrum (CDCl<sub>3</sub>) indicates a rotation hindrance of the phenyl groups. This result could be expected from steric overcrowding between the *o*-H atoms of the phenyl groups and the other atoms at the 1,2-dithiacyclohexane ring assuming that the phenyl groups

If a ring-opened valence tautomer B' behaves as a reactive species, even in an extremely low equilibrium concentration, the 1,2-dithiines 6 should be able to condense with hydrazine hydrate with the formation of pyridazines 22 according to path [i] of Scheme 8. The formation of 22, indeed takes place, e.g. in ethanol/pyridine along with the formation of thiophenes 12 in about equal amounts and hydrogen sulfide. Path [i], however, seems not to be involved but path [ii]: The formation of the pyridazines 22 proceeds more likely via reduction of 6 to the (Z,Z)-1,4-dimercaptobutadienes 5 or 10, respectively (as is known from non-cyclic disulfides)<sup>27</sup> followed by condensation to the dihydropyridazines K and final



oxidation. Reaction of 10 with hydrazine hydrate gives 22 in even better yields. The additional formation of the thiophenes 12 in the reaction of 6 with hydrazine hydrate possibly results from a competing base-catalyzed sulfur extrusion (cf. below).

More detailed information is provided by polarographic studies in dimethylformamide (SCE, NEt4I). Under air exclusion the mixture of **6h** and hydrazine hydrate shows a polarographic half wave potential at  $E_{1/2} = -1.95$  V, corresponding to the reduction of **K** ( $R^1 = R^2 = C_6H_5$ ). After admittance of air the latter half wave potential vanishes and that indicative of the reduction of **22** ( $R^1 = R^2 = C_6H_5$ ) at  $E_{1/2} = -1.51$  V gradually appears. The latter value should correspond to attack at the "azo subunit" whereas the reduction of (*E,E*)-1,4-diphenylbutadiene appears under the same conditions at  $E_{1/2} = -2.15$  V (cf. half-wave potential of the S-S fission of **6h** in table 2). Analogous results were obtained using 1,2-dibenzoylethane, the oxygen counterpart of **10h**. On the other hand, (*Z*)-1,2-dibenzoylethene, the oxygen analogue of **B'** ( $R^1 = R^2 = C_6H_5$ ), reacts immediately with hydrazine hydrate under the same conditions to produce **22** ( $R^1 = R^2 = C_6H_5$ ), indicated as well by  $E_{1/2} = -1.51$  V.

This reaction represents rather a special instance in the transformation possibilities of the 1,2-dithiine ring. Normally the reaction potential of the 3,6-disubstituted 1,2-dithiines **6** is dominated by the propensity to extrude sulfur producing thiophenes. For example all attempts to use **6h** as diene or dienophile in Diels-Alder reactions or in addition reactions with nucleophiles (e.g. amines) failed due to the predominance of sulfur extrusion. The 3,6-disubstituted 1,2-dithiines **6** undergo this stabilization process smoothly in the solid state on heating near the melting point or in solution above 100°C. Sulfur extrusion from **6** in solution at room temperature under normal day-light is also an unusual feature of the chemistry of these compounds. Additionally, it should be emphasized that the parent compound **6a** undergoes polymerisation more readily, especially under base catalysis (possibly via ring-opening by S-S fission), whilst di-anellated 1,2-dithiines are surprisingly photochemically stable and require forcing thermal conditions for sulfur extrusion.  $l_{i,k,n}$  Various more detailed results, obtained from our preliminary studies on the sulfur extrusion, are outlined below.

By monitoring the thermal sulfur extrusion process by <sup>1</sup>H NMR, it was possible to deduce that the reaction kinetics were (pseudo)first order.<sup>28</sup> The process is significantly influenced by the substituents, as demonstrated by some half-life values determined in dimethylsulfoxide-d<sub>6</sub>: For **6f** t<sub>1/2</sub> = 6219 s (120°C), for **6g** t<sub>1/2</sub> = 2810 s (120°C), and for **6h** t<sub>1/2</sub> = 13562 (90°C), 3338 (105°C), 525 (120°C), and 203 s (135°C). The reaction

rate is significantly increased by additional conjugation, as exemplified by **6h**. From the temperature dependence of the extrusion of **6h** the values  $\Delta H^{\ddagger} = 115.7 \text{ kJ} \cdot \text{mol}^{-1}$  and  $\Delta S^{\ddagger} = -10.2 \text{ J} \cdot \text{mol}^{-1} \cdot \text{K}^{-1}$  were deduced, indicating only a slight enhancement of order in the transition phase.



**Figure 4.** Correlation between the quantum yield  $\Phi$  of the photolysis and the UV/Vis spectrum of **6h** in ethanol

On the other hand, the light-induced sulfur extrusion depends significantly on the wavelength  $\lambda_{irr.}$  of the irradiated light and is unequivocally characterized by the quantum yields  $\Phi$  as illustrated in *Figure 4* for the case of **6h** [determined at 20°C and based on the decrease of the extinction of the long wave absorption].<sup>29a</sup> Absorption in the visible region is chiefly responsible for this process. For example, the quantum yields  $\Phi$  (in ethanol) are for **6d**: 0.56 ( $\lambda_{irr.} = 365$  nm), 0.77 ( $\lambda_{irr.} = 436$  nm); for **6f**: 0.14 ( $\lambda_{irr.} = 405$  nm), 0.21 ( $\lambda_{irr.} = 436$  nm), 0.14 ( $\lambda_{irr.} = 525$  nm); for **6g**: 0.30 ( $\lambda_{irr.} = 405$  nm), 0.47 ( $\lambda_{irr.} = 254$  nm), 0.33 ( $\lambda_{irr.} = 365$  nm), <sup>29b</sup> 0.63 ( $\lambda_{irr.} = 405$  nm), 0.73 ( $\lambda_{irr.} = 436$  nm), 0.68 ( $\lambda_{irr.} = 525$  nm). These results are

in line with an ab initio study.<sup>6i,j</sup> According to this study the determined  $n \rightarrow \sigma^*$  character of the longest wave transition should weaken the S-S bond and, hence, favour ring-opening, whilst the  $\pi \rightarrow \pi^*$  transitions in the short wave range should be less able to weaken the S-S bond and should be characterized, consequently, by smaller quantum yields. The influence of the substituents seems to be comparable with those in thermolysis (see above). Additionally, the choice of reaction solvent (e.g. heptane versus ethanol) had no significant influence on the quantum yield. Oxygen, known to be a good quencher of triplet states, did not influence the quantum yields (almost identical values under argon as under oxygen). Hence, the extrusion of sulfur appears to be due to an electronically excited singlet state.



A rationalization of the sulfur extrusion<sup>30a</sup> is shown in Scheme 9. On the basis of well documented valence isomerizations in the heterocyclohexa-1,3-diene series<sup>31</sup> the ring-opened valence isomer **B'** of 6 (analogous **B**) undergoes, in the twisted form **B''**, an intramolecular  $[\pi 4+\pi 2]$ cycloaddition to the episulfide  $L^{30b}$  which then extrudes sulfur affording the thiophenes 12 in an irreversible final step (*pathway i*). This stabilization process parallels the sulfur extrusion in thiepine<sup>32</sup> via thianorcaradien and 1,4dithiocine<sup>33</sup> via benzenebisepisulfide.<sup>34</sup> Moreover, this sequence explains the amazing resistance of the di-anellated 1,2-dithiines to extrude sulfur under the above noted conditions due to the steric hindrance in **B''**. An intermediate ring opened thioxo isomer **B'**, however, could not be

detected by flash photolysis in the microsecond range at room temperature. An alternative assumption according to *pathway ii* avoids an intermediate ring opening to **B'** but involves a 1,5-electrocyclization  $M \rightarrow N^{35}$  proceeding via the thiosulfoxide **O** (ref.<sup>30</sup>, p. 172) as possibly competing precursor for 12.

The final process from L or O, respectively, and the nature of the primarily extruded sulfur are still unclear. In preliminary attempts with **6h** to trap  $S_n$  (n = 1, 2) with excess cyclohexene (1:300 molar ratio) and with 2,3-dimethylbutadiene (1:2 molar ratio, benzene) in day-light at 20°C, a S<sub>1</sub> or S<sub>2</sub>-adduct could be detected in traces only ( $\leq 1\%$ ; GC-MS). The major products were thiophene **12h** and oligomeric sulfur (mainly S<sub>8</sub>). Moreover, in the <sup>1</sup>H NMR monitoring of an attempted reaction of **6h** with norbornene in DMSO-d<sub>6</sub> at 120°C (with exclusion of day-light) only the gradual formation of **12h** was observed. Thus, the attractive possibility that 1,2-dithines could function as sulfur transfer reagents remains questionable. The primary extrusion or transfer of singlet sulfur should require too much energy as suggested in ref.<sup>36</sup>. On the other hand, a sulfur transfer via addition of the supposed intermediates L or O, respectively, to the olefinic substrates is obviously minimized due to reactions of these intermediates with themselves. Thus, a successive concatenation of sulfur atoms and their final elimination in a thermodynamically favoured cyclic form according to the sequence (a)-(c) in *Scheme 10* (just as proposed for the sulfur extrusion from thiepine to benzene via thianorcaradiene)<sup>37</sup> should compete successfully.<sup>38</sup>



#### Conclusions

The title system A represents formally an  $8\pi$ -electron system, a sulfur-expanded thiophene and a butadiene-bridged disulfide. Special attention has been focussed on the parent compound and the 3,6-disubstituted derivatives 6, which as *non-anellated* compounds differ essentially from their anellated counterparts in some fundamental aspects of their synthesis and behaviour. Unusual features are represented by the light absorption in the visible region, by the structure with respect to the questionable existence of the valence isomer **B** and by the tendency for the sulfur extrusion both under thermal and photochemical conditions.

A general synthesis, based on the base-assisted nucleophilic bis-addition of benzyl- or t-butylmercaptans, respectively, to butadiynes 1 forming the corresponding (Z,Z)-1,4-dithiobutadienes 3, subsequent deprotection to (Z,Z)-butadiene-1,4-dithiols 10 and finally disulfide oxidation, was developed. An alternative approach, exemplified by the lithium-halogen exchange at (Z,Z)-1,4-diphenylbutadiene (19) and successive sulfurization, was limited by the accessibility of the precursor.

The colour, arising from a  $n \rightarrow \sigma^*$  transition, is influenced by the substituents and bathochromic shifts are especially caused by conjugation effects. The structure, a twisted cyclic disulfide, is well established in solution by NMR spectroscopy and in the solid state by X-ray analysis. The presence of any ring-opened thioxo valence isomer **B** is unequivocally excluded, unlike the situation in the 1,2-dithiete series. The structure **B**, however, can be involved as a transient species in thermally as well as photochemically induced sulfur extrusion with formation of the correspondingly substituted thiophenes. The quantum yields of the photo-induced sulfur extrusion depend crucially on the wave length of the light emloyed and attain their maximum value at about the absorption maximum in the visible region. A significant dependence on the substituents is also demonstrated from a consideration of the half-life values in the thermally induced process. Based on the activation parameters, a low ordered transition state is suggested. The extrusion process appears to involve a complex sequence of events prior to the expulsion of oligomeric sulfur. The use of these compounds as preparatively useful sulfur transfer reagents is excluded b at this time.

The high tendency to extrude sulfur under mild conditions predominates over the general reaction behaviour of the 1,2-dithine ring. The 1,2-dithine ring behaves preponderantly as an olefinic cyclic disulfide, and the compounds readily undergo reductive S-S ring fission. In the reaction with hydrazine hydrate, a primary reduction to (Z,Z)-butadiene-1,4-dithiols occurs which subsequently condense with excessive hydrazine to afford the pyridazines 22. Bromine can be added to the C=C double bonds to form an unstable tetrabromo adduct.

#### **Experimental Part**

NMR spectra: Varian Unity 500 (<sup>1</sup>H: 499.84 MHz, <sup>13</sup>C: 125.71 MHz), Bruker WP 200 (<sup>1</sup>H: 200.13 MHz, <sup>13</sup>C: 50.3327 MHz), Bruker AC 80 (<sup>1</sup>H: 80.13 MHz, <sup>13</sup>C: 20.149 MHz). <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded with TMS as internal standard. – MS: Varian MAT CH6, AMD Intectra 402 (70 eV). – IR: Carl Zeiss Jena Specord 71 and 75. – UV: Beckman DK-2A; Perkin-Elmer Lambda 16 – Column chromatography (CC): silica gel 60 [70-230 and 230-400 mesh (Merck)]. – HPLC: Merck Hitachi L-4000 (UV detector). – Melting points: Heating stage microscope (Boetius M; all temperatures quoted are not corrected). – X-ray analyses: Diffractometer STADI4 (Stoe, MoK<sub>α</sub> radiation,  $3 < 2\Theta < 54^{\circ}$ ). – Elemental analyses: Carlo Erba (automatic apparatus).

#### Nucleophilic addition of thiols 2 and 7 at butadiynes 1

(Z,Z)-1,4-Di(benzylthio)-1,3-butadiene (4a; cf. ref.<sup>1a,c,f</sup>). – *Procedure I*: A mixture of 252 g (2 mol) benzyl chloride, 152 g (2 mol) thiourea and 400 ml ethanol was heated at reflux with stirring 6 h, then treated with 320 g (8 mol) NaOH

and 2 l ethanol and the resulting mixture heated at reflux for further 3 h (generation of 2) From a separate generator and under argon, a stream of 1a (prepared by the dropwise addition of 600 g of a 40% aqueous NaOH solution to 200 (1.45 mol) 1,4-dichlorobutyne in 300 ml ethanol) was slowly (2-3 h) bubbled into the above solution of 2. After 3-4 h the mixture was diluted with two volumes of water and the separated solid filtered and recrystallized from EtOH/AcOEt (1:1). ~ Colourless leaflets; yield 200-300 g (66-78%); m.p. 128-129°C.  $^{-1}$ H NMR (CDCl<sub>3</sub>):  $\delta = 3.88$  (s; 4H; 2 SCH<sub>2</sub>), 6.00 (2 d; 2H; =CH-SCH<sub>2</sub>); 6.29 (2 d; 2H; =CH-CH=; J<sup>3</sup> = 5.8, J<sup>4</sup> = 1.9 Hz), 7.28 (m; 10H; aromat. H); cf. data in ref.<sup>1f.</sup> - <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 38.2$  (-SCH<sub>2</sub>-), 123.2 (=CH-SCH<sub>2</sub>), 126.9, 127.2 (=CH-, C<sub>p-aromat.</sub>), 128.6, 128.8 (C<sub>o,m-aro</sub>-

mat.), 137.67 ( $C_{i-aromat.}$ ); cf. data in ref.<sup>11</sup>. – MS (70 eV) m/z = 298 (8 [M]<sup>+</sup>), 207 (18 [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>]), 175 (6 [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>] – S]), 149 (17 [M/2]<sup>+</sup>). – C<sub>18</sub>H<sub>18</sub>S<sub>2</sub> (298.5): calcd. C 72.44, H 6.08, S 21.48; found: C 72.54, H 5.94, S 21.40. – *Procedure II* (*in situ* reaction of 1a): Analogous to procedure I using a 1 mole charge in 200 ml ethanol with stirring and under an inert atmosphere. After heating under reflux for 6 h, 280 g KOH was added, the mixture warmed until the commencement of reflux and 62 g (0.5 mol) 1,4-dichlorobutyne were added over a 1 h period. The mixture was then processed as above. 45-60 g (30-40%) yield.

 $(Z_zZ)$ -5,8-Di(benzylthio)dodeca-5,7-diene (4b). – A solution of 5.6 g (100 mmol) KOH and 0.264 g (1 mmol) 18crown-6 in 15 ml water was stirred 15 min under argon. Subsequently 1.5 g (12 mmol) 2 in 15 ml benzene and 0.81 g (5 mmol) 1b in 20 ml benzene were added with stirring. The mixture was refluxed 20 h. After separation of the layers the aqueous phase was extracted with benzene. The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The residue was purified by CC (silica gel, benzene). – Pale yellow oil, yield: 1.9 g (93%). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.91-0.95 (t, 6 H, CH<sub>3</sub>), 1.38-1.53 (m, 8 H, CH<sub>2</sub>), 2.22-2.29 (t, 4 H, CH<sub>2</sub>), 3.60 (s, 4 H, SCH<sub>2</sub>), 7.21-7.36 (m, 12 H, C<sub>6</sub>H<sub>5</sub>, CH=). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 13.5 (CH<sub>3</sub>), 18.9, 21.9, 30.4 (CH<sub>2</sub>), 37.0 (SCH<sub>2</sub>), 127.4, 128.5, 128.9, 129.4, 136.8, 137.5 (C<sub>6</sub>H<sub>5</sub>, =CH-, >C=). – MS (70 eV) m/z = 184 (10) [M<sup>+</sup> – 2 C<sub>7</sub>H<sub>7</sub> – 2 CH<sub>3</sub>CH<sub>2</sub>], 162 (100) [M<sup>+</sup> – 2 C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>SH – C<sub>3</sub>H<sub>7</sub>], 105 (90) [M<sup>+</sup> – 2 C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>SH – CH<sub>3</sub>], 133 (24) [M<sup>+</sup> – 2 C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>SH – C<sub>2</sub>H<sub>5</sub>], 119 (62) [M<sup>+</sup> – 2 C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>SH – C<sub>3</sub>H<sub>7</sub>], 105 (90) [M<sup>+</sup> – 2 C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>SH – C<sub>4</sub>H<sub>9</sub>]. – C<sub>26</sub>H<sub>34</sub>S<sub>2</sub> (410.7): calcd. C 76.04, H 8.35. S 15.61; found: C 76.29, H 8.06, S 15.62.

(Z)-3-Benzylthio-2,2,7,7-tetramethyl-3-octen-5-yne (3c). – A mixture of 2.56 ml (20 mmol) 2, 300 mg powdered KOH, 1.62 g (10 mmol) 1c and 40 ml DMF was stirred 2 h (solution). After 24 h the mixture was poured on to ice and the product was washed with water and the residue was recrystallized from ethanol. – Colourless needles; yield: 35%; m.p. 59-60°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.23$  (s, 9 H, *t*-Bu), 1.35 (s, 9 H, *t*-Bu), 3.48 (s, 2 H, SCH<sub>2</sub>), 6.48 (s, 1 H, =CH-), 7.26-7.32 (m, 5 H, C<sub>6</sub>H<sub>5</sub>). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 29.1$ , 30.8 [C(CH<sub>3</sub>)<sub>3</sub>], 28.4, 38.6 [C(CH<sub>3</sub>)<sub>3</sub>], 37.9 (SCH<sub>2</sub>), 77.3 ( $\equiv$ C-CH=), 99.8 ( $\equiv$ C-*t*-Bu), 104.3 (=CH-), 127.4 (C<sub>0-aromat</sub>), 128.6 (C<sub>m-aromat</sub>), 129.1 (C<sub>p-aromat</sub>), 135.9 (C<sub>i-aromat</sub>), 159.0 (=C-S). – MS (70 eV) m/z = 286 (55) [M<sup>+</sup>], 271 (21) [M<sup>+</sup> – CH<sub>3</sub>], 229 (39) [M<sup>+</sup> – C<sub>4</sub>H<sub>9</sub>], 215 (11) [M<sup>+</sup> – C<sub>4</sub>H<sub>9</sub> – CH<sub>2</sub>], 173 (8) [M<sup>+</sup> – C<sub>4</sub>H<sub>9</sub> – CH<sub>2</sub> – C<sub>3</sub>H<sub>6</sub>], 91 (100) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>], 57 (55) [C(CH<sub>3</sub>)<sub>3</sub><sup>+</sup>]. – C<sub>19</sub>H<sub>26</sub>S (286.5): calcd. C 79.66, H 9.15, S 11.19; found: C 79.73, H 9.02, S 11.24.

 $(Z_zZ)$ -3,6-Di(benzylthio)-2,7-dihydroxy-2,7-dimethyl-3,5-octadiene (4d). – A mixture of 2.9 ml (24.4 mmol) 2, 332 mg KOH, 1.85 g (11.1 mmol) 1d and 40 ml DMF was stirred 180 min at room temperature. The mixture was diluted with 150 ml water and extracted twice with AcOEt. The combined organic phases were washed with water and the residue obtained on concentration was recrystallized from *n*-hexane. – Colourless needles; yield 3.45 g (75 %); m.p. 95-96°C. – IR (KBr):  $\bar{v} = 3500$ -3140 cm<sup>-1</sup> (s, OH, free and assoc.) – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.36$  (s, 12 H, CH<sub>3</sub>), 2.16 (s, 2 H, OH), 3.81 (s, 4 H, SCH<sub>2</sub>), 7.08 (s, 2 H, =CH-), 7.21-7.28 (m, 10 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 29.4$  (CH<sub>3</sub>), 41.6 (SCH<sub>2</sub>), 74.7 [C(CH<sub>3</sub>)<sub>3</sub>OH], 127.3, 128.6, 129.1, 131.5, 140.0, 147.2 (C<sub>olef</sub>, C<sub>aromat</sub>). – MS (70 eV) m/z = 414 (1) [M<sup>+</sup>], 305 (2) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – H<sub>2</sub>O], 265 (19) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – H<sub>2</sub>O – C<sub>3</sub>H<sub>4</sub>], 247 (78.6) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – 2 H<sub>2</sub>O – C<sub>3</sub>H<sub>4</sub>], 207 (16) [M/2<sup>+</sup>], 91 (100) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>24</sub>H<sub>30</sub>S<sub>2</sub>O<sub>2</sub> (414.5): calcd. C 69.56, H 7.25 S 15.46; found C 69.45, H 7.45, S 15.45.

 $(Z_zZ)$ -2,5-Di(benzylthio)-1,6-dipyrrolidino-2,4-hexadiene (4e). – To a mixture of 2.56 ml (22 mmol) 2, 56 mg powdered KOH and 50 ml dry diglyme 2.17 g (10 mmol) 1e (m.p. 48-49°C) were added with stirring over 30 min. Stirring was continued 3 h at room temperature. The mixture was poured into 150-200 ml ice-water and then extracted twofold with AcOEt and once with Et<sub>2</sub>O. After washing the combined organic layers with water (NaCl) and evaporation, the residue was recrystallized twice from ethanol. – Light beige needles; yield: 1.86 g (40%); m.p. 113-114°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.46$  (m, 8 H, N-CH<sub>2[cycl.]</sub>), 2.48 (m, 8 H, CH<sub>2[cycl.]</sub>), 3.22 (s, 4 H, N-CH<sub>2</sub>), 4.03 (s, 4 H, S-CH<sub>2</sub>), 7.16-7.29 (m, 10 H, aromat.). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 23.7$  ([CH<sub>2]2</sub>), 36.5 (S-CH<sub>2</sub>), 53.8 (N-CH<sub>2[cycl.]</sub>), 62.7 (N-CH<sub>2</sub>), 126.9 (=CH-), 128.2 (>C=), 128.4 (C<sub>0-aromat.</sub>), 128.8 (C<sub>m-aromat.</sub>), 136.1 (C<sub>p-aromat.</sub>), 138.5 (C<sub>i-aromat.</sub>). – MS (70 eV) m/z = 464 (0.3) [M<sup>+</sup>], 373 (27) [M<sup>+</sup> – CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 302 (50) [M<sup>+</sup> – CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> – HN(CH<sub>2</sub>)<sub>4</sub>], 270 (8) [M<sup>+</sup> – CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> – HN(CH<sub>2</sub>)<sub>4</sub> – S], 244 (6) [M<sup>+</sup> – CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> – HN(CH<sub>2</sub>)<sub>4</sub> – S – C<sub>2</sub>H<sub>5</sub>], 91 (53) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>], 84 (100) [pyrrolidino-CH<sub>2</sub><sup>+</sup>]. – C<sub>28</sub>H<sub>36</sub>N<sub>2</sub>S<sub>2</sub> (464.6): calcd. C 72.39, H 7.81, N 6.03, S 13.80; found C 72.22, H 7.84, N 6.13, S 13.82.

 $(Z_zZ)$ -2,5-Di(benzylthio)-1,6-dipiperidino-2,4-hexadiene (4f). – As described for 4e with 2.44 g (10 mmol) 1f (m.p. 60°C). – Light beige needles; yield: 2.66 g (54%); m.p. 90-91°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.42-1.47 (m, 4 H, CH<sub>2[cycl.C4]</sub>), 1.57-1.62 (m, 8 H, CH<sub>2[cycl.C3/5]</sub>), 2.35-2.37 (m, 8 H, CH<sub>2[cycl.C2/6]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 2.35-2.37 (m, 8 H, CH<sub>2[cycl.C2/6]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 2.35-2.37 (m, 8 H, CH<sub>2[cycl.C2/6]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 2.35-2.37 (m, 8 H, CH<sub>2[cycl.C3/6]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, CH<sub>2[cycl.C3/5]</sub>), 3.09 (s, 4 H, N-CH<sub>2</sub>), 4.12 (s, 4 H, N-CH<sub>2[cycl.C3/5]</sub>), 3.09 (s

S-CH<sub>2</sub>), 6.73 (s, 2 H, =CH-), 7.19-7.33 (m, 10 H, aromat. H).  $-^{13}$ C NMR (CDCl<sub>3</sub>):  $\delta = 24.6$  (CH<sub>2[cycl.C4]</sub>), 26.2 (CH<sub>2[cycl.C3/5]</sub>), 36.3 (S-CH<sub>2</sub>), 54.4 (N-CH<sub>2[cycl.1</sub>), 126.8(3) (=CH-), 127.8(6) (>C=), 128.4 (C<sub>o-aromat.</sub>), 128.8 (C<sub>m-aromat.</sub>), 135.5 (C<sub>p-aromat.</sub>), 138.6 (C<sub>i-aromat.</sub>). - MS (70 eV) m/z = 492 (1) [M<sup>+</sup>], 401 (22) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 316 (48) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - piperidine], 284 [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - piperidine - S], 258 (4) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - piperidine - S - C<sub>2</sub>H<sub>2</sub>], 233 (6) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - piperidine - S - 2 C<sub>2</sub>H<sub>2</sub>], 98 (100) [piperidino-CH<sub>2</sub><sup>+</sup>]. - C<sub>30</sub>H<sub>40</sub>N<sub>2</sub>S<sub>2</sub> (492.8): calcd. C 73.14, H 8.18, N 5.69, S 13.02; found C 72.95, H 7.89, N 5.59, S 13.05.

 $(\textbf{Z,Z})-2,5-Di(benzylthio)-1,6-dimorpholino-2,4-hexadiene (4g). - As described for 4e with 2.48 g (10 mmol) 1g (m.p. 107-108°C). - Light beige needles; yield: 2.08 g (42%); m.p. 116-117°C. - <sup>1</sup>H NMR (CDCl<sub>3</sub>): <math>\delta = 2.39-2.41$  (t, 8 H, N-CH<sub>2[cycl.]</sub>), 3.10 (s, 4 H, N-CH<sub>2</sub>), 3.68-3.70 (t, 8 H, O-CH<sub>2[cycl.]</sub>), 4.07 (s, 4 H, S-CH<sub>2</sub>), 6.72 (s, 2 H, =CH-), 7.18-7.30 (m, 10 H, aromat. H). - <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 36.1$  (S-CH<sub>2</sub>), 53.4 (N-CH<sub>2[cycl.]</sub>). 65.1 (N-CH<sub>2</sub>), 67.1 (O-CH<sub>2[cycl.]</sub>), 127.1 (=CH-), 128.0 (>C=), 128.5 (C<sub>0-aromat.</sub>), 128.7 (C<sub>m-aromat.</sub>), 134.7 (C<sub>p-aromat.</sub>), 138.2 (C<sub>i-aromat.</sub>). - MS (70 eV) m/z = 496 (0.4) [M<sup>+</sup>], 405 (20) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 318 (57) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - C<sub>4</sub>H<sub>9</sub>NO], 286 (9) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - C<sub>4</sub>H<sub>9</sub>NO - S], 260 (3) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - C<sub>4</sub>H<sub>9</sub>NO - S - C<sub>2</sub>H<sub>2</sub>], 234 (8) [M<sup>+</sup> - CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - C<sub>4</sub>H<sub>9</sub>NO - S - 2 C<sub>2</sub>H<sub>2</sub>], 100 (100) [morpholino-CH<sub>2</sub><sup>+</sup>]. - C<sub>28</sub>H<sub>36</sub>N<sub>2</sub>O<sub>2</sub>S<sub>2</sub> (496.7): calcd. C67.72, H 7.31, N 5.64, S 12.91; found C 67.62, H 7.40, N 5.73, S 13.04.

(Z,Z)-1,4-Di(benzylthio)-1,4-diphenyl-1,3-butadiene (4h; cf. ref.<sup>1c,f</sup>). – *Procedure I*: After refluxing (1 h) a mixture of 30 g (0.15 mol) benzylisothiouronium chloride and 30 g (0.75 mol) NaOH in 160 ml ethanol, 10.1 g (0.05 mol) 1h was gradually added (30 min) under argon and the mixture heated under reflux for a further 6 h. At 20°C 500 ml water was added; the isolated solid was crystallized from EtOH/AcOEt – Pale yellow needles; yield: 12.5 g (55%); m.p. 117-118°C. – <sup>1</sup>H NMR (CDCI<sub>3</sub>):  $\delta$  = 3.61 (s; 4H; S-CH<sub>2</sub>), 7.04 (m; 4H; aromat. H), 7.13 (s; 2H; =C-H), 7.18 (m; 6H; aromat. H), 7.35 (m; 6H; aromat. H), 7.50 (m; 4H; aromat. H); cf. data in ref.<sup>1f. 13</sup>C NMR (CDCI<sub>3</sub>):  $\delta$  = 37.7 (S-CH<sub>2</sub>), 126.9 (CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>[p]), 128.0(8) (C<sub>6</sub>H<sub>5</sub>[p)), 128.1(3), 128.2, 128.4, 128.7 (C<sub>6</sub>H<sub>5</sub>[o,m]), 132.2(7) (=CH-), 138.3(2), 139.2, 140.0 (>C=); cf. data in ref.<sup>1f.</sup> – MS (70 eV): m/z = 450 (13) [M]<sup>+</sup>, 359 (80) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 237 (27), 236 (94) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 91 (100) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>30</sub>H<sub>26</sub>S<sub>2</sub> (450.7): calcd. C 79.95, H 5.82, S 14.23; found C 79.43, H 5.60, S 14.47. – *Procedure II*: To a mixture of 56 mg KOH or 112 mg KO-tBu, respectively, and 50 ml diglyme under argon 2.56 ml (22 mmol) **2** was added drowise and, after stirring for 30 min at 20°C, the mixture was treated slowly with 2.02 g (10 mmol) **1b**. After further stirring at 20°C (2.5 h) the red coloured mixture was poured on ice-water, the separated yellow solid was extracted with AcOEt, and the organic layer evaporated i. vac.: 58% (using KOH), 61% (using KO-tBu).

(Z,Z)-1,4-Di(benzylthio)-1,4-di-(4-bromophenyl)-1,3-butadiene (4i). – As described for 4h/procedure I using 1i (m.p. 269°C): Colourless prisms; yield: 56%; m.p. 191-192°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 3.58$  (s, 4 H, CH<sub>2</sub>), 6.96-7.50 (m, 20 H, aromat. H), 7.05 (s, 2 H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 37.7$  (CH<sub>2</sub>), 122.2 (C<sub>i</sub> [C<sub>6</sub>H<sub>4</sub>Br], 127.0 (=CH-), 128.3, 128.7. 129.6, 131.6 (aromat. C), 132.9 (=CH-), 138.1, 138.4, 139.9 (>C=). – MS (70 eV): m/z = 608 (8), 517 (62) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 394 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>]. – C<sub>30</sub>H<sub>24</sub>Br<sub>2</sub>S<sub>2</sub> (608.4): calcd. C 59.22, H 3.98, Br 26.26, S 10.54; found C 59.07, H 3.93, Br 26.38, S 10.22.

(Z,Z)-1,4-Di(benzylthio)-1-phenyl-4-(p-tolyl)-1,3-butadiene (4j; cf. ref.<sup>1d</sup>). – As described for 4h/procedure I using 1j (m.p. 110°C): Pale yellow needles; yield: 55%; m.p. 120-121°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.38 (s, 3 H, CH<sub>3</sub>), 3.60 (s, 4 H, S-CH<sub>2</sub>), 7.10, 7.11 (s, s, 2 H, =CH-), 7.02 (m, 4 H, aromat. H), 7.18 (m, 7 H, aromat. H), 7.41 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 21.2 (CH<sub>3</sub>), 37.7 (S-CH<sub>2</sub>), 126.8 (CH<sub>2</sub>C<sub>6</sub>H<sub>5[p]</sub>), 128.0 (C<sub>6</sub>H<sub>5[p]</sub>), 128.0, 128.1, 128.2, 128.4, 128.8, 129.1 (aromat. C), 131.7 (H-C<sup>3</sup>=), 132.5 ((H-C<sup>1</sup><sup>2</sup>=), 137.2, 138.1, 138.3(9), 138.4(4), 138.6, 139.3, 140.1 (>C=). – MS (70 eV): m/z = 464 (6) [M<sup>+</sup>], 373 (56) [M<sup>+</sup> – C<sub>7</sub>C<sub>7</sub>], 296 (46) [M<sup>+</sup> – C<sub>7</sub>C<sub>7</sub> – C<sub>6</sub>H<sub>5</sub>], 281 (31) [M<sup>+</sup> – C<sub>7</sub>C<sub>7</sub> – C<sub>6</sub>H<sub>5</sub>], 91 (88) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>31</sub>H<sub>28</sub>S<sub>2</sub> (464.7): calcd. C 80.12, H 6.08; found C 79.87, H 5.99.

(Z,Z)-1,4-Di(benzylthio)-1,4-di(p-tolyl)-1,3-butadiene (4k; cf. ref.<sup>1c</sup>). – As described for 4h/procedure I with 11.5 g (0.05 mol) 1k (m.p. 183°C) resulting in 6.5 g (27%) product. The evaporation of the mother liquor gave 14.35 g of an oily product (mono-adduct 3k) which was subsequently treated with 5.46 g (0.044 mol) 2 and 600 mg KOH in 120 ml DMF under stirring 2 h at 46°C. After addition of water further 7.8 g (40.5%) product was isolated and recrystallized from EtOH/AcOEt (1:1): Pale yellow needles; yield (total): 14.3 g (59.7%); m.p. 135-136°C.  $^{-1}$ H NMR (CDCl<sub>3</sub>):  $\delta = 2.38$  (s, 6 H, CH<sub>3</sub>), 3.61 (s, 4 H, S-CH<sub>2</sub>), 7.03 (m, 4 H, CH<sub>2</sub>C<sub>6</sub>H<sub>5[o]</sub>), 7.11 (s, 2 H, =CH-), 7.22 (m, 10 H, aromat. H), 7.41 (m, 4 H, aromat. H).  $^{-13}$ C NMR (CDCl<sub>3</sub>):  $\delta = 21.2$  (CH<sub>3</sub>), 37.7 (S-CH<sub>2</sub>), 126.8 (aromat. C<sub>[p]</sub>), 128.0, 128.2, 128.8, 129,1 (aromat. C<sub>[0,m]</sub>), 131.8 (=CH-), 137.2, 138.0, 138.5, 138.7 (>C=). – MS (70 eV): m/z = 478 (9) [M<sup>+</sup>], 387 (55) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub>], 264 (100) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 91 (41) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>].  $^{-}C_{32}H_{32}S_{2}$  (478.7): calcd. C 80.29, H 6.32; found C 79.93, H 6.42. – From the oily intermediate the mono-adduct [(Z)-1-benzylthio-1,4-di(p-tolyl)-1-buten-3-yne (3k; cf. ref.<sup>1c</sup>)] could be isolated by crystallization from ethanol: Colourless needles, m.p. 90-91°C; MS (70 eV): m/z = 354 (12) [M<sup>+</sup>]; C<sub>25</sub>H<sub>22</sub>S (354.5): calcd. C 84.70, H 6.26; found 84.28, 6.36.

 $(Z_zZ)$ -1,4-Di(benzylthio)-1-(p-methoxyphenyl)-4-phenyl-1,3-butadiene (4l; cf. ref.<sup>1d</sup>). – As described for 4h/procedure I using 1l m.p. (110°C): Pale yellow needles; yield: 55%; m.p. 108°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.61 (s, 4 H, S-

#### W. SCHROTH et al.

CH<sub>2</sub>), 3.85 (s, 3 H, CH<sub>3</sub>), 6.91 (m, 2 H, aromat. H), 7.04 (d, 1H, J = 6.56 Hz,  $-C^2H=$ ), 7.05 (m, 4 H, aromat. H), 7.11 (d, 1H, J = 6.56 Hz,  $-C^3H=$ ), 7.20 (m, 6 H, aromat. H), 7.30-7.54 (m, 7 H, aromat. H).  $-^{13}C$  NMR (CDCl<sub>3</sub>):  $\delta = 37.6(5)$ , 37.6(9) (S-CH<sub>2</sub>), 55.4 (O-CH<sub>3</sub>), 113.8, 126.8, 127.9(4), 128.0(7), 128.2(1), 128.2(5), 128.4, 128.7(4), 128.7(5), 129.3 (aromat. CH), 131.0 (=CH-), 132.5 (C<sub>1</sub>[C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]), 132.6 (=CH-), 138.2, 138.4(1), 138.4(4), 139.0, 140.2 (>C=), 159.8 (C-OCH<sub>3</sub>). – MS (70 eV): m/z = 480 (9) [M<sup>+</sup>], 389 (74) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 267 (38), 266 (99) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> – CH<sub>3</sub>], 91 (100) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>31</sub>H<sub>28</sub>OS<sub>2</sub> (480.7): calcd. C 77.46, H 5.90; found C 76.92, H 6.12.

(Z,Z)-1,4-Di(benzylthio)-1,4-di(p-methoxyphenyl)-1,3-butadiene (4m; cf. ref.<sup>1c</sup>). – A mixture of 5.24 g (0.02 mol) 1m (m.p. 140°C), 5.1 g (0.041 mol) 2, 0.5 g (1.9 mmol) 18-crown-6 (1.9 mmol), 1.12 g (0.02 mol) KOH, 30 mł water and 60 mł benzene was refluxed for 5 h with stirring. The oily evaporation residue obtained from the organic layer was again treated with 3.6 g (0.029 mol) 2 and 0.56 g (0.01 mol) KOH in 60 ml DMF and stirred for 90 min at 50-60°C. After the addition of 15 ml water to the red-orange solution the product was separated and recrystallized from EtOH/AcOEt (1:1). – Pale yellow needles; yield: 3.9 g (38%); m.p. 156-157°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.61 (s, 4 H, S-CH<sub>2</sub>), 3.84, (s, 6 H, OCH<sub>3</sub>), 6.89 (m, 4 H, o-H [C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]), 7.04 (m, 4 H, phenyl-H), 7.06 (s, 2 H, =CH-), 7.20 (m, 6 H, phenyl-H), 7.43 (m, 4 H, m-H [C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 37.7 (S-CH<sub>2</sub>), 55.3 (OCH<sub>3</sub>), 113.9 (o-C [C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]), 126.8 (p-C [phenyl]), 128.2, 128.7, 129.3 (o-C, m-C [phenyl]; m-C [C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]), 131.3 (=CH-), 132.7, 137.9, 138.5 (>C=), 159.6 (C-OCH<sub>3</sub>). – MS (70 eV): m/z = 510 (3%) [M<sup>+</sup>], 419 (24) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub>], 296 (100) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 281 (32) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> - CH<sub>3</sub>], 91 (37) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>32</sub>H<sub>30</sub>O<sub>2</sub>S<sub>2</sub> (510.7): calcd. C 75.26, H 5.92, S 12.56; found C 75.00, H 6.10, 12.50. – From the oily intermediate the mono-adduct [(Z)-1-benzylthio-1,4-di(p-methoxyphenyl)-1-buten-3-yne (3m; cf. ref.<sup>1c</sup>)] could be separated by fractional crystallization from ethanol: Colourless needles, m.p. 72-73°C; MS (70 eV): m/z = 386 (8) [M<sup>+</sup>]; C<sub>25</sub>H<sub>22</sub>O<sub>2</sub>S (386.5): calcd. C 77.69, 5.74; found C 78.10, H 6.12.

(Z,Z)-1,4-Di(benzylthio)-1,4-di(p-*n*-butylphenyl)-1,3-butadiene (4n). – As described for 4h/procedure I using 15.7 g (0.05 mol) 1n (m.p. 72°C): Pale yellow needles (EtOH/AcOEt); yield: 7.6 g (27%); m.p. 79-80°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta \approx 0.96$  (t, 6 H, CH<sub>3</sub>), 1.30-1.71 (m, 8 H, CH<sub>2</sub>), 2.64 (t, 4 H, CH<sub>2</sub>), 3.62 (s, 4 H, CH<sub>2</sub>), 7.02-7.45 (m, 20 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta \approx 14.0$  (CH<sub>3</sub>), 22.4, 33.6, 35.4 (CH<sub>2</sub>), 37.7 (S-CH<sub>2</sub>), 126.8, 128.0, 128.2, 128.5, 128.8, 131.8, 137.4, 138.5, 138.8, 143.0 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 562 (10), [M<sup>+</sup>], 471 (51) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub>], 438 (8) [M<sup>+</sup> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 348 (100) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub> - SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 305 (27) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub>C<sub>6</sub>H<sub>5</sub> - HSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>]. – C<sub>38</sub>H<sub>42</sub>S<sub>2</sub> (562.8): calcd. C 81.09, H 7.52, S 11.39; found C 80.77, H 7.51, S 11.33.

(Z,Z)-1,4-Di(benzylthio)-1,4-di(p-t-butylphenyl)-1,3-butadiene (40). – As described for 4h/procedure I using 15.7 g (0.05 mol) 10 (m.p. 90°C): Pale yellow needles (EtOH/AcOEt[1:1]; yield: 22.5 g (80%); m.p. 166-167°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.45 (s, 18 H, CH<sub>3</sub>), 3.73 (s, 4 H, CH<sub>2</sub>), 7.13, 7.58 (m, 20 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 31.3 (CH<sub>3</sub>), 34.6 (>C<), 37.7 (CH<sub>2</sub>), 125.3, 126,5, 127,7, 127.8, 128,2, 128.8, 131,2, 131,7, 138.5, 138.7 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 562 (14) [M<sup>+</sup>], 471 (50) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 438 (11) [M<sup>+</sup> – HSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 380 (4) [M<sup>+</sup> – 2 C<sub>7</sub>H<sub>7</sub>], 348 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub> – CH<sub>3</sub>]. – C<sub>38</sub>H<sub>42</sub>S<sub>2</sub> (562.8): calcd. C 81.09, H 7.52, S 11.39; found C 81.31, H 7.50, S 11.21.

(*Z*,*Z*)-1,4-Di(benzylthio)-1,4-di(p-cyclohexylphenyl)-1,3-butadiene (4p). – As described for 4h/procedure I using 18.3 g (0.05 mol) 1p (m.p. 232°C) and 28 h reaction time: Pale yellow prisms (EtOH/AcOEt/benzene [1:1:1]); yield: 15.6 g (51%); m.p. 172°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.29$ -1.89 (m, 22 H, cyclohexyl), 3.62 (s, 4 H, S-CH<sub>2</sub>), 7.02-7.46 (m, 20 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 26.2$ , 26.9, 34.4 (CH<sub>2</sub>), 37.7 (S-CH<sub>2</sub>), 44.3 (CH), 126.8, 128.0, 128.2, 128.8, 131.8, 137.6, 138.6, 138.8, 148.2 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 614 (24) [M<sup>+</sup>], 523 (76) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 490 (35) [M<sup>+</sup> – HSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 400 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>]. – C4<sub>2</sub>H<sub>4</sub>S<sub>2</sub> (614.9): calcd. C 80.03, H 7.54, S 10.43; found C 80.03, H 7.43, S 10.40. – By fractional crystallization (EtOH/AcOEt [1:1]) the mono-addition product [(*Z*)-1-benzylthio-1,4-di(p-cyclohexylphenyl)-1-buten-3-yne (3p)] could also be separated: Pale yellow needles; yield 6.8 g (28%); m.p. 117°C (EtOH/AcOEt [1:1]). – IR (KBr):∇ = 2160 cm<sup>-1</sup> (C=C). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.24$ -2.48 (m, 22 H, cyclohexyl), 3.90 (s, 2 H, S-CH<sub>2</sub>), 74.5 (m, 14 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 26.1$ , 26.3, 26.9, 34.2, 34.3, 34.4 (CH<sub>2</sub>), 37.5 (S-CH<sub>2</sub>), 44.3, 44.5 (CH), 87.2, 110.0 (-C≡), 120.9, 126.9, 127.9, 128.0, 128.2, 128.8, 128.9, 131.4, 132.5, 136.5, 138.0, 148.4, 148.5, 148.9, (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 490 (90) [M<sup>+</sup>], 366 (100) [M<sup>+</sup> – HSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>]; C<sub>35</sub>H<sub>38</sub>S (490.7): calcd. C 85.66, H 7.81, S 6.53; found C 85.49, H 7.85, S 6.72.

(Z,Z)-1,4-Di(benzylthio)-1,4-di[2-(5,6,7,8-tetrahydronaphthyl)]-1,3-butadiene (4q; cf. ref.<sup>1d</sup>). – As described for 4h/procedure I using 15.5 g (0.05 mol) 1q (m.p. 163°C) and 40 h reaction time: Pale yellow needles (EtOH/ AcOEt [3:2]); yield: 17.8 g (64%); m.p. 152°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.53-2.78 (m, 16 H, C<sub>4</sub>H<sub>8</sub>), 3.61 (s, 4 H, S-CH<sub>2</sub>), 6.99-7.29 (m, 18 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 23.2, 29.3, 29.5 (CH<sub>2</sub>), 37.8 (S-CH<sub>2</sub>), 125.3, 126.8, 128.2, 128.7, 128.8, 129.0, 129.1, 137.1, 137.4, 138.7, 138.8, 141.6 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 558 (3) [M<sup>+</sup>], 434 (73) [M<sup>+</sup> – HSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 344 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>]. – C<sub>38</sub>H<sub>38</sub>S<sub>2</sub> (558.8): calcd. C 81.67, H 6.85, S 11.48; found C 81.45, H 6.92, S 10.94.

 $(Z_2Z)$ -1,4-Di(benzylthio)-1,4-di(4-biphenylyl)-1,3-butadiene (4r; cf. ref.<sup>1d</sup>). – *Procedure I*: 3.54 g (0.01 mol) 1r (m.p. 231°C), 3.5 g (0.025 mol) 2, 2.4 g (0.06 mol) NaOH and 26 ml dry ethanol were heated at 80°C for 40 h in a sealed tube. Isolation as described for 4h/procedure I. – Yellow needles (benzene; very poorly soluble); yield: 1.99 g (33%); m.p. 213-214°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.67 (s, 4 H, S-CH<sub>2</sub>), 7.06, 7.21, 7.45, 7.64 (m, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 37.9 (S-CH<sub>2</sub>), 126.9 (127.0 (9), 127.4, 128.3, 128.5, 128.7 (9), 128.8 (3) (aromat. C), 132.5 (=CH-), 138.4, 138.8 (7), 139.0 (3), 140.6, 140.9 (>C=). – MS (70 eV): m/z = 602 (4) [M<sup>+</sup>], 511 (21) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 389 (53), 388 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – S-CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 91 (78) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>42</sub>H<sub>34</sub>S<sub>2</sub> (602.8): calcd. C 83.67, H 5.68; found C 83.42, H 6.06. – *Procedure II*: A mixture of 1.28 g (10.32 mmol) **2**, 0.16 g (2.85 mmol) KOH and 20 ml dry DMF was stirred 30 min. After the addition of 1.6 g (4.5 mmol) **1r** stirring was continued 3 h at room temperature. The precipitate was filtered by suction and recrystallized from benzene, 0.7 g (26%).

(Z,Z)-1,4-Di(benzylthio)-1,4-di(1-naphthyl)-1,3-butadiene (4s; cf. ref.<sup>1d</sup>). – *Procedure I*: As described for 1r/procedure I using 3.02 g (0.01 mol) 1s (m.p. 177°C): Colourless needles; yield 1.71 g (31%); m.p. 189-190°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.39 (s, 4 H, S-CH<sub>2</sub>), 6.87 (m, 4 H, o-H [benzyl]), 7.07 (m, 8 H, aromat. H, =CH-), 7.49 (m, 8 H), 7.85 (m, 4 H), 8.26 (m, 2 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 36.6 (S-CH<sub>2</sub>), 125.1, 125.9, 126.0, 126.1, 126.8, 127.4 (aromat. C), 128.1 (phenyl C<sub>0/m</sub>), 128.2, 128.4 (aromat. C), 128.7 (phenyl C<sub>0/m</sub>), 129.3 (aromat. C), 131.8, 133.7 (>C=), 137.5(8), 137.6(0), 137.6(3) (>C= [naphthyl]). – MS (70 eV): m/z = 550 (26) [M<sup>+</sup>], 459 (24) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub>], 36 (100) [M<sup>+</sup> – C<sub>7</sub>H<sub>7</sub> – SCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 91 (79) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. – C<sub>38</sub>H<sub>30</sub>S<sub>2</sub> (550.8): calcd. C 82.87, H 5.49; found C 82.82, H 5.71. – *Procedure II*: A mixture of 2.8 g (9.3 mmol) 1s, 3.0 g (24 mmol) 2, 11 g (196 mmol) KOH, 0.52 g (2 mmol) 18-crown-6, 30 ml water and 60 ml benzene was heated under reflux for 7 h. The organic phase was washed with water and concentrated. The resulting oily layer was again treated with 2.0 g (15.9 mmol) 2 and 300 mg (5.35 mmol) KOH in 10 ml dry DMF and stirred at 40°C for 1 h. The precipitated solid was recrystallized from benzene, yield: 2.25 g (44%).

 $(\textbf{Z,Z})-1,4-Di(benzylthio)-1,4-di(2-thienyl)-1,3-butadiene (4t; cf. ref.^{1c}). - As described for 4h/procedure I using 10.7 g (0.05 mol) 1t (m.p. 92°C): Yellow needles; yield: 13.4 g (58%); m.p. 115-116°C. - <sup>1</sup>H NMR (CDCl<sub>3</sub>): <math>\delta$  = 3.81 (s, 4 H, S-CH<sub>2</sub>), 7.00-7.28 (m, 18 H, aromat. H, thienyl H, =CH-). - <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 39.2 (S-CH<sub>2</sub>), 125.6, 126.2, 127.0, 127.8 (aromat. C, thienyl C), 128.3, 128.8 (phenyl C<sub>0/m</sub>), 131.1 (>C=), 131.9 (=CH-), 138.1, 146.2 (>C=). - MS (70 eV): m/z = 462 (14) [M<sup>+</sup>], 371 (72) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub>], 248 (100) [M<sup>+</sup> - C<sub>7</sub>H<sub>7</sub> - SSCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>], 91 (89) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. - C<sub>26</sub>H<sub>22</sub>S<sub>4</sub> (462.7): calcd. 67.48, H 4.79; found C 66.97, H 4.82.

(Z,Z)-1,4-Di(*t*-butylthio)-1,4-diphenyl-1,3-butadiene (8h). – A mixture of 1.8 g (20 mmol) 7, 300 mg KOH and 40 ml dry DMF was stirred at room temperature for 30 min and then treated slowly with 2.02 g (10 mmol) 1h. Stirring was continued 30 min (after 10 min a solid precipitates), then 200 ml water were added. The product was filtered by suction and recrystallized from trichloroethene or EtOH/CHCl<sub>3</sub>. – Colourless needles; yield:3.35 g (87.7%); m.p. 238-239°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.15$  (s, 18 H, *t*-Bu), 7.73 (s, 2 H, =CH-), 7.32-7.70 (m, 10 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 31.8$  (C[CH<sub>3</sub>]<sub>3</sub>), 48.2 (C[CH<sub>3</sub>]<sub>3</sub>), 127.8 (=CH-), 128.0(5), 128.1(2) (aromat. C<sub>0,m</sub>), 138.0 (>C=), 138.9 (aromat. C<sub>p</sub>), 144.0 (aromat. C<sub>i</sub>). – MS (70 eV): m/z = 382 (33.5) [M<sup>+</sup>], 325 (16) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub>], 269 (58) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - C<sub>4</sub>H<sub>8</sub>], 237 (100) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - S-C<sub>4</sub>H<sub>8</sub>], 204 (11) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - S-C<sub>4</sub>H<sub>8</sub> - SH]. – C<sub>24</sub>H<sub>38</sub>S<sub>2</sub> (382.6): calcd. C 75.34, H 7.90, S 16.76; found C 75.16, H 7.71, S 17.25.

(Z,Z)-1,4-Di(*t*-butylthio)-1,4-di(p-tolyl)-1,3-butadiene (8k). – As described for 8h using 2.3 g (10 mmol) 1k: Colourless needles (AcOEt); yield: 3.4 g (83.2%); m.p. 244-245°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.17$  (s, 18 H, *t*-Bu), 2.37 (s, 6 H, CH<sub>3</sub>C<sub>6</sub>H<sub>5</sub>], 7.77 (s, 2 H, =CH-), 7.15 (*pseudo*-d, J = 8 Hz, 4 H, aromat. H [C<sub>0</sub>H]), 7.60 (*pseudo*-d, 4 H, J = 8 Hz, aromat. H [CH<sub>m</sub>]). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 21.1$  (CH<sub>3</sub>-C<sub>6</sub>H<sub>5</sub>), 31.7 (C[CH<sub>3</sub>]<sub>3</sub>), 47.9 (C[CH<sub>3</sub>]<sub>3</sub>), 127.8, 128.7 (C<sub>0,m</sub> [tolyl]), 137.4 (=CH-), 137.5 (>C=), 138.2 (C<sub>i</sub> [tolyl]), 141.1 (C<sub>i</sub> [phenyl]). – MS (70 eV): m/z = 410 (21) [M<sup>+</sup>], 353 (18) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub>], 297 (55) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - C<sub>4</sub>H<sub>8</sub>], 265 (100) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - S-C<sub>4</sub>H<sub>8</sub>], 232 (8) [M<sup>+</sup> - C<sub>4</sub>H<sub>9</sub> - S-C<sub>4</sub>H<sub>8</sub>] – SH]. – C<sub>26</sub>H<sub>34</sub>S<sub>2</sub> (410.7): calcd. C 76.04, H 8.34, S 15.62; found C 76.02, H 8.26, S 15.70.

 $(\textbf{Z,Z})-1,4-Di(t-butylthio)-1,4-di(p-methoxyphenyl)-1,3-butadiene (8m). - As described for 8h using 2.62 g (10 mmol) 1m: Colourless needles (trichloroethene); yield: 3.60 g (81.5%); m.p. 137-138°C. - <sup>1</sup>H NMR (CDCl<sub>3</sub>): <math>\delta = 1.16$  (s, 18 H, t-Bu), 3.82 (s, 6 H, OCH<sub>3</sub>), 7.71 (s, 2 H, =CH-), 6.87 (2 d, 4 H, J = 2 Hz, aromat. H<sub>o</sub> [CH<sub>3</sub>O-C<sub>6</sub>H<sub>4</sub>]), 6.30 (2 d, 4 H, J = 2 Hz, aromat. H<sub>m</sub> [CH<sub>3</sub>O-C<sub>6</sub>H<sub>4</sub>]). - <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 31.8$  (C[CH<sub>3</sub>]<sub>3</sub>), 48.0 (C[CH<sub>3</sub>]<sub>3</sub>), 55.3 (OCH<sub>3</sub>), 113.5, 129.2 (aromatic. C<sub>0,m</sub>), 136.6 (=CH-), 136.8 (>C=), 137.4 (aromat. C<sub>i</sub> {=C[S]-C=}), 159.5 (aromat. C<sub>i</sub> [=C-OCH<sub>3</sub>]). - C<sub>26</sub>H<sub>34</sub>O<sub>2</sub>S<sub>2</sub> (442.7): calcd. C 70.54, H 7.74, S 14.49; found C 70.68, H 7.58, S 14.51.

**Transformation to 1,2-dithiines 6** (all preparative and isolation steps must be carried out with the exclusion of air): **1,2-Dithiine (6a**; cf. ref.<sup>1a,c</sup>). – To a solution of 50 g (308 mmol) anhydrous FeCl<sub>3</sub> in 250 ml dry methanol was dropped (2 h; 20°C) with stirring a solution of 15.4 g (130 mmol) **10a** (see below) in 100 ml dry methanol. The mixture was diluted with 1.5 l water and extracted twice with 300 ml petroleum ether (b.p. 30-50°C; stabilization with a small amount of N-cyclohexylmaleinimide). After careful evaporation i. vac. (bath temp. at -10 till -20°C) the residue was distilled i. vac. (to reduce polymerization caused by alkali traces the apparatus was previously treated with conc. H<sub>2</sub>SO<sub>4</sub> at 200°C, washed with water and dryed at 150°C). – Dark red oil; yield: 1,2 g (8%; a major part remains in the flask as viscous material [polymer or polydisulfide?]). – UV/Vis (cyclohexane):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 222 (3.22), 267 (3.18), 274 (3.20), 355 (2.25; sh), 457 (1.98) nm. – IR (CHCl<sub>3</sub>): $\overline{v}$  = 620, 1300 (s, m; cis-HC=CH), 1300, 1540 (w-m; C=C), 3005 (m; =CH) cm<sup>-1. – 1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 6.05 (m, 2 H, C<sup>3</sup> = <sup>6</sup>H), 6.27 (m, 2H, C<sup>4</sup> = <sup>5</sup>H); AA'BB'-type: <sup>3</sup>J<sub>3,4</sub> = <sub>6.5</sub> = 9.4, <sup>4</sup>J<sub>3,5</sub> = <sub>6.4</sub> = 0, <sup>3</sup>J<sub>4,5</sub> = 5.5, <sup>5</sup>J<sub>3,6</sub> = 2.0 Hz. – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 119.5 (C<sup>3</sup> = <sup>6</sup>H; <sup>1</sup>J<sub>C,H</sub> = 176.2 Hz), 129.8 (C<sup>4</sup> = <sup>5</sup>H; <sup>1</sup>J<sub>C,H</sub> = 162.9 Hz). – MS (70 eV): m/z = 116 (100) [M<sup>+</sup>], 84 (81) [M<sup>+</sup> – S], 71 (66) [M<sup>+</sup> – S – CH], 58 (30) [M<sup>+</sup> – S – 2 CH]. – C<sub>4</sub>H<sub>4</sub>S<sub>2</sub> (116.2): calcd. C 41.35, H 3.47, S 55.18; found C 41,45, H 3.80, S 54.92.

**3,6-Di**(*n*-butyl)-1,2-dithiine (6b). – To 11.1 g (27 mmol) powdered 4b in 200 ml liquid ammonia was added 2.53 g (110 mmol) Na at -70°C over 90 min. Solid NH<sub>4</sub>Cl was added until the blue colour of the solution had disappeared. After evaporation, the residue was dissolved in 200 ml of water and washed twice with two 20 ml portions of Et<sub>2</sub>O. The resulting aqueous solution of 5b was covered with 100 ml portions of Et<sub>2</sub>O and acidified at -5°C with 25% aqueous HCl solution. The organic phase 10b was evaporated under reduced pressure (bath temp. < 15°C), the resulting pale yellow oil dissolved in 30 ml MeOH and a solution of FeCl<sub>3</sub> in 20 ml MeOH slowly added with stirring under an argon atmosphere. The mixture was stirred for a further 2 h at room temperature, diluted with 100 ml water and extracted with cyclohexane. On concentration 4.0 g (65%) of a deep red oil was obtained (R<sub>f</sub> = 0.65 [silica gel, benzene]) which had a high tendency for sulfur extrusion. – UV/Vis (MeCN):  $\lambda_{max} = 246$ , 440 nm (compare, however, the product in ref.<sup>10</sup>). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 0.92$  (t<sub>c</sub>, 6 H, CH<sub>3</sub>), 1.01-1.83 (m, 8 H, CH<sub>2</sub>), 2.06-2.41 (m, 4 H, CH<sub>2</sub>), 6.62 (s, 2 H, =CH-). – C<sub>12</sub>H<sub>20</sub>S<sub>2</sub> (228.4): calcd. C 63.10, H 8.83, S 28.07; found C 62.86, H 8.89, S 27.90.

**3,6-Di(2-hydroxy-2-propyl)-1,2-dithiine** (6d). – Approximately 1 g (43 mmol) of Na was added slowly to a solution 0f 3.32 g (8 mmol) 4d in 120 ml liquid ammonia at -70°C until a deep blue colour persisted. After 2 h further stirring at this temperature the mixture was neutralized by the addition of NH<sub>4</sub>Cl (decolourization) and the solution allowed to evaporate. The resulting residue was dissolved in 40 ml 2N NaOH solution, insoluble material removed by filtration and the product purified by extraction with Et<sub>2</sub>O. The orange aqueous phase 5d was slowly added with stirring to a 5% aqueous solution of K<sub>3</sub>[Fe(CN)<sub>6</sub> (6.6 g [20 mmol]) and the precipitated product recrystallized from *n*-hexane/CHCl<sub>3</sub> (3-4:1). – Yellow-orange needles; yield: 1.1 g (59.1%); m.p. 108-109°C. – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 262 (3.44), 412 (2.60) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.49 (s, 12 H, CH<sub>3</sub>), 1.74 (s, 2 H, OH), 6.39 (s, 2 H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 29.0 (CH<sub>3</sub>), 73.4 (-C[CH<sub>3</sub>]<sub>2</sub>OH), 122.6 (=CH-), 144.2 (>C=). MS (70 eV): m/z = 232 (57) [M<sup>+</sup>], 200 (11) [M<sup>+</sup> - S], 185 [M<sup>+</sup> - S - CH<sub>3</sub>], 167 (13) [M<sup>+</sup> - S - CH<sub>3</sub> - H<sub>2</sub>O], 159 (100) [M<sup>+</sup> - S - CH<sub>3</sub> - H<sub>2</sub>O - C<sub>2</sub>H<sub>2</sub>], 59 (56) [C(CH<sub>3</sub>)<sub>2</sub>OH<sup>+</sup>]. – C<sub>10</sub>H<sub>16</sub>O<sub>2</sub>S<sub>2</sub> (232.4): calcd. C 51.69, H 6.94, S 27.60; found 51.89, H 7.20, S 27.27.

**3,6-Di(pyrrolidinomethyl)-1,2-dithine** (**6e**). – As described for **6d** using 2.10 g (4.5 mmol) **4e**, 75 ml liquid NH<sub>3</sub>, 529 mg (23 mmol) Na, 25 ml 2N NaOH solution and 3.95 g (12 mmol) K<sub>3</sub>[Fe(CN)<sub>6</sub> (in a 5% aqueous solution), recrystallization from AcOEt. – Bright yellow needles; yield: 661 mg (52%); m.p. 72-73°C;<sup>39</sup> R<sub>f</sub> = 0.18 (silica gel, *n*-hexane(AcOEt [1:1]). – UV/Vis (*n*-hexane):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 268 (3.67), 4.23 (2.19) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.69-1.85 (m, 8 H, CH<sub>2[cycl.C3/4]</sub>), 2.46-2.54 (m, CH<sub>2[cycl.C2/5</sub>), 3.31 (s, 4 H, >N-CH<sub>2</sub>), 6.05 (s, 2 H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 23.7 (CH<sub>2[cycl.C3/4]</sub>), 54.0 (CH<sub>2[cycl.C2/5</sub>), 61.4 (>N-CH<sub>2</sub>), 125.1 (=CH-), 134.4 (>C=). – MS (70 eV): m/z = 282 (12) [M<sup>+</sup>], 250 (4) [M<sup>+</sup> – S], 180 (11) [M<sup>+</sup> – S – C<sub>4</sub>H<sub>8</sub>N], 111 (7) [M<sup>+</sup> – S – 2 C<sub>4</sub>H<sub>8</sub>N], 84 (100) [pyrrolidino-CH<sub>2</sub><sup>+</sup>]. – C<sub>14</sub>H<sub>22</sub>N<sub>2</sub>S<sub>2</sub> (282.5): calcd. C 59.53, H 7.85, N 9.92, S 22.70; found C 59.42, H 7.93, N 9.80, S 22.85.

**3,6-Di(piperidinomethyl)-1,2-dithime (6f)**. – As described for **6d/6e** using 2.22 g (4.5 mmol) **4f**; recrystallization from AcOEt – Bright yellow needles; yield: 800 mg (55%); m.p. 112-113°C;<sup>39</sup> R<sub>f</sub> = 0.42 (*n*-hexane/AcOEt [1:1]). – UV/Vis (*n*-hexane):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 282 (3.64), 425 (2.31) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.38-1.40 (m, 4 H, CH<sub>2[cycl.C4]</sub>), 1.51-1.57 (m, 8 H, CH<sub>2[cycl.C3/5]</sub>), 2.37-2.39 (m, 8 H, CH<sub>2[cycl.C2/6]</sub>), 3.15 (s, 4 H, >N-CH<sub>2</sub>), 6.12 (s, 2 H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 24.3 (CH<sub>2[cycl.C4]</sub>), 26.0 (CH<sub>2[cycl.C3/5]</sub>), 54.4 (CH<sub>2[cycl.C2/6]</sub>), 64.7 (>N-CH<sub>2</sub>), 125.3 (=CH-), 134.3 (>C=). – MS (70 eV): m/z = 310 (23) [M<sup>+</sup>], 278 (5) [M<sup>+</sup> – S], 194 (14) [M<sup>+</sup> – S – C<sub>5</sub>H<sub>10</sub>N], 111 (6) [M<sup>+</sup> – S – 2 C<sub>5</sub>H<sub>10</sub>N], 98 (100) [piperidino-CH<sub>2</sub><sup>+</sup>]. – C<sub>16</sub>H<sub>26</sub>N<sub>2</sub>S<sub>2</sub> (310.5): calcd. C 61.88, H 8.44, N 9.02, S 20.65; found C 61.14, H 8.31, N 9.11, S 20.64.

**3,6-Di(morpholinomethyl)-1,2-dithilne (6g)**. – As described for **6d/6e** using 2.24 g (4.5 mmol) **4g**; recrystallization from AcOEt. – Bright yellow needles; yield: 849 mg (60%); m.p. 159-160°C;  $^{39}$  R<sub>f</sub> = 0.24 (*n*-hexane/AcOEt [1:1]). – UV/Vis (*n*-hexane):  $\lambda_{max}$  (lg  $\epsilon$ ) = 280 (3.53), 423 (2.53) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.40-2.51 (m, 8 H, CH<sub>2[cycl.C2/6]</sub>), 3.21 (s, 4 H, >N-CH<sub>2</sub>), 3.64-3.75 (m, 8 H, CH<sub>2[cycl.C3/5]</sub>), 6.17 (s, 2 H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 53.3 (CH<sub>2[cycl.C2/6]</sub>), 64.3 (>CH<sub>2</sub>), 67.0 (CH<sub>2[cycl.C3/5]</sub>), 126.0 (=CH-, J<sup>1</sup><sub>C,H</sub> = 159.6 Hz), 133.5 (>C=). – MS (70 eV): m/z = 314 (40) [M<sup>+</sup>], 282 (6) [M<sup>+</sup> – S], 196 (14) [M<sup>+</sup> – S – C<sub>4</sub>H<sub>8</sub>NO], 111 (7), [M<sup>+</sup> – S – 2 C<sub>4</sub>H<sub>8</sub>NO], 100 (100) [morpholino-CH<sub>2</sub><sup>+</sup>]. – C<sub>14</sub>H<sub>22</sub>N<sub>2</sub>O<sub>2</sub>S<sub>2</sub> (314.5): calcd. C 53.47, H 7.05, N 8.91, S 20.39; found C 53.42, H 7.08, N 8.37, S 20.41.

**3,6-Diphenyl-1,2-dithiine (6h**; cf. ref.<sup>1c,f</sup>). – *Method I*: A solution of 2.25 g (5 mmol) of **4b** in 100 ml liquid ammonia was treated slowly with about 0.5 g (22 mmol) of Na, then stirred for a further 2 h, neutralized by the addition of NH<sub>4</sub>Cl (disappearance of the blue colour) and the solution evaporated. The residue was dissolved in 25 ml 2N NaOH under an argon atmosphere and filtered to remove insoluble material. The resulting solution of **5h** was added drowise with stirring to a 5% aqueous solution of K<sub>3</sub>Fe(CN)<sub>6</sub> (4.5 g [13.5 mmol]). The product precipitated immediately. Alternatively

if a current of air was passed through the above alkaline solution (3 h), the product separated as a fine crystalline material. – Purple leaflets (AcOEt); yield: 724 mg (54%); m.p. 144-145°C;<sup>39</sup>  $R_f = 0.77$  (*n*-hexane/AcOEt [3:1]). – UV/Vis  $(\text{cyclohexane}): \lambda_{\text{max}} (\text{Ig } \epsilon) = 237 (4.23), 312 (4.13), 468 (3.51) \text{ nm}; (MeCN): 241 (4.15), 310 (4.19), 464 (3.72) \text{ nm};$ (EtOH): 233 (4.25), 307 (4.22), 461 (3.55) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 6.90 (s, 2 H, =CH-), 7.29-7.82 (m, 10 H, aromat. H).  $- {}^{13}C$  NMR (CDCl<sub>3</sub>): see data in ref. <sup>1f</sup>, p. 606. - MS (70 eV): m/z = 268 (100) [M<sup>+</sup>], 236 (31) [M<sup>+</sup> - S], 202 (9)  $[M^+ - H_2S]$ , 147 (7)  $[M^+ - C_6H_5CS]$ , 121 (19)  $[C_6H_5CS^+]$ .  $-C_{16}H_{12}S_2$  (268.4): calcd. C 71.60, H 4.51, S 23.89; found C 71.43, H 4.54, S 23.32; molar weight (cryoscopic in benzene: 256). - Method II: A mixture of 1.91 g (5 mmol) 8h, 25 ml trichloroethene and 25 ml AcOH was stirred and treated portionwise with 11.89 g 2-nitrophenylsulfenyl chloride and the resulting pale yellow solution stored at 60°C for 6 h. After concentration at reduced pressure, the solids obtained were washed with water, dried and the crude 9h used for subsequent transformations to 6h; (Z,Z)-1,4-Di(o-nitrophenyldithio)-1,4-diphenyl-1,3-butadiene (9h). - Yellow leaflets; yield: 2.55 g (92%); m.p. 185-186°C (rapid heating); very poorly soluble in most solvents; in trichloroethene, DMF and DMSO colour change to red occurs with precipitation of  $6h_{-}$  MS (70 eV): m/z = 268 (25) [ $6h^{+}$ ], 308 (25) [di(2-nitrophenyl) disulfide<sup>+</sup>], 236 (100) [ $6h^{+} - S$ ], 154 (100) [di(2-nitrophenyl) disulfide/2<sup>+</sup>]; no molpeak recognizable. - [C<sub>28</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub>S<sub>4</sub> (576.7)]. - To a suspension of 9h in 20 ml dry THF was added with stirring 8 mg (0.08 mmol) NEt<sub>3</sub> and 6 mg (0.08 mmol) 2-mercaptoethanol. A red solution was obtained after 20 min. The mixture was allowed to stand at room temperature for a further 30 min, the solvents removed at reduced pressure and the residue dissolved in Et<sub>2</sub>O. The solution was washed twice with 5% NaHCO<sub>3</sub> solution and then with water to remove any thiol. The etheral solution was concentrated and the residues recrystallized from AcOEt to eliminate the co-produced di(2-nitrophenyl) disulfide. Yield of 6h: 300 mg (58%). - Method III: A suspension of 0.72 g (2 mmol) 19 in 30 ml dry diethyl ether was treated dropwise at -10°C with 3 ml (4.8 mmol) of a 1.6M *n*-BuLi solution in 20 ml diethyl ether under an argon atmosphere. After stirring for 30 min at  $-10^{\circ}$ C and 20 min at room temperature the mixture was cooled to -40°C and an excess of finely powdered elemental sulfur (0.32 g) was added. Stirring was continued at the same temperature for 2 h. After standing overnight at room temperature, 40 ml of a 10% NaOH solution was added at 0 to 5°C. The residue obtained after evaporation of the solvents was washed with n-hexane and recrystallized from EtOAc; yield of 6h: 0.376 g (70%).

**3-Phenyl-6-(p-tolyl)-1,2-dithine (6j**; cf. ref.<sup>1d</sup>). – As described for **6h**/method I using 2.32 g (5 mmol) **4j**: Purple needles (AcOEt); yield: 1.0 g (71%); m.p. 126-127°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 238 (4.09), 316 (4.14), 464 (3.42) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.37 (s, 3 H, CH<sub>3</sub>), 6.85, 6.90 (2 d [J = 6.7 Hz], 2 H, =CH-), 7.18-7.70 (m, 4 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 21.2 (CH<sub>3</sub>), 125.1 (=CH-), 126.2, 127.7, 128.7, 128.8, 129.4 (aromat. CH), 133.7 (>C<sup>1</sup>= [tolyl]), 134.0 (>C<sup>6</sup>= [dithine]), 135.1 (>C<sup>3</sup>= [dithine]), 136.8 (>C= [phenyl]), 139.1 (>C<sup>4</sup>= [tolyl]). – MS (70 eV): m/z = 282 (M<sup>+</sup>], 250 (100) [M<sup>+</sup> – S], 267 (8) [M<sup>+</sup> – CH<sub>3</sub>], 216 (8) [M<sup>+</sup> – S – H<sub>2</sub>S]. – C<sub>17</sub>H<sub>14</sub>S<sub>2</sub> (282.4): calcd. C 72.30, H 5.00, S 22.70; found C 72.24, H 5.12, S 23.16.

**3,6-Di(p-tolyl)-1,2-dithine (6k**; cf. ref.<sup>1c</sup>). – *Method I*: As described for **6h**/method I using 2.39 g (5 mmol) **4k**: Purple prisms (AcOEt); yield: 711 mg (48%); m.p. 161-162°C<sup>39</sup>;  $R_f = 0.60$  (benzene/*n*-hexane 1:1). – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 241 (4.23), 316 (4.25), 464 (3.64). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 2.36$  (s, 6 H, CH<sub>3</sub>), 6.85 (s, 2 H, =CH-), 7.19, 7.56 (2 d [J = 8 Hz)], 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 21.2$  (CH<sub>3</sub>), 125.3 (=CH-), 127.7, 129.4 (aromat. CH), 133.4 (>C<sup>1</sup> = [toly1]), 134.4 (>C= [dithine]), 139.1 (>C<sup>4</sup> = [toly1]). – MS (70 eV): m/z = 296 (35) [M<sup>+</sup>], 264 (100) [M<sup>+</sup> - S], 248 (28), [M<sup>+</sup> - S - CH<sub>4</sub>], 215 (28), [M<sup>+</sup> - S - CH<sub>4</sub> - SH]. – C<sub>18</sub>H<sub>16</sub>S<sub>2</sub> (296.4): calcd. 72.93, H 5.44, S 21.63; found C 73.01, H 5.50, S 21.50. – *Method II*: As described for **6h** /method II using 2.05 g (5 mmol) **8k** and yielding first (**Z,Z)-1,4-Di(o-nitrophenyldithio)-1,4-di[p-toly1]-1,3-butadiene (9k**). – Yellow leaflets; yield: 2.8 g (93%); m.p. 179-181°C (rapid heating up); very poorly soluble in most solvents. – MS (70 eV): m/z = 308 (10) [di(2-nitrophenyl) disulfide<sup>+</sup>], 296 (32) [**6 k**<sup>+</sup>], 264 (100) [**6 k** - S], 154 (15) [di(2-nitrophenyl) disulfide/2<sup>+</sup>]; no molpeak recognizable. – [C<sub>30</sub>H<sub>24</sub>N<sub>2</sub>O<sub>4</sub>S<sub>4</sub> (604.8)]. – Transformation using 1.2 g (2 mmol) **9k**; yield of **6k**: 361 mg (61%).

**3-(p-Methoxyphenyl)-6-phenyl-1,2-dithine (6**]; cf. ref.<sup>1d</sup>). – As described for **6h**/method I using 2.40 g (5 mmol) **4**]: Dark-red needles (AcOEt); yield: 985 mg (66%); m.p. 140-141°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 241 (4.15), 320 (4.19), 464 (3.60) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.83 (s, 3 H, CH<sub>3</sub>), 6.80, 6.89 (2 d [J = 6.7 Hz], 2 H, =CH-), 6.93, 7.62 (2 d [J = 8.9 Hz], 2 H, aromat. H [C<sub>6</sub>H<sub>4</sub>OCH<sub>3</sub>]), 7.31-7.70 (m, 7 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 55.38 (CH<sub>3</sub>), 114.2, 124.2, 126.4, 127.6, 128.7 (aromat. C), 129.2 (=CH- [dithiine]), 129.2 (>C= [phenyl]), 133.1 (>C<sup>6</sup>= [dithiine]), 135.1 (>C<sup>1</sup>= [methoxyphenyl]), 136.9 (>C<sup>3</sup>= [dithiine]), 160.4 (>C<sup>4</sup>= [methoxyphenyl]). – MS (70 eV): m/z = 298 (29) [M<sup>+</sup>], 266 (100) [M<sup>+</sup> - S], 283 (14) [M<sup>+</sup> - CH<sub>3</sub>], 251 (90) [M<sup>+</sup> - S - CH<sub>3</sub>], 223 (40) [M<sup>+</sup> - S - CH<sub>3</sub> - CO]. – C<sub>17</sub>H<sub>14</sub>OS<sub>2</sub> (298.4): calcd. C 68.42, H 4.73, S 21.49; found C 68.44, H 5.03, S 21.85.

**3,6-Di(p-methoxyphenyl)-1,2-dithine (6m**; cf. ref.<sup>1</sup>c). – *Method I*: As described for **6h**/method I using 2.55 g (5 mmol **4m**: Purple needles (AcOEt); yield: 1.18 g (72%); m.p. 177-178°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 244 (4.15), 324 (4.31), 463 (3.65) nm. – <sup>1</sup>H NMR (acetone-d<sub>6</sub>):  $\delta$  = 3.84 (s, 6 H, CH<sub>3</sub>), 6.98 (s, 2 H, =CH- [dithiine]), 6.68 (2 pseudo-d [J = 2 H], 4 H, aromat. H), 7.01 (2 pseudo-d [J = 2 Hz], 4 H, aromat. H). – <sup>13</sup>C NMR (acetone-d<sub>6</sub>):  $\delta$  = 55.8 (CH<sub>3</sub>), 115.2 (aromat. C<sup>3</sup>), 125.6 (aromat. C<sup>2</sup>), 127.4 (=CH- [dithiine]), 129.7 (aromat. C<sup>1</sup>), 133.6 (>C= [dithiine]), 162 (aromat. C<sup>4</sup>). – MS (70 eV): m/z = 328 (38) [M<sup>+</sup>], 313 (11) [M<sup>+</sup> – CH<sub>3</sub>], 296 (100) [M<sup>+</sup> – S], 281 (96) [M<sup>+</sup> – S – CH<sub>3</sub>],

266 (11)  $[M^+ - S - 2 CH_3]$ , 238 (11)  $[M^+ - S - 2 CH_3 - CO]$ .  $-C_{18}H_{16}O_2S_2$  (328.4): calcd. C 65.82, H 4.91, S 19.52; found C 66.10, H 4.85, S 19.20. – *Method II*: As described for **6h**/method II using 2.21 g (5 mmol) **8m** and yielding first (**Z,Z)-1,4-Di(2-nitrophenyldithio)-1,4-di[p-methoxyphenyl]-1,3-butadiene (9m**). – Yellow leaflets; yield: 2.8 g (89.7%); m.p. 168-172°C (rapid heating ). – MS (70 eV): m/z = 328 (17) [**6m**<sup>+</sup>], 308 (10) [di(2-nitrophenyl) disulfide<sup>+</sup>], 296 (100) [**6m**<sup>+</sup> - S], 154 (65) [di(2-nitrophenyl) disulfide/2<sup>+</sup>]; no molpeak recognizable. – [C<sub>30</sub>H<sub>24</sub>N<sub>2</sub>O<sub>6</sub>S<sub>4</sub> (636.8)]. – Transformation using 637 mg (1 mmol) **9m**; yield of **6m**: 185 mg (57%).

**3,6-Di(p-***n***-butylphenyl)-1,2-dithiine (6n)**. – As described for **6h**/method I using 2.81 g (5 mmol **4n**: Felted purple needles (AcOEt); yield: 875 mg (46%); m.p. 105-108°C (nematic phase at 106°C);  $R_f = 0.76$  (benzene) – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 230 (4.11), 328 (4.30), 460 (3.50) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.93 (t<sub>c</sub>, 6 H, CH<sub>3</sub>), 1.27-1.68 (m, 8 H, CH<sub>2</sub>), 2.63 (t<sub>c</sub>, 4 H, CH<sub>2</sub>), 6.87 (s, 2 H, =CH-), 7.17-7.62 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 13.94 (CH<sub>3</sub>), 22.3, 33.5, 33.5, 35.4 (CH<sub>2</sub>), 125.5, 127.7, 128.9, 142.3, 144.0 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 380 (8) [M<sup>+</sup>], 348 (100) [M<sup>+</sup> – S], 305 (53) [M<sup>+</sup> – C<sub>3</sub>H<sub>7</sub>]. – C<sub>24</sub>H<sub>28</sub>S<sub>2</sub> (380.6): calcd. C 75.74, H 7.42, S 16.85; found C 75.36, H 7.36, S 16.58.

**3,6-Di(p-t-butylphenyl)-1,2-dithime (60).** – As described for **8h**/method I using 2.81 g (5 mmol **40**: Rectangular purple leaflets (AcOEt); yield: 818 mg (43%); m.p. 160-161°C;<sup>39</sup> R<sub>f</sub> = 0.72 (benzene). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.33 (s, 18 H, CH<sub>3</sub>), 6.87 (s, 2 H, =CH-), 7.23-7.64 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 31.2 (CH<sub>3</sub>), 34.7 (>C<), 125.4, 125.7, 127.5, 128.3, 134.0, 143.2 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 380 (48) [M<sup>+</sup>], 365 (16) [M<sup>+</sup> – CH<sub>3</sub>], 348 (100) [M<sup>+</sup> – S], 333 (60) M<sup>+</sup> – S – CH<sub>3</sub>]. – C<sub>24</sub>H<sub>28</sub>S<sub>2</sub> (380.6): calcd. C 75.74, H 7.42, S 16.85; found C 75.34, H 7.27, S 16.43.

**3,6-Di(p-cyclohexylphenyl)-1,2-dithime (6p)**. – As described for **6h**/method I using 3.07 g (10 mmol) **4p**: Rectangular purple leaflets (CHCl<sub>3</sub>, also crystals for X-ray analysis<sup>23</sup> [cf. Figure 3]); yield: 104 mg (48%); m.p. 221°C;<sup>39</sup> R<sub>f</sub> = 0.70 (benzene). – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 239 (4.14), 300 (3.93), 465 (3.26) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.21-1.84 (m, 22 H, C<sub>6</sub>H<sub>11</sub>), 6.85 (s, 2 H, =CH-), 7.21-7.81 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 26.1, 26.8, 34.3 (CH<sub>2</sub>), 44.4 (>CH), 125.4, 127.2, 127.8, 134.4, 149.1 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 432 (8), [M<sup>+</sup>], 399 (100) [M<sup>+</sup> – SH], 357 (15) [M<sup>+</sup> – S – C<sub>3</sub>H<sub>7</sub>], 331 (10) [M<sup>+</sup> – S – C<sub>5</sub>H<sub>9</sub>]. – C<sub>28</sub>H<sub>32</sub>S<sub>2</sub> (432.7): calcd. C 77.73. H 7.45, 14.82; found C 77.22, H 7.35, S 14.52.

**3,6-Di[2-(5,6,7,8-tetrahydronaphthyl]-1,2-dithiine (6q)**. – As described for **6h**/method I using 2.79 g (5 mmol) **4q**: Rectangular purple leaflets (AcOEt); yield: 979 mg (52%); m.p.  $165^{\circ}C$ ;<sup>39</sup> R<sub>f</sub> = 0.83 (benzene). – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\epsilon$ ) = 239 (4.14), 300 (3.93), 465 (3.26) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.53-2.91 (m, 16 H, CH<sub>2</sub>), 6.83 (s, 2 H, =CH-), 7.09-7.40 (m, 6H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 23.1, 29.3, 29.5 (CH<sub>2</sub>), 124.8, 125.2, 128.6, 129.4, 134.1, 134.4, 137.5, 139.4 (aromat. H, =CH-, >C=). – MS (70 eV): m/z = 376 (7) [M<sup>+</sup>], 344 (15) [M<sup>+</sup> – S], 256 (100) [S<sub>8</sub><sup>+</sup>]. – C<sub>24</sub>H<sub>24</sub>S<sub>2</sub> (376.6): calcd. C 76.55, 6.42, S 17.03; found C 76.36, H 6.47, S 17.42.

**3,6-Di(p-biphenyly)-1,2-dithime (6r;** cf. ref.<sup>1d</sup>). – As described for **6h**/method I using 3.01 g (5 mmol) **4r**; major amount of unsoluble solid after evaporation of NH<sub>3</sub>: Reddish brown needles (dissolution in benzene and reprecipitation by AcOEt); yield: 533 mg (25%); m.p. 229-230°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 277, 350, 460 nm (poorly soluble). – <sup>1</sup>H NMR (DMSO-d<sub>6</sub>):  $\delta$  = 6.88 (2 H, =CH-), 7.39-7.85 (m, 18 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>; 12 h accumulated):  $\delta$  = 126.0, 127.1, 127.4, 127.7, 128.2, 128.9 (aromat. C, =CH-), 134.4, 135.7, 140.3, 141.7 (>C=). – MS (70 eV): m/z = 420 (18) [M<sup>+</sup>], 388 (100) [M<sup>+</sup> – S], 354 (5) [M<sup>+</sup> – S – H<sub>2</sub>S], 195 (65) [M<sup>+</sup> – S – C<sub>6</sub>H<sub>5</sub>-C<sub>6</sub>H<sub>4</sub> – CSH]. – C<sub>28</sub>H<sub>20</sub>S<sub>2</sub> (420.6): calcd. C 79.96, H 4.79, S 15.25; found C 79.32, H 4.99, S 14.86.

**3,6-Di(1-naphthyl)-1,2-dithiine** (6s; cf. ref.<sup>1d)</sup>. – As described for 6h/method I using 2.75 g (5 mmol) 4s: Orange red prisms (coformed thiophene 12s was precipitated by addition of AcOEt to the solution of the crude product in few benzene and precipitating with AcOEt (isolation of 6s from the mother liquor); yield: 655 mg (35%); m.p. 148-149°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (Ig  $\varepsilon$ ) = 317 (3.94), 453 (3.06) nm. – <sup>1</sup>H NMR (DMSO-d<sub>6</sub>):  $\delta$  = 6.71 (s, 2 H, =CH-), 7.36-8.55 (m, 14 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>; 12 h accumulated):  $\delta$  = 125.3, 126.1, 126.3, 126.6, 127.4, 128.5, 129.4, 130.2 (aromatic C, =CH- [dithiine]), 131.1, 132.8, 133.9, 136.2 (>C=). – MS (70 eV): m/z = 368 (31) [M<sup>+</sup>], 336 (100) [M<sup>+</sup> – S], 302 (16) [M<sup>+</sup> – S – H<sub>2</sub>S]. – C<sub>24</sub>H<sub>16</sub>S<sub>2</sub> (368.5): calcd. C 78.22, H 4.38, S 17.40; found C 77.95, H 4.48, S 17.56.

**3.6-Di(2-thienyl)-1,2-dithiine** (6t; cf. ref.<sup>1c</sup>). – As described for 6h/method I using 2.31 g (5 mmol) 4t: Violet needles (AcOEt); yield: 101 mg (72%); m.p. 122-123°C.<sup>39</sup> – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 252 (3.87), 339 (4.23), 477 (3.79). – <sup>1</sup>H NMR (DMSO-d\_6):  $\delta = 6.83$  (s, 2 H, =CH-), 7.03-7.07 (m, 2 H, thienyl-H), 7.31-7.35 (m, 4 H, thienyl-H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 124.0$  (=CH- [dithiine]), 126.5, 126.9, 127.5 (thienyl-CH), 127.9 (>C= [dithiine]), 141.1 (>C= [thienyl]). – MS (70 eV): m/z = 280 (30) [M<sup>+</sup>], 248 (100) [M<sup>+</sup> – S], 214 (10) [M<sup>+</sup> – S – H<sub>2</sub>S], 203 (20) [M<sup>+</sup> – S – H<sub>2</sub>S – CSH]. – C<sub>12</sub>H<sub>8</sub>S<sub>4</sub> (280.4): calcd. C 51.39, H 2.88, S 45.73; found C 51.54, H 2.88, S 46.00.

Access to 1,3-butadiene-1,4-dithiols 10 and 1,4-di(acetylthio)-1,3-butadienes 11

A) By reductive fission of (Z,Z)-1,4-Di(benzylthio)-1,3-butadienes 4 via 1,4-dithiolates 5: (Z,Z)-1.3-Butadiene-1,4dithiol (10a; cf. ref.<sup>1a,c,f</sup>). – A mixture of 40 g (134 mmol) 4a and 1 l liquid ammonia was cautiously treated with Na (maximum 15.4 g [670 mmol]) at -70°C over a 2 h period and then with NH<sub>4</sub>Cl until the blue colour of the solution was discharged. After evaporation of the solvent, the product was dissolved in 500 ml water and washed repeatedly with Et<sub>2</sub>O. The resulting aqueous solution of 5a was covered with 200 ml of Et<sub>2</sub>O, and acidified at -20°C with 25% aqueous HCl. The aqueous phase was extracted several times with Et<sub>2</sub>O, the combined extracts dried (Na<sub>2</sub>SO<sub>4</sub>), the solvents removed at reduced pressure (receiver at -70°C) and the residue distilled under vacuum (reveiver at -70°C). - Pale yellow lachrymatory liquid with a highly disagreeable odour; yield: 8.2 g 52%; b.p.  $38^{\circ}C/0.3$  Torr. (m.p.  $\approx -30^{\circ}C) - {}^{1}H$  NMR  $(CDCl_3)$ :  $\delta = 2.92$  (d, 2H SH; J = 9 Hz), 6.17 (m, 2H; =CHSH), 6.41 (d, 2H, -CH=CHSH; J = 8.3 Hz). - <sup>13</sup>C NMR  $(CDCl_3): \delta = 118.3 (=CHSH), 122.8 (HC=C). - MS (70 eV): m/z = 118 (46) [M^+], 116 (35) [M^+ - 2 H], 85 (100) [M^+ - 2 H], 85 (100$ SH], 84 (48)  $[M^+ - H_2S]$ , 71 (32)  $[M^+ - H_2S - CH]$ , 58 (23)  $[M^+ - H_2S - 2 CH]$ ,  $-C_4H_6S_2$  (118.2); calcd. C 40.64, H 5.12, S 54.24; found C 40.63, H 5.46, S 55.05; molar weight (cryoscopic in benzene: 126). - Conversion into (Z,Z)-1,4di(acetylthio)-1,3-butadiene (11a; cf. ref.<sup>1a,c,f</sup>): A) An aqueous solution of 5a (obtained as above) was treated dropwise at 0°C with Ac<sub>2</sub>O until precipitation of the product took place (acidification). - B) A solution, prepared from 1.18 g (0.01 mol) 10a and 15 ml 2N NaOH, was treated with Ac<sub>2</sub>O at 0°C as described above. - Colourless needles (EtOH/AcOEt [1:1]); yield: 19 g (70%); m.p. 147-148°C. – IR (KBr): $\tilde{v} = 1700 \text{ cm}^{-1}$  (C=O). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1000 \text{ cm}^{-1}$ 2.40 (s, 6 H, CH<sub>3</sub>), 6.33-6.37 (2 d, 2 H, =CH-CH=;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz,  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz),  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz),  $J^2 = 8.75$  Hz), 6.74-6.78 (2 d, 2 H, =CH-SCOCH<sub>3</sub>;  $J^1 = 2.1$  Hz),  $J^2 = 8.75$  Hz), J^2 = 8.75 Hz),  $J^2 = 8.75$  Hz),  $J^2 = 8.75$  Hz), J^2 = 8.75 Hz), J^2 = 8.75 Hz),  $J^2 = 8.75$  Hz), J^2 = 8.75 Hz), J^2 = 8.75 Hz), J^2 = 8.75 Hz),  $J^2 = 8.75$  Hz), J^2 = 8.75 Hz), J^2 = 8. 2.1 Hz,  $J^2 = 8.76$  Hz); cf. data in ref.<sup>1f. - 13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 30.8$  (CH<sub>3</sub>), 121.0 (=CH-), 124.8 (=CH-S), 190.4 (C=O). - MS (70 eV): m/z = 202 (10) [M<sup>+</sup>], 160 (10) [M<sup>+</sup> - COCH<sub>2</sub>], 118 (30) [M<sup>+</sup> - 2 COCH<sub>2</sub>], 85 (24) [M<sup>+</sup> - 2 COCH<sub>2</sub> - HS], 84 (21) [M<sup>+</sup> - 2 COCH<sub>2</sub> - H<sub>2</sub>S], 43 (100) [COCH<sub>3</sub><sup>+</sup>]. - C<sub>8</sub>H<sub>10</sub>O<sub>2</sub>S<sub>2</sub> (202.3): calcd. C 47.50, H 4.98, S 31.70; found C 47.56, H 4.98, S 31.67.

(*Z*,*Z*)-1,4-Diphenyl-1,3-butadiene-1,4-dithiol (10h; cf. ref.<sup>1c</sup>). – A 2N NaOH-solution (containing 5h), obtained by the procedure for 6h/method I, was covered with 60 ml Et<sub>2</sub>O and carefully acidified with 25% HCl-solution; the aqueous solution was washed twice with 20 ml Et<sub>2</sub>O and the combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated. The residue was recrystallized from petroleum ether: Pale yellow needles; yield: 1.11 g (82%); m.p. 99-100°C.<sup>39</sup> – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.13 (s; 2H, SH), 6.89 (s, 2H; =CH-), 7.25-7.72 (m, 10H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 125.4, 127.1, 128.5, 128.5 (aromat. C), 134.0, 142.5 (>C=); cf. data in ref.<sup>1f</sup>. – MS (70 eV): m/z = 270 (9) [M<sup>+</sup>], 237 (100) [M<sup>+</sup> – SH], 236 (89) [M<sup>+</sup> – H<sub>2</sub>S], 204 (25) [M<sup>+</sup> – H<sub>2</sub>S – S], 202 (15) [M<sup>+</sup> – 2 H<sub>2</sub>S]. – C<sub>16</sub>H<sub>14</sub>S<sub>2</sub> (270.4): calcd. C 71.06, H 5.22, S 23.71; found C 71.30, H 5.22, S 23.62. – Conversion into (*Z*,*Z*)-1,4-di(acetylthio)-1,4-diphenyl-1,3-butadiene (11h; cf. ref.<sup>1c</sup>): The above noted alkaline solution of 5b was stirred and treated dropwise with AcCl until neutral: Colourless needles (benzene); yield: 1.10 g (62%); m.p. 158-159°C. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.37 (s, 6 H, CH<sub>3</sub>), 7.48 (s, 2 H, =CH-), 7.32-7.62 (m, 10 H, aromat. H). – IR (KBr):  $\overline{v}$  = 1690 cm<sup>-1</sup>. – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 30.5 (CH<sub>3</sub>), 127.2, 128.5, 128.7, 134.5 (aromat. C, =CH-), 134.8, 140.36 (>C=), 192.4 (C=O); cf. data in ref.<sup>1f</sup>. – MS (70 eV): m/z = 354 (5) [M<sup>+</sup>], 311 (28), [M<sup>+</sup> – COCH<sub>3</sub>], 268 (32) [M<sup>+</sup> – 2 COCH<sub>3</sub>], 236 (100) [M<sup>+</sup> – 2 COCH<sub>3</sub> – S]. – C<sub>20</sub>H<sub>18</sub>O<sub>2</sub>S<sub>2</sub> (354.5): calcd. C 67.76, H 5.12, S 18.09; found 67.60, H 5.20, S 18.12.

(Z,Z)-1,4-Di(p-tolyl)-1,3-butadiene-1,4-dithiol (10k; cf. ref.<sup>1c</sup>). – As described for 10h using 2.39 g (5 mmol) 4k: Pale yellow leaflets (Et<sub>2</sub>O); yield: 925 mg (62%); m.p. 137-138°C<sup>39</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 2.35 (s, 6 H, CH<sub>3</sub>), 3.07 (s, 2 H, SH), 6.83 (s, 2 H, =CH-), 7.15-7.54 (2 d, 8 H, aromat. H; J = 8.2 Hz). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 21.2 (CH<sub>3</sub>), 124.9, 127.0, 129.1, 133.5, 138.5, 139.7 (aromat. C, =CH-). – MS (70 eV): m/z = 298 (3) [M<sup>+</sup>], 264 (100) [M<sup>+</sup> – H<sub>2</sub>S], 249 (5) [M<sup>+</sup> – H<sub>2</sub>S - CH<sub>3</sub>]. – C<sub>18</sub>H<sub>18</sub>S<sub>2</sub> (298.5): calcd. C 72.47, H 6.08, S 21.49; found C 72.47, H 6.08, S 21.52. – Conversion into (*Z,Z*)-1,4-di(acetylthio)-1,4-di(p-tolyl)-1,3-butadiene (11k; cf. ref.<sup>1c</sup>) as described for 11h: Colourless needles (benzene); m.p. 192-193°C. – IR (KBr): $\tilde{v}$  = 1730 cm<sup>-1</sup>. – C<sub>22</sub>H<sub>22</sub>O<sub>2</sub>S<sub>2</sub> (382.6): calcd. C 69.07, H 5.79; found C 69.33, H 5.66.

(Z,Z)-1-(p-Methoxyphenyl)-4-phenyl-1,3-butadiene-1,4-dithiol (10l). – As described for 10h using 2.4 g (5 mmol) 4l: Pale yellow needles (petroleum ether in the presence of some NaBH<sub>4</sub>); yield: 901 mg (60%); m.p. 97-98°C<sup>39</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.80 (s, 2 H, SH), 3.82 (s, 3H, OCH<sub>3</sub>), 6.84-6.90 (2 d, 2 H, =CH-; J = 2.76 Hz), 7.31-7.62 (m, 9 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 55.4 (OCH<sub>3</sub>), 124.2, 125.7, 127.1, 127.7, 128.3, 128.4, 132.9 (aromat. C, =CH-), 132.9, 133.7, 134.9, 142.5, 160.0 (>C=). – MS (70 eV): m/z = 300 (4) [M<sup>+</sup>], 266 (100) [M<sup>+</sup> – H<sub>2</sub>S], 251 (77) [M<sup>+</sup> – H<sub>2</sub>S – CH<sub>3</sub>]. – C<sub>17</sub>H<sub>16</sub>OS<sub>2</sub> (300.4): calcd. C67.96, H 5.37, S 21.34; found C 68.06, H 5.29, S 21.29. – Conversion into (Z,Z)-1,4-di(acetylthio)-1-(p-methoxyphenyl)-4-phenyl-1,3-butadiene (111) as described for 11h: Pale yellow needles (benzene); m.p. 132-133°C. – IR (KBr): $\tilde{v}$  = 1710 cm<sup>-1</sup>. – C<sub>21</sub>H<sub>20</sub>O<sub>3</sub>S<sub>2</sub> (384.5): calcd. C 65.59, H 5.24; found C 65.92, H 5.58.

(Z,Z)-1,4-Di(p-methoxyphenyl)-1,3-butadiene-1,4-dithiol (10m; cf. ref.<sup>1c</sup>). – As described for 10h using 2.55 g (10 mmol) 4m: Pale yellow needles (petroleum ether); yield: 122 mg (74%); m.p. 147-148°C<sup>39</sup>. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.05 (s, 2 H, SH), 3.82 (s, 6 H, OCH<sub>3</sub>), 6.79 (s, 2 H, =CH-), 6.87-7.59 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 55.4 (OCH<sub>3</sub>), 113.8, 124.5, 128.3 (aromat. C, =CH-), 132.6, 135.0 (>C=), 159.9 (C-OCH<sub>3</sub>). – MS (70 eV): m/z = 330 (5) [M<sup>+</sup>], 296 (100) [M<sup>+</sup> – H<sub>2</sub>S], 281 (93) [M<sup>+</sup> – H<sub>2</sub>S – CH<sub>3</sub>], 266 (18) [M<sup>+</sup> – H<sub>2</sub>S – 2 CH<sub>3</sub>]. – C<sub>18</sub>H<sub>18</sub>O<sub>2</sub>S<sub>2</sub> (330.5): calcd. C 65.42, H 5.49, S 19.41; found C 65.33, H 5.45, S 19.50. – Conversion into (Z,Z)-1,4-di(acetylthio)-1,4-di(p-methoxyphenyl)-1,3-butadiene (11m; cf. ref.<sup>1c</sup>) as described for 11h: Pale yellow needles (benzene); m.p. 166-167°C. – IR (KBr):  $\tilde{v}$  = 1695 cm<sup>-1</sup>. C<sub>22</sub>H<sub>22</sub>O<sub>4</sub>S<sub>2</sub> (414.6): calcd. C 63.72, H 5.35; found C 63.30, H 5.31.

B) By reduction of 1,2-dithilnes 6: A solution of 1 mmol 6 in 5ml benzene/20ml EtOH/1ml H<sub>2</sub>O was treated at room temperature with 380 mg (10 mmol) NaBH<sub>4</sub>. The colour of the reaction mixture changed from red to yellow. The mixture was carefully acidified with 2N H<sub>2</sub>SO<sub>4</sub>, 15 ml benzene added and the organic phase recovered and concentrated. The residue was recrystallized from petroleum ether, isolated yields  $\approx 85\%$  (examples: 10h,k,l,m).

#### Transformation to thiophenes 12.

A) By action of acids on 4 (cf. ref.<sup>1e</sup>): A solution of 3 mmol 4 in 20 ml dioxane was treated with a current of dry HCl gas for 2 h. The mixture was allowed to stand overnight, the solvents evaporated and the residue recrystallized from EtOH or AcOH. - Examples: 2,5-Diphenylthiophene (12h), 65%, m.p. 152-153°C; 2-phenyl-5-(p-tolyl)thiophene (12j), 63%, m.p. 148-149°C; 2,5-di(p-tolyl)thiophene (12k), 61%, m.p. 173-174°C; 2-phenyl-5-(pmethoxyphenyl)thiophene (121), 52%, m.p. 165-166°C, 2,5-di(p-methoxyphenyl)thiophene (12m), 50%, m.p. 218-219°C. – The same reaction with 3k yields 55% 12k. – B) By sulfur extrusion from 6: A solution of 100 mg 6 in 30 ml AcOEt was exposed for 4-5 h at room temperature to day-light whereupon complete decolouration occurs (cf. also the thermal sulfur extrusion, e.g in DMSO above 100°C in »Results and Discussion« and ref.<sup>28</sup>). After evaporation the residue was recrystallized in order to separate the product from coformed sulfur; Yields almost quantitative. Examples: **2,5-Di(p-***n***-butylphenyl)thiophene (12n)**, m.p. 142°C (no occurrence of a nematic phase in opposite to **6n**),  $R_f = 0.79$ (silica gel, benzene); UV/ Vis (MeCN):  $\lambda_{max}$  (lg  $\epsilon$ ) = 233 (4.11), 327 (4.40) nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 0.93 (t<sub>c</sub>, 6 H, CH<sub>3</sub>), 1.24-1.45 (m, 4 H, CH<sub>2</sub>), 1.53-1.68 (m, 4 H, CH<sub>2</sub>), 2.61 (t<sub>c</sub>, 4 H, CH<sub>2</sub>), 7.15-7.54 (m, 10 H, aromat. H, =CH-); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 13.9 (CH<sub>3</sub>), 22.3, 35.3, 37.5 (CH<sub>2</sub>), 125.4, 127.3, 128.2, 131.5, 138.7, 143.3 (aromat. C, =CH-, >C=); MS (70 eV): m/z = 348 (100) [M<sup>+</sup>], 306 (43) [M<sup>+</sup> - C<sub>3</sub>H<sub>6</sub>]; C<sub>24</sub>H<sub>28</sub>S (348.5): calcd. C 82.70, H 8.10, S 9.20; found H 82.91, H 8.43, S 9.36. -2,5-Di(p-t-butylphenyl)thiophene (120), m.p. 164°C,  $R_f = 0.77$  (silica gel, benzene); UV/ Vis (MeCN):  $\lambda_{max}$  (lg  $\epsilon$ ) = 235 (4.15), 326 (4.47) nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.33 (s, 18 H, CH<sub>3</sub>), 7.22-7.57 (m, 10 H, aromat, H, =CH-); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 31.3 (CH<sub>3</sub>), 34.6 (>C<), 123.5, 125.3, 125.8, 131.6, 143.2, 146.9 (aromat. C, =CH-, >C=); MS (70 eV): m/z = 348 (100) [M<sup>+</sup>], 333 (55) [M<sup>+</sup> - CH<sub>3</sub>]; C<sub>24</sub>H<sub>28</sub>S (348.5): calcd. C 82.70, H 8.10, S 9.20; found C 82.71, H 7.84, S 9.24. - 2,5-Di(p-cyclohexylphenyl)thiophene (12p), m.p. 233°C, Rf = 0.86 (silica gel, benzene); UV/ Vis (MeCN):  $\lambda_{max}$  (lg  $\epsilon$ ) = 232 (4.20), 330 (4.44) nm; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.30-2.50 (m, 22 H, C<sub>6</sub>H<sub>1</sub>), 7.19-7.55 (m, 10 H, aromat. H, =CH-); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta \approx 26.2, 26.9, 34.4$  (CH<sub>2</sub>), 44.3, (>CH-), 123.4, 125.6, 127.3, 132.0, 143.3, 147 .5, (aromat. C, =CH-, >C=); MS (70 eV):  $m/z = 400 (100) [M^+]$ , 331 (13)  $[M^+ - C_5H_9]$ ; C28H32S (400.6): calcd. C 83.94, H 8.05, S 8.01; found C 83.87, H 7.98, S 8.06. - 2,5-Di[2-(5,6,7,8-tetrahydronaphthyl]thiophene (12q), m.p 164°C,  $R_f = 0.77$  (silica gel, benzene); UV/ Vis (MeCN):  $\lambda_{max}$  (lg  $\epsilon$ ) = 231 (4.28), 328 (4.54); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 1.22-1.51 (m, 16 H, C<sub>4</sub>H<sub>8</sub>), 7.23-7.63 (m, 8 H, aromat. H, =CH-); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.4, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.6, 30.8, 36.7$  (CH<sub>2</sub>), 127.0, 127.3, 128.3, 128.6, 128.8, 129.3, 137.8, 149.9 (aromat. C, =CH-, CDCl<sub>3</sub>):  $\delta = 22.0, 30.8, 36.7$  (CH<sub>2</sub>):  $\delta = 22.0, 30.8$ >C=); MS (70 eV): m/z = 344 (100) [M<sup>+</sup>], 310 (17) [M<sup>+</sup> - H<sub>2</sub>S]; C<sub>24</sub>H<sub>24</sub>S (344.5): calcd. C 83.67, H 7.03, S 9.30; found C 83.07, H 7.70, S 9.81.

C) By thermolysis of 10 (H<sub>2</sub>S-elimination): Heating of 10 for several minutes at the melting point or refluxing a solution of 10 in a high-boiling solvent (e.g. xylene) for 5-10 min gave almost quantitative yields (examples: 12h,j). Alkylation of 5 and 10 (11)

A) The alkaline aqueous solution of 5 or 10, obtained from the reductive debenzylation of 4 (described in the procedure for 6h) was diluted with the equal amount EtOH and subsequently treated with two molar equivalents of the organic halide (benzyl chloride, methyl iodide, ethyl iodide). The precipitated products (4, 13, 14) were filtered by suction and recrystallized; yield: 70-90%. - B) 10 mmol 11 were dissolved (saponified) in a solution of 3.4 g (60 mmol) KOH in 20 ml MeOH, after addition of the organic halide at elevated temperature ( $\approx 40^{\circ}$ C) the products precipitate; yield: 80-95%. - (Z,Z)-1,4-Di(methylthio)-1,4-diphenyl-1,3-butadiene (13h), m.p. 134°C (EtOH/AcOEt [1:1]), Rf = 0.68 (benzene); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 2.03$  (s, 6 H, CH<sub>3</sub>), 7.23-7.63 (m, 12 H, aromat. H, =CH-); <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 16.4$  (CH<sub>3</sub>), 128.0, 128.1, 128.4, 129.4, 139.6. 140.9 (aromat. C, =CH-, >C=); MS (70 eV): m/z = 298 (21) [M<sup>+</sup>], 251 (56) [M<sup>+</sup> -CH<sub>3</sub>S], 236 (85) [M<sup>+</sup> – CH<sub>3</sub>S – CH<sub>3</sub>], 202 (51) [M<sup>+</sup> – CH<sub>3</sub>S – CH<sub>3</sub> – H<sub>2</sub>S], 121 (100) [C<sub>6</sub>H<sub>5</sub>CS<sup>+</sup>], 77 (81) [C<sub>6</sub>H<sub>5</sub><sup>+</sup>]; C18H18S2 (298.4): calcd. C 72.44, H 6.07, S 21.49; found C 72.41, H 6.19, S 21.55; identical with product obtained by thiol addition to 1h using 16.1 g (75 mmol) S-methylisothiouronium sulfate according to procedure for 4h/method I (yield: 64%). - Further examples: 13a, b.p. 93-94°C/3 Torr [C<sub>6</sub>H<sub>10</sub>S<sub>2</sub> (146.3)]; 13j, m.p. 140-141°C [C<sub>19</sub>H<sub>20</sub>S<sub>2</sub> (312.5)]; 13k, m.p. 187-188°C [C<sub>20</sub>H<sub>22</sub>S<sub>2</sub> (326.5)]; 13l, m.p. 106-107°C [C<sub>19</sub>H<sub>20</sub>OS<sub>2</sub> (328.5)]; 13m, m.p. 163-164°C [C<sub>20</sub>H<sub>22</sub>O<sub>2</sub>S<sub>2</sub> (358.5)]; 13t, m.p. 111-112°C [C<sub>14</sub>H<sub>16</sub>S<sub>4</sub> (312.5)]. - (Z,Z)-1,4-Di(ethylthio)-1,4-diphenyl-1,3-butadiene (14h), m.p. 99-100°C (EtOH),  $R_f = 0.72$ ; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 1.09$  (t<sub>c</sub>, 6 H, CH<sub>3</sub>), 2.48 (q<sub>c</sub>, 4 H, CH<sub>2</sub>), 7.32-7.68 (m, 12 H, aromat. H, =CH-);  ${}^{13}$ C NMR (CDCl<sub>3</sub>):  $\delta$  = 15.1 (CH<sub>3</sub>), 27.1 (CH<sub>2</sub>), 128.1, 128.5, 131.5, 138.4, 139.6, 140.3 (aromat. C, =CH-); MS (70 eV): m/z = 326 (10) [M+], 265 (25) [M+ - C<sub>2</sub>H<sub>5</sub>S], 236 (100) [M+ - C<sub>2</sub>H<sub>5</sub>S - C<sub>2</sub>H<sub>5</sub>], 202 (33)  $[M^{+} - C_{2}H_{5}S - C_{2}H_{5} - H_{2}S]$ , 121 (89)  $[C_{6}H_{5}CS^{+}]$ ;  $C_{20}H_{22}S_{2}$  (326.5): calcd. C 73.57, H 6.79, S 19.64; found C 73.96, H 6.78, S 19.36; identical with product obtained by thiol addition to 1h using 27.75 g (150 mmol) S-ethylisothiouronium bromide according to procedure for 4h/method I (yield: 76%). - Further examples: 14a, b.p. 95-96°C/0.9 Torr [C<sub>8</sub>H<sub>12</sub>S<sub>2</sub> (174.3)]; 14j, m.p. 96-97°C [C<sub>21</sub>H<sub>24</sub>S<sub>2</sub> (340.6)]; 14k, m.p. 140-141°C [C<sub>22</sub>H<sub>26</sub>S<sub>2</sub> (354.6)]; 14l, m.p. 7678°C [C<sub>21</sub>H<sub>24</sub>OS<sub>2</sub> (356.6)]; **14m**, m.p. 113-114°C [C<sub>22</sub>H<sub>26</sub>O<sub>2</sub>S<sub>2</sub> (386.6)]; **14t**, m.p. 108-109°C [C<sub>16</sub>H<sub>18</sub>S<sub>4</sub> (338.6)]. – Compounds **14h**, **14j**, **14k**, **14l**, **14m** and **14t** were also produced by reaction of the corresponding 1,2-diaroylethenes with EtSH under acidic conditions (1,4-dioxane/HCl; p-toluenesulfonic acid,  $\Delta$ ) as previously described in ref.<sup>1e</sup>.

(*Z,Z,Z,Z*)-1,4-Diphenyl-1,4-di(β-styrylthio)-1,3-butadiene (15). – A solution of 540 mg (2 mmol) 10h, 225 mg (4 mmol) KOH and 408 mg (4 mmol) phenylacetylene in 6 ml MeOH was refluxed 7 h. Oily droplets gradually separated and crystallized. The hot methanolic solution was decanted and the residue after washing with H<sub>2</sub>O and MeOH recrystallized from AcOEt/EtOH (1:1). – Yellow needles; yield: 500 mg (52%); m.p. 159-160°C. – UV/Vis (EtOH):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 257 (4.47), 292 (4.55), 381 (4.53) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 6.02 (d, 2 H), 6.37 (d, 2 H; H<sup>α</sup>, H<sup>β</sup>; J = 10.8 Hz), 7.23 (m, 2H, aromat. H), 7.31 (s, 2 H, -CH=C[S]-C<sub>6</sub>H<sub>5</sub>), 7.36 (m, 10 H, aromat. H), 7.50, 7.54, 7.61 (m, 8 H, aromat. H). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 125.1, 126.4, 127.1 (=CH-), 128.4, 128.6, 128.7, 129.5 (aromat. C), 136.4, 139.3, 139.4 (>C=). – MS (70 eV): m/z = 474 (40) [M<sup>+</sup>], 339 (100) [M<sup>+</sup> – C<sub>6</sub>H<sub>5</sub>CHCHS], 261 (44) [M<sup>+</sup> – C<sub>6</sub>H<sub>5</sub>CHCHS – C<sub>6</sub>H<sub>5</sub>], 121 (31) [C<sub>6</sub>H<sub>5</sub>CS<sup>+</sup>]. – C<sub>32</sub>H<sub>26</sub>S<sub>2</sub> (474.7): calcd. C 80.97, H 5.52, S 13.51; found C 81.10, H 5.73, S 13.89.

7,10-Diphenyl-3,4-benzo-1,6-dithiacyclododeca-3,7,9-triene (16). – A solution of 840 mg (15 mmol) KOH in 4 ml MeOH and 300 ml DMF was treated dropwise with solutions of 900 mg (2.5 mmol) 11h in 50 ml DMF and of 660 mg (2.5 mmol) o-di(bromomethyl)benzene in 50 ml DMF at 55°C with stirring under an argon atmosphere. After further stirring at this temperature for 3 h, the solvents were evaporated, the residues dissolved in 50 ml water and the solution extracted three times with 30 ml portions of CH<sub>2</sub>Cl<sub>2</sub>. The combined extracts were concentrated and the residue was recrystallized from AcOEt. – Pale yellow leaflets; yield: 100 mg (11%); m.p. 142°C; R<sub>f</sub> = 0.67 (silica gel, benzene). – UV/Vis (MeCN):  $\lambda_{max}$  (lg  $\varepsilon$ ) = 268 (4.21), 343 (4.25) nm. – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  = 3.44 (s, 4 H, CH<sub>2</sub>), 7.09-7.22 (m, 16 H, aromat. H, =CH-). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta$  = 36.5 (CH<sub>2</sub>), 126.9, 127.2, 128.2, 130.7, 134.7, 138.0, 138.5, 139.9 (aromat. C, =CH-, >C=). – MS (70 eV): m/z = 273 (10) [M<sup>+</sup>], 236 (96) [C6<sub>4</sub>5<sub>5</sub>CCHCHCSC<sub>6</sub>H<sub>5</sub>+], 136 (68) [SCH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub><sup>+</sup> (dihydroisothionaphthene)], 121 (100) [C<sub>6</sub>H<sub>5</sub>CS<sup>+</sup>], 104 (69) [CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub><sup>+</sup>]. – C<sub>2</sub>H<sub>2</sub>OS<sub>2</sub> (372.5): calcd. C 77.38, H 5.41, S 17.21; found C 77.28, H 5.22, S 17.23. – From the mother liquor 12h (32%) and dihydroisothionaphthene were isolale (TLC, R<sub>f</sub> = 0.59 [silica gel, benzene]).

*Reductive defunctionalization of* **14h** *to* (*E*)-**1,4-Diphenyl-2-butene** (**18**): 6.53 g (20 mmol) **14h** was treated with 2.0 g (87 mmol) Na in 120 ml liquid NH<sub>3</sub> at -70°C and worked up as described for **6h**/method I. After addition with 100 ml 2N NaOH to the evaporation residue, the precipitated white solid was removed by suction filtration and recrystallized from EtOH (the filtrate shows no reaction with FeCl<sub>3</sub>). – White needles; yield: 2.8 g (67%); m.p. 43-44°C (ref.<sup>40a</sup>: 43-45°C; ref.<sup>40b</sup>: 45.5-46°C); identical with the product obtained from the analogous treatment of (*Z*,*Z*)-1,4-diphenyl-1,3-butadiene (72%). – IR (nujol):  $\tilde{v} = 970 \text{ cm}^{-1. - 1}\text{H NMR (CDCl_3)}$ :  $\delta = 3.35-3.41$  (m, 4 H, CH<sub>2</sub>), 5.64-5.73 (m, 2 H, =CH-), 7.13-7.28 (m, 10 H, aromat. H). – <sup>13</sup>C NMR (CDCl\_3):  $\delta = 39.0$  (CH<sub>2</sub>), 126.0, 128.4, 128.6, 130.5, 140.8 (aromat. C, =CH-). – MS (70 eV): m/z = 208 (36) [M<sup>+</sup>], 117 (100) [C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CHCH<sup>+</sup>], 91 (35) [C<sub>7</sub>H<sub>7</sub><sup>+</sup>]. C<sub>16</sub>H<sub>16</sub> (208.3).

Bromine addition:  $3\alpha_4\beta_5\alpha_6\beta$ -Tetrabromo-3,6-diphenyl-1,2-dithiane (21). – A suspension of 0.27 g (1 mmol) 6h in 4 ml CCl<sub>4</sub> was treated at -5°C with 0.32 g (2 mmol) Br<sub>2</sub>. A clear solution was obtained and a solid precipitated. The solid was collected, washed with ice-cold CCl<sub>4</sub> and quickly dried i. vac. – Buff-coloured crystals; yield: 0.38 g (65%); m.p. 83-84°C (dec., from 77°C formation of gas and colour change to orange red). – <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta = 5.40$  (s, 2 H, >CHBr), 7.47 m<sub>c</sub>, 6 H aromat. H<sub>[m,p]</sub>), 7.88 (m<sub>c</sub>[broad, hindrance of rotation], 4 H, aromat. H<sub>[o]</sub>). – <sup>13</sup>C NMR (CDCl<sub>3</sub>):  $\delta = 65.5$  (>CHBr), 81.8 >CBrPh), 127.7 (aromat. C<sub>[p]</sub>), 128.9, 130.2 (aromat. C<sub>[o,m]</sub>), 138.8 (aromat. C<sub>[i]</sub>). – C<sub>16</sub>H<sub>12</sub>Br<sub>4</sub>S<sub>2</sub> (588.0); storage stability even at -20°C only 3-4 d.

**Condensation reaction to 3,6-disubstituted pyridazines 22.** – A) To a mixture of 3 ml EtOH and 5 ml pyridine (liberated from oxygen by previous heating under an argon atmosphere) was added 0.27 g (1 mmol) **6h** and 0.12 ml (2.4 mmol) 98% hydrazine hydrate. The mixture was heated under exclusion of day-light and oxygen 2 h on a wather bath (yellow brown solution). After chilling **3,6-diphenylpyridazine (22h)** precipitated which was separated and recrystallized from EtOH. – Colourles leaflets; yield: 100 mg (43%); m.p. 223°C (ref.<sup>41a</sup>: 219-220°C); C<sub>16</sub>H<sub>12</sub>N<sub>2</sub> (232.3). – The residues were concentrated and treated with water to afford a solid which was recrystallized from a little EtOH to give 110 mg (47%) 2,5-diphenylthiophene (**12h**) (pale yellow plates; m.p. 148-149°C). – Further examples: **22j** (48%; m.p. 185-186°C [ref.<sup>41a</sup>: 185-186°C]; C<sub>17</sub>H<sub>14</sub>N<sub>2</sub> [246.3]), **22k** (38%; m.p. 233-234°C [ref.<sup>41b</sup>: 231-232°C]; C<sub>18</sub>H<sub>16</sub>N<sub>2</sub> [260.3]), **22l** (45%); m.p. 195-196°C [ref.<sup>41a</sup>: 195-196°C]; **C**<sub>14</sub>H<sub>4</sub>N<sub>2</sub>O [262.3]). – B) As described precedingly with 1 mmol **10** or **11**, 0.5 ml EtOH, 1.5 ml pyridine and 75 mg (1.5 mmol) hydrazine hydrate. Hydrogen sulfide was produced and the reaction was completed by heating on a wather bath for 1 h. The product was obtained by dilution of the solution with water: Yields 75-85% (from 10), 70-75% (from **11**), examples: **22h**, **22k**, **22h**, **22k**, **22h**, **22h**, **22m** (m.p. 236-237°C [ref.<sup>41a</sup>: 234-235°C]; C<sub>18</sub>H<sub>16</sub>ON<sub>2</sub> [276.3]), **22t** (m.p. 178-179°C; C<sub>12</sub>H<sub>8</sub>N<sub>2</sub>S<sub>2</sub> [244.3]).

Acknowledgements: Financial support by the Fonds der Chemischen Industrie, Frankfurt/Main, and the Deutsche Forschungsgemeinschaft is acknowledged. We are very grateful to Dr. Allan Dunn, Frankfurt/Main, for editorial assistance.

## **References and Notes**

## Dedicated to Professor Siegfried Hunig on the occasion of his 75th birthday

- 1. Preceding parts I: a) Schroth, W.; Langguth, H.; Billig, F. Z. Chem. 1965, 5, 352-353. II: b) Schroth, W.; Billig, F.; Langguth, H. ibid. 1965, 5, 353-354. - III: c) Schroth, W.; Billig, F.; Reinhold, G. Angew. Chem. 1967, 79, 685-686; Angew. Chem. Int. Ed. Engl. 1967, 6, 698-699. - IV: d) Schroth, W.; Billig, F.; Zschunke, A. Z. Chem. 1969, 9, 184-185. - V: e) Schroth, W.; Billig, F.: Reinhold, G. Z. Chem. 1969, 9, 229-230. - VI: f) Radeglia, R.; Poleschner, H.; Schroth, W. Z. Naturforsch. 1988, 43b, 605-610. - VII: g) Zschunke, A.; Mügge, C.; Hintzsche, E.; Schroth, W. J. Prakt. Chem. 1992, 334, 141-146. - VIII: h) Schroth, W.; Hintzsche, E.; Viola, H.; Winkler, R.; Klose, H.; Boese, R.; Kempe, R.; Sieler, J. Chem. Ber. 1994, 127, 401-408. - IX: i) Schroth, W.; Hintzsche, E.; Spitzner, R.; Irngartinger, H.; Siemund, V. Tetrahedron Lett. 1994, 35, 1973-1976. - X: j) Schroth, W.; Felicetti, M.; Hintzsche, E.; Spitzner, R.; Pink, M. ibid. 1994, 35, 1977-1980. - XI: k) Schroth, W.; Hintzsche, E.; Felicetti, M.; Spitzner, R.; Sieler, J.; Kempe, R. Angew. Chem. 1994, 106, 808-810; Angew. Chem. Int. Ed. Engl. 1994, 33, 739-741. - XII: 1) Schroth, W.; Jordan, H.; Spitzner, R. Tetrahedron Lett. 1995, 36, 1421-1424. - XIII: m) Schroth, W.; Ströhl, D.; Thondorf, I.; Brandt, W.; Felicetti, M.; Gelbrich, T. Tetrahedron 1995, 51, 8853-8862. - XIV: n) Schroth, W.; Hintzsche, E.; Spitzner, R.; Ströhl, D.; Sieler, J. ibid. 1995, 51, 13247-13260. - XV: o) Schroth, W.; Hintzsche, E.; Spitzner, R.; Ströhl, D.; Schmeiß, K.; Sieler, J. ibid. 1995, 51, 13261-13270.
- 2. a) Ph.D. Thesis, University of Halle 1989 (cf. Billig, F. Ph.D. Thesis, University of Leipzig, 1969).
- 3. Author to whom inquires concerning the X-ray structure analyses should be addressed (cf. ref.<sup>22,23</sup>).
- a) Bohlmann, F.; Kleine, K.-M. Chem. Ber. 1965, 98, 3081-3086. b) Mortensen, J. T.; Sørensen, J. S.; Sørensen, N. A. Acta Chem. Scand. 1964, 18, 2392-2394. - Cf. also review on naturally occurring 1,2dithiines: c) Freeman, F.; Aregullin, M.; Rodriguez, E. Rev. Heteroatom Chem. 1993, 9, 1-19.
- a) Antibiotic properties: Towers, G. H. N.; Abramowski, Z.; Finlayson, A. J.; Zucconi, A. Planta Med. 1985, 225-229. – Antiviral properties: b) Hudson, J. B.; Graham, E. A.; Fong, R.; Finlayson, A. J.; Towers, G. H. N. *ibid.* 1986, 51-54. – Antibacterial and antifungal properties: c) Constabel, C. P.; Towers, G. H. N. *ibid.* 1989, 55, 35-37. – Cf. also d) Rodriguez, E.; Aregullin, M.; Nishidida, T.; Uehara, S.; Wrangham, R.; Abramowski, Z.; Finlayson, A.; Towers, G. H. N. *Experientia* 1985, 41, 419-420.
- 6. a) Koreeda, M.; Yang, W. Synlett 1994, 201-203. Synthesis of thiarubrine A: b) Koreeda, M.; Yang, W. J. Am. Chem. Soc. 1994, 116, 10973-10974. Synthesis of thiarubrine B: c) Block, E.; Guo, C.; Thiruvazhi, M.; Toscano, P. J. J. Am. Chem. Soc. 1994, 116, 9403-9404. Comprehensive substituent variations and investigation of antifungal activity: d) Bierer, D. E.; Dener, J. M.; Dubenko, L. G.; Gerber, R. E.; Litvak, J.; Peterli, S.; Peterli-Roth, P.; Truong, T. V.; Mao, G.; Bauer, B. E. J. Med. Chem. 1995, 38, 2628-2648. Other analogues: e) Freeman, F.; Kim, D. S. H. L.; Rodriguez, E. J. Org. Chem. 1992, 57, 1722-1727. f) Freeman, F.; Kim, D. S. H. L.; Rodriguez, E. ibid. 1993, 58, 2317-2319. Theoretical investigations: g) Borsdorf, R.; Hofmann, H.-J.; Köhler, H.-J.; Scholz, M.; Fabian, J. Tetrahedron 1970, 26, 3227-3231. h) Fabian, J.; Birner, P. Coll. Czech. Chem. Commun. 1988, 53, 2096-2115. i) Cimiraglia, R.; Fabian, J.; Hess jr., B. A. J. Mol. Struct. (Theochem) 1991, 230, 287-293. j) Mann, M.; Fabian, J. ibid. 1995, 331, 51-61. Cf. reviews: k) Eisner, U.; Krishnamurthy, T. Int. J. Sulfur Chem., B, 1972, 7, 101-107; l) Freeman, F.; Kim, D. S. H. L.; Rodriguez, E. Sulfur Rep. 1989, 9, 207-256.
- Cf., for example: a) Küsters, W.; de Mayo, P.; J. Am. Chem. Soc. 1973, 95, 2383-2384. b) Schulz, R.; Schweig, A.; Hartke, K.; Köster, J. *ibid.* 1983, 105, 4519-4528. - c) Diehl. F.; Meyer, H.; Schweig, A. *ibid.* 1989, 111, 7651-7653. - d) Yu, H.; Chan, W. T.; Goddart, J. D. *ibid.* 1990, 112, 7529-7537.
- Cf. exceptions according to Scheme 6 and in: a) Behringer, H.; Meinetsberger, E. Liebigs Ann. Chem. 1981, 1729-1750; ibid. 1981, 1928-1959; ibid. 1982, 315-341. - b) Moran, J. R.; Huisgen, R.; Kalwinsch, I. Tetrahedron Lett. 1985, 26, 1849-1852.
- A resemblance of this approach to the presumed biosynthesis of the naturally occurring C<sub>13</sub>-1,2-dithiines should be emphasized; namely by an analogous 1,4-bis-addition of *natural* sulfur suppliers at a butadiyne-subunit within a C<sub>13</sub>-polyyne but in a different orientation within this chain; cf.: a) Bohlmann, F. Fortschr. Chem. Org. Naturst. 1967, 1-62, especially 9-10. – b) Bohlmann, F.; Zdero, C. Chem. Heterocycl. Compds. (Weissberger, A., ed.) 1985, 44, Part 1, 261-324, especially 299-300. – c) Bohlmann, F.; Burkhardt, T.; Zdero, C. Naturally Occurring Acetylenes; Academic Press, London, New York: 1973, especially 61, 65-66. – d) ref.<sup>4c</sup>, p. 2-3. – Here, the terminal positions of the polyyne chain are generally

avoided, whereas in the nucleophilic thiol additions these positions are in principle preferred (cf. ref. 1g).

- Dshemilew, U. M.; Selimow, F. A.; Chafisow, W. R.; Chalikow, L. M.; Tolstikow, G. A. *Izv. Akad.* Nauk SSSR 1986, 1211-1212. – Dshemilew, U. M.; Baibulatowa, N. S.; Tkatschenko, T. K.; Kumakowa, R. W. *ibid.* 1987, 1918. – Distillable non-coloured oil without any extrusion of sulfur, obtained by a cobalt catalyzed reaction of 1-pentyne with carbon disulfide or propyne with sulfur/butyl chloride under vigorous conditions (150°C, 6 h). In accord with the indication of that two isomers 2,4- and 2,5-dibutyl-1,4-dithiines should be present.
- Cf. Review: a) Shostakovskii, M. F.; Bogdanova, A. V. The Stereochemistry of Diacetylenes, John Wiley & Sons, New York, Toronto, 1974, especially 109-118. Cf. deviations from this regiochemistry in the reaction with amines: b) Schroth, W.; Peschel, J.; Zschunke, A. Z. Chem. 1969, 9, 108-109, 110-111, 143.
- a) Truce, W. E.; Simms, J. A. J. Am. Chem. Soc. 1956, 78, 2756-2759. b) Baliak, V.; Rathinasamy, T. K. Indian J. Chem. 1971, 9, 220-225.
- a) Freeman, F.; Lu, H.; Zeng, Q. J. Org. Chem. 1994, 59, 4350-4354. b) Furthermore, the use of LiAlH<sub>4</sub> in THF seems to be a convenient method in the aromatic substituted series: Freeman, F.; Lu, H.; Rodriguez, E. Sulfur Lett. 1995, 18, 243-257.
- 14. Pastuszak, J. J.; Chimiak, A. J. Org. Chem. 1981, 46, 1868-1873.
- 15. a) Cf. also information in ref.<sup>1a</sup>. b) As already pointed out, <sup>1n</sup> the predominance of the dithiols **10** over the butan-1,4-dithione tautomers, in contrast to the situation in the oxygen-series, represents an impressive example for the disadvantage of  $p_z$ -overlapping between the sulfur atom with the adjacent carbon atom.
- 16. a) The reaction of 2,2'-dimercaptobiphenyl with the same dichloro-C<sub>2</sub>-building blocks affords 5,6,7,8-dibenzo-1,4-dithiocine whereas the reaction of (Z)-ethene-1,2-dithiol or o-xylene-4,5-dithiol with diacetylene yields the corresponding 2-vinyl-1,4-dithiines via a 1,2-orientation of the bis-addition at the C<sub>4</sub>-component, as already reported in ref.<sup>1d</sup>. On the other hand, the reaction of (Z)-1,2-ethenedithiol with (Z)-1,2- or 1,1-dichloroethene gives 1,4-dithiine without any problems: Schroth, W.; Mögel, L. Z. Chem. 1981, 21, 30-31. Cf. attempts by direct sulfurization of 1,4-diketones: b) Freeman, F.; Kim, D. S. H. L.; Rodriguez, E. J. Org. Chem. 1992, 57, 1722-1727.
- a) Reich, H. J.; Reich, I. L. J. Org. Chem. 1975, 40, 2248-2250. b) Atwell, W. H.; Weyenberg, D. R.; Gilman, H. ibid. 1967, 32, 885-888.
- 18. See, for example, the long-wave maximum  $\lambda_{max} = 509 \text{ nm}$  (lg  $\varepsilon = 3.75$ ) of 3,6-bis(diphenylmethylen)amino-4,5-dicyano-1,2-dithine: a) Moran, J. R.; Huisgen R.; Kalwinsch, I. *Tetrahedron Lett.* **1985**, *26*, 1849-1852.  $-\lambda_{max} = 490 \text{ nm}$  (lg  $\varepsilon = 4.45$ ) of 3-exo,3'-exo-(1R,1'R)-bithiocampher: b) ref.<sup>1n</sup>.
- Thus, 1,8-naphthalene disulfide likewisely shows an absorption in the visible region, discussed as π → σ\* transition (λ<sub>max</sub> = 425, 460, 490 nm): a) Zweig, A.; Hoffmann, A. K. J. Org. Chem. 1965, 30, 3997-4001 Compare: b) Riga, J; Verbist, J. J. J. Chem. Soc. Perkin Trans. II 1983, 1545-1551.
- 20. Cf. Review: Kobayashi K.; Gajurel, C. L. Sulfur Rep. 1986, 7, 123-152.
- 21. a) Cf.: Cook, M. J.; Katritzky, A. R.; Linda, P. Adv. Heterocycl. Chem. 1974, 17, 255-356, and that 331. - b) Freeman, F.; Lu, H.; Ziller, J. W. Acta Cryst. 1996, C52, 1207-1209.
- 22.  $C_{14}H_{22}N_2O_2S_2$ ; deep yellow, orthorhombic crystals; crystal size: 0.30.0.26.0.24 mm; space group: Pbcn(Nr.60); unit cell: a = 5.9219(9), b = 18.479(4), c = 14.320(3) Å; volume: 1567.0(5) Å<sup>3</sup>; Z = 4; d<sub>calcd</sub>. = 1.333 Mg/m<sup>3</sup>; absorption coefficient: 0.343 mm<sup>-1</sup>; wavelength: 0.71069 Å;  $\Theta$  range for data collection: 2.20 to 27.02°; reflections<sub>collectd</sub>: 3440; final R indices [I > 2 $\sigma$  > (I)]: R1 = 0.0369, wR2 = 0.0900; R indices (all data): R1 = 0.0736, wR2 = 0.1039; diffractometer: STOE. — Tables of the atomic coordinates, thermal parameters, bond lengths, and angles have been deposited at the Cambridge Data Centre, University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW. The X-ray data are available on request from the Director of the CCDC by quoting the literature citation of this paper.
- C<sub>28</sub>H<sub>32</sub>S<sub>2</sub>; red, thin plates; crystal size: 0.2·0.25·0.08 mm; space group: PnaS<sub>1(Nr. 33)</sub>; a = 13.382(2), b = 6.201(1), c = 28.261(5) Å; volume: 2325.2 Å<sup>3</sup>; Z = 4; d<sub>calcd.</sub> = 1.22 g/cm<sup>3</sup>; absorption coefficient: 2.35 cm<sup>-1</sup>; MoK<sub>α</sub> radiation; scan-type: Θ-2Θ; 2 theta range: 3-52; reflections<sub>collectd.</sub>: 3158; reflections<sub>observd.</sub>: 1139; final R indices: R = 0.056, R<sub>w</sub> = 0.054; diffractometer: HUBER; program: SHELX76, SHELXS86: Further details of the crystal structure analysis may be obtained from the Fachinformationszentrum Karlsruhe, D-76344 Eggenstein-Leopoldshafen (FRG), on quoting the depository number CSD-400847 and the names of the authors (J. Sieler).
- 24. Cf. further X-ray indications on related (partly anellated) 1,2-dithiines in: a) ref. 1h,i,k. b) Rapp, J. The-

sis, Univ. of München 1988, p. 23. (cf. ref.<sup>18a</sup>). – After completion of this paper, we learned that also microwave spectroscopy reveals a nonplanar structure of the parent compound **6a** with a torsion angle of 53.9°: c) Block, E; Page, J.; Toscano, J., P; Wang, C.-X.; Zhang, X.; DeOrazio, R.; Guo, C.; Sheridan, R. S.; Towers, G. H. N. J. Am. Chem. Soc. 1996, 118, 4719-4720.

 Cf., for example, EtS as substituent: a) Hoffmann, R.; Hartke, K. Chem. Ber. 1980, 113, 919-933. - b) Hartke, K.; Pfleging, E. Liebigs Ann. Chem. 1988, 933-941. - Moreover, attention should be focused to the blue-black 1,2-(thiobenzoyl)benzene Q which does not exist as valence isomeric 2,3-benzodithiine R



due to disadvantegeous benzoanellation (cf. however the existence of 1,2-benzodithiine)<sup>11</sup>. Nevertheless, **Q** tends to extrude sulfur with formation of benzo[c]thiophene **S**: c) Schönberg, A.; Frese, E. *Chem. Ber.* **1968**, *101*, 701- 715 (especially p. 707).

- a) Polarographic measurements were carried out with an apparatus GWP 673 (ZWG of the former Academy of Sciences Berlin). Cf. further: b) Hall, M. E. Anal. Chem. 1953, 25, 556-561. c) Colichman, E. L.; Love, D. L. J. Am. Chem. Soc. 1953, 75, 5736-5737. d) Barber, J.; Smiles, S. J. Chem. Soc. 1928, 1141-1149.
- Maiti, S. N.; Spevak, P.; Singh, M. P. Micetich, R. G.; Reddy, A. V. N. Synth. Commun. 1988, 18, 575-581.
- 28. The process was directly monitored by the decrease of the 1,2-dithiine resonance as well as by the synchronous appearance or increase, respectively, of the thiophene signals under use of a deuterium-lock and exclusion of day-light.
- 29. a) Measurements at 298 K with stirring and irradation with monochromatic light (interference monochromatic filter) and light intensities of 1·10<sup>-10</sup> Einstein/sec in a cross-ray apparatus (cf.: Baumann, H.; Lindenlaub, W; Timpe, H.-J. J. Prakt. Chem. 1978, 320, 825-839, especially 836-837); for 6h: 3.7·10<sup>-4</sup> mol·dm<sup>-3</sup>/solvent. The quantum yields are constant up to conversions of 70%. b) Corrected value in comparison with the value given in ref.<sup>11</sup> (there note 14); error limits generally ± 0.1.
- 30. a) Review: Williams, C. R.; Harpp, D. N. Sulfur Rep. 1990, 10, 103-191. b) According to informations after completion of this paper, episufides (2,6-dithiabicyclo[3.1.0]hex-3-enes) of type L (R<sup>1</sup> = R<sup>2</sup> = H, C<sub>6</sub>H<sub>5</sub>) are, indeed, obtainable by brief irradiation with visible light at -60 to 75°: see ref.<sup>24c</sup>.
- 31. George, M. V.; Mitra, A.; Sukumaran, K. B. Angew. Chem. 1980, 92, 1005-1014; Angew. Chem. Int. Ed. Engl. 1980, 19, 973.
- 32. Cf.: Murata, I. Phosphorus, Sulfur and Silicon 1989, 43, 243-259.
- 33. Cf.: Vogel, E.; Schmidbauer, E.; Altenbach, H.-A. Angew. Chem. 1974, 86, 818-819; Angew. Chem. Int. Ed. Engl. 1974, 13, 736.
- A further possible relationship, the suggested course of the reaction of ethoxycarbonyl azide with thiophenes to N-ethoxycarbonylpyrroles, expelling sulfur, should be considered: Hafner, K.; Kaiser, W. Tetrahedron Lett. 1964, 2185-2190. Lindley, J. M.; Meth-Cohn, O.; Suschitzky, H. J. Chem. Soc. Perkin Trans. I 1978, 1198-1204. Colburn, V. M.; Iddon, B.; Suschitzky, H.; Gallagher, P. T. ibid. 1979, 1337-1340.
- 35. Huisgen, R. Angew. Chem. 1980, 92, 979-1005; Angew. Chem. Int. Ed. Engl. 1980, 19, 947.
- 36. Miller, K. J.; Moschner, K. F.; Potts, K. T. J. Am. Chem. Soc. 1983, 105, 1705-1712.
- Gleiter, R.; Krennrich, G.; Cremer, D.; Yamamoto, K.; Murata, I. J. Am. Chem. Soc. 1985, 107, 6874-6879.
- 38. a) In this connection it should be of interest that we could detect free radicals at irridiation of 6h (polymerization of present acrylonitrile; ESR indication at low temperature photolysis [-196°C]). - b) In view of the inability of 6 to act as sulfur transfer reagent the very recent report of the use of a special thioozonide as efficient sulfur transfer reagent seems most surprisingly: Adam, W.; Weinkötz, S. Chem. Commun. 1976, 177-178.
- 39. Rapid heating, otherwise formation of the corresponding thiophene 12.
- a) Crotti, P.; Feretti, M.; Macchia, F.; Stoppioni, A. J. Org. Chem. 1984, 49, 4706-4711. b) Blum, J.; Becker, Y. J. Chem. Soc. Perkin Trans. 2, 1977, 982-989.
- 41. a) cf. Baddar, F. G.; El-Habashi, A.; Fateen, A. K. J. Chem. Soc. (London) 1965, 3342-3348. b) Campbell, N.; Khanna, N. M. *ibid.* 1949 (S), 33-36.

(Received in Germany 27 June 1996; accepted 9 August 1996)