This article was downloaded by: [Florida Atlantic University] On: 12 November 2014, At: 09:59 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



### Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

# REGIOSELECTIVE SYNTHESIS OF $1,3,5-{}^{13}C_3$ AND $2,4-{}^{13}C_2$ -LABELED 2-DEOXYRIBONOLACTONES

Paul J. Hatala<sup>a</sup>, James Kallmerten<sup>b</sup> & Philip N. Borer<sup>c</sup> <sup>a</sup> Department of Chemistry, Syracuse University, Syracuse, NY, 13244-4100, U.S.A. <sup>b</sup> Department of Chemistry, Syracuse University, Syracuse, NY, 13244-4100, U.S.A. <sup>c</sup> Department of Chemistry, Syracuse University, Syracuse, NY, 13244-4100, U.S.A. Published online: 17 Aug 2006.

To cite this article: Paul J. Hatala , James Kallmerten & Philip N. Borer (2001) REGIOSELECTIVE SYNTHESIS OF 1,3,5-<sup>13</sup>C<sub>3</sub> AND 2,4-<sup>13</sup>C<sub>2</sub>-LABELED 2-DEOXYRIBONOLACTONES, Nucleosides, Nucleotides and Nucleic Acids, 20:12, 1961-1973

To link to this article: http://dx.doi.org/10.1081/NCN-100108326

#### PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

## REGIOSELECTIVE SYNTHESIS OF 1,3,5-<sup>13</sup>C<sub>3</sub> AND 2,4-<sup>13</sup>C<sub>2</sub>-LABELED 2-DEOXYRIBONOLACTONES

Paul J. Hatala, James Kallmerten,\* and Philip N. Borer\*

Department of Chemistry, Syracuse University, Syracuse, NY 13244-4100, USA

#### ABSTRACT

The synthesis of  $1,3,5^{-13}C_3$ - and  $2,4^{-13}C_2$ -labeled 5-O-bromobenzyl-2-deoxyribonolactones **2**, precursors to <sup>13</sup>C-enriched nucleoside phosphoramidites for solid-phase synthesis of DNA oligonucletides, is described. An equimolar combination of these two multiply labeled lactones affords a "population-labeled" mixture of isotopomers which exhibits an approximately 50-fold increase in the sensitivity of <sup>13</sup>C-NMR compared to natural abundance measurements. The <sup>13</sup>C-<sup>13</sup>C 2-bond and 4-bond coupling constants are reported for the lactones; all are < 2Hz, confirming that this labeling scheme should be especially useful for NMR-relaxation measurements.

#### INTRODUCTION

As part of our ongoing effort to investigate the structure and dynamics of DNA oligomers<sup>1</sup>, we recently reported the preparation of the series of regioselectively <sup>13</sup>C<sub>1</sub>-deoxyribonolactones<sup>2</sup>. These labeled deoxyribose derivatives are pivotal intermediates in a novel strategy for <sup>13</sup>C labeling of synthetic oligonucleotides, in which a <sup>13</sup>C-enriched oligonucleotide is prepared from phophoramidite monomers which are themselves synthesized from equimolar mixtures of the <sup>13</sup>C<sub>1</sub>-labeled isotopomers of deoxyribose. Each

Copyright © 2001 by Marcel Dekker, Inc.

<sup>\*</sup>Corresponding authors.

deoxyribose subunit of the oligonucleotide derived from such "population labeled" precursors will contain a single, randomly distributed <sup>13</sup>C atom, resulting in a net <sup>13</sup>C enrichment of approximately 21% at each of the carbon atoms of the phosphodeoxyribose backbone.

The success of our synthetic efforts directed at monolabeled deoxyribose derivatives has prompted consideration of a second generation strategy for the population labeling of an oligonucleotide, in which incorporation of multiple <sup>13</sup>C labels into deoxyribose precursors would enhance both efficiency of our synthetic scheme and the degree to which <sup>13</sup>C label is incorporated into synthetic oligonucleotides. Toward this end, we have examined the preparation of  $1,3,5^{-13}C_3$ - and  $2,4^{-13}C_2$ -labeled deoxyribose systems, reasoning that incorporation of an equimolar mixture of these intermediates into synthetic oligonucleotides would effectively label each of the carbohydrate backbone carbons of the resulting DNA polymer at an overall enrichment of  $\sim$  50-fold over natural abundance. Besides being an appealing alternative for nucleotide structural determination<sup>3</sup>, the  $1,3,5-{}^{13}C_{3}$ - and 2,4-<sup>13</sup>C<sub>2</sub>-labeled deoxyribose systems maximize the contribution of one-bond <sup>13</sup>C-<sup>1</sup>H dipolar effects to NMR relaxation properties. These isotopomers remove one- and three-bond <sup>13</sup>C-<sup>13</sup>C coupling effects that complicate the interpretation of relaxation properties; only natural abundance <sup>13</sup>C-<sup>13</sup>C adjacencies remain. The labeling scheme also provides obvious advantages in sensitivity and reduction of spectral crowding, as well as presenting interesting possibilities for spectral editing and efficient measurement of residual dipolar couplings in oriented systems<sup>4</sup>. Furthermore, recent advances in solid phase oligonucleotide synthesis, indicated that incorporation of precious labeled phosphoramidites into oligonucleotides could be achieved much more efficiently using only 1.75 equivalents<sup>5</sup> of labeled reagent, as opposed to the 10-fold excess normally required<sup>6</sup>. In contrast to the work of Serianni and Bondo<sup>7</sup>, and Agrofoglio et al.<sup>8</sup>, whose methodology can also be used to create <sup>13</sup>C-enriched 2-deoxy-D-ribose, our method does not require the synthesis of D-ribose followed by deoxygenation at C2 to create 2-deoxy-Dribose.

Our previously reported<sup>2</sup> route to the isotopomeric series of  ${}^{13}C_1$ -labeled deoxyribose derivatives, while designed to maximize economy of synthetic effort, is intrinsically limited by the necessity of preparing five unique isotopomers of deoxyribose. An additional liability of our initial approach was the undesired introduction of the minor enantiomer of each isotopomer as the consequence of an early asymmetric transformation that proceeds with high, but not exclusive, stereoselectivity. To address these concerns, we have considered alternative <sup>13</sup>C-labeled precursors to population-labeled oligonucleotides that can be accessed with increased efficiency and an improved control of absolute stereochemistry. Attractive in both regards is the labeling scheme depicted in Fig. 1, in which an equimolar mixture of  $1,3,5-{}^{13}C_3$  and  $2,4-{}^{13}C_2$ -deoxyribonolactones  $1,3,5-{}^{13}C_3-2$  and  $2,4-{}^{13}C_2-2$  would yield the



*Figure 1.* Retrosynthesis of 50% mixture of  $1,3,5^{-13}C_3$ - and  $2,4^{-13}C_2$ -deoxyribonolactones (top center). The labels originate from 2-<sup>13</sup>C-bromoacetic acid (bottom center) and K<sup>13</sup>CN.

population-labeled deoxyribonolactones *pl*-1. Oligonucleotides derived from *pl*-1 would be expected to exhibit an impressive <sup>13</sup>C sensitivity enhancement over unlabeled oligomers and over the previously reported monolabeled systems<sup>2</sup>. Herein we describe the short and stereocontrolled routes to deoxyribonolactones 1,3,5-<sup>13</sup>C<sub>3</sub>-2 and 2,4-<sup>13</sup>C<sub>2</sub>-2, as well as preliminary spectroscopic observations on these compounds and the population-labeled mixtures obtained there from.

#### **RESULTS AND DISCUSSION**

Our integrated synthetic plan for preparing multiple <sup>13</sup>C-labeled deoxyribose derivatives, based on our previously reported conversion of differentiated 2,3-epoxy butanediols **3** to the corresponding lactones via a modified Payne homologation methodology, is shown in Fig. 1. The substrates for Payne homologation, epoxy alcohols  $1,3-^{13}C_2$ -**3** and  $2,4-^{13}C_2$ -**3**, were envisioned deriving from a common starting material, the readily available  $2-^{13}C_2$ bromoacetic acid. For our preparation of  $1,3,5-^{13}C_3$ -**2**, we anticipated introducing the final <sup>13</sup>C label at C1 from <sup>13</sup>C-labeled cyanide ion. Incorporation of the bromobenzyl ether system for protection of the C4 alcohol was expected to facilitate removal of minor enantiomeric impurities by crystallization, providing optically pure final products<sup>9</sup>.

Our synthesis of  $1,3,5^{-13}C_3$ -2 is shown in Scheme 1. O-Alkylation of 2-<sup>13</sup>C-bromoacetic acid with 4-bromobenzyl alcohol followed by treatment of the crude product with diazomethane yielded methyl ester 4. Reduction to aldehyde 5, was achieved with DIBAL at -78 °C, trace amounts (<10%) of the over-reduced alcohol were separated by from 5 by chromatography.



Scheme 1. Reagents: (a) KH, BrBnOH, THF,  $\Delta$ ; (b) CH<sub>2</sub>N<sub>2</sub>, Et<sub>2</sub>O, 0°C; (c) DIBAL, THF, -78°C; (d) NaH, DME, 0°C; (e) DIBAL, THF, 0°C; (f) TBHP, Ti(O*i*Pr)<sub>4</sub>, D-(-)-DIPT, DCM, -20°C; (g) K<sup>13</sup>CN, KI, EtOH-H<sub>2</sub>O,  $\Delta$ , then HCl, H<sub>2</sub>O.

Homologation of **5** by Wadsworth-Emmons-Horner addition with labeled phosphonate **6**<sup>10</sup> furnished the unsaturated ester (not shown), which was directly reduced with DIBAL to allylic alcohol 2,4-<sup>13</sup>C<sub>2</sub>-7. Sharpless asymmetric epoxidation<sup>11</sup> of 2,4-<sup>13</sup>C<sub>2</sub>-7 gave epoxide 2,4-<sup>13</sup>C<sub>2</sub>-3 as a single enantiomer<sup>12</sup> after one recrystallization. Payne rearrangement of epoxide 2,4-<sup>13</sup>C<sub>2</sub>-3 and homologation<sup>13</sup> with K<sup>13</sup>CN, produced a hydroxy nitrile which after hydrolysis and lactonization with aqueous HCl, afforded 1,3,5-<sup>13</sup>C<sub>3</sub>-2.

For our synthesis of 2,4-<sup>13</sup>C<sub>2</sub>-2, we proposed to exploit the latent symmetry of the differentiated butendiol 7, as depicted in Scheme 2. Protection of 2-<sup>13</sup>C-bromoacetic acid as the 4-methoxybenzyl ether and esterification with diazomethane afforded methyl ester 8. Reduction to the corresponding aldehyde 9, addition of phosphonate 6, and reduction with DIBAL yielded the 4-methoxybenzyl protected butenediol derivative 10. At this juncture, the diol system of 10 was inverted by protection of the free



Scheme 2. Reagents: (a) KH, PMBOH, THF,  $\Delta$ ; (b) CH<sub>2</sub>N<sub>2</sub>, Et<sub>2</sub>O, 0°C; (c) DIBAL, THF, -78°C; (d) NaH, DME, **6**, 0°C; (e) DIBAL, THF, 0°C; (f) NaH, THF, BrBnBr, 0°C; (g) DDQ, 9:1 DCM:H<sub>2</sub>O, 0°C; (h) TBHP, Ti(O*i*Pr)<sub>4</sub>, D-(-)-DIPT, DCM, -20°C; (i) KCN, KI, EtOH-H<sub>2</sub>O,  $\Delta$ , then HCl, H<sub>2</sub>O.

#### 2-DEOXYRIBONOLACTONES

alcohol as the 4-bromobenzyl ether, and selective cleavage of the 4-methoxybenzyl residue<sup>14</sup> gave allylic alcohol 1,3-<sup>13</sup>C<sub>2</sub>-7. Conversion to 2,4-<sup>13</sup>C<sub>2</sub>-2 was then accomplished as described for Scheme 1<sup>15</sup>.

To complete our labeling strategy, equimolar amounts of  $1,3,5^{-13}C_3$ -2 and  $2,4^{-13}C_2$ -2 were combined to produces the population labeled mixture 50% *pl*-1. The HSQC spectrum of *pl*-1 was acquired and compared with the HSQC spectrum for  $2,4^{-13}C_2$ -2 (Fig. 2). All <sup>13</sup>C-<sup>1</sup>H correlations appear in the spectrum of *pl*-1 (panel a), but only the correlations for C2-H2,H2' and C4-H4 appear in panel b.

Since we are primarily concerned with the interpretation of relaxation properties, we determined the <sup>2</sup>J coupling constants. Multiple-bond pathways through the ring oxygen may also contribute to these small coupling constant values. The values are shown in Table 1. We also measure a mutiple-bond <sup>13</sup>C-<sup>13</sup>C coupling between C1 and C5. Selected spectra are shown (Fig. 3). The larger two-bond and smaller multiple-bond couplings are clearly present for C1 (Fig. 3a). For C3, both  ${}^{2}J_{C3-C5}$  and  ${}^{2}J_{C3-C1}$  couplings are visible (Fig. 3b, left), but for C5, the  ${}^{2}J_{C5-C3}$  and the  ${}^{3}J/{}^{4}J_{C5-C1}$  are closer in magnitude so only a triplet appears (Fig. 3b, right). Of course C2 is split by  ${}^{2}J_{C2-C4}$  as shown (Fig. 2c); the identical coupling constant value was observed at C4 (not shown).

#### CONCLUSION

We have successfully prepared a new class of population-labeled deoxyribonolactones with ~ 50-fold increase in <sup>13</sup>C sensitivity over natural abundance. The regioselectively labeled lactones are prepared enantiospecifically, and display <sup>13</sup>C-<sup>13</sup>C couplings of no greater than 1.6Hz. The individual deoxyribonolactones are highly stable intermediates and can be stored for extended periods at -20°C, prior to combination and conversion to nucleoside phosphoramidites for use in DNA synthesis. The overall synthetic scheme is sufficiently flexible to allow preparation of all <sup>13</sup>C-isotopomers of DNA sugars in high yield. In our previous paper, the syntheses of five x-<sup>13</sup>C<sub>1</sub> and four 1,x-<sup>13</sup>C<sub>2</sub>-deoxyribonolactone<sup>17</sup> isotopomers is demonstrated. This paper describes the 1,3,5-<sup>13</sup>C<sub>3</sub> and 2,4-<sup>13</sup>C<sub>2</sub>-lactones; simple substitution of unlabeled cyanide affords the 3,5-<sup>13</sup>C<sub>2</sub>-lactone. This accounts for 12 of the 32 isotopomers of 2-deoxy-D-ribose. Further work in progress demonstrates procedures for the other isotopomers as well as attractive population-labeled mixtures.

#### **EXPERIMENTAL SECTION**

General: IR spectra were recorded on a Perkin Elmer Paragon 1000 FT spectrometer, with sodium chloride plates, and are reported in wavenumbers (cm<sup>-1</sup>). <sup>1</sup>H- and <sup>13</sup>C NMR were taken on a Bruker Avance 300 unless



Figure 2. HSQC spectra of 50% pl-1 (panel a), and  $2,4^{-13}C_2$ -2 (panel b).

	$^{2}J$	$^{3}J/^{4}J$
C1-C3	1.6	_
C3-C5	0.9	_
C2-C4	0.8	_
C1-C5	-	0.5

Table 1. <sup>13</sup>C-<sup>13</sup>C Coupling Constants (in Hz)<sup>16</sup>

otherwise noted. Proton chemical shifts are reported in  $\delta$ , using the residual CHCl<sub>3</sub>, as internal reference (7.26 ppm), unless otherwise noted. J values are given in Hz. Carbon chemical shifts are reported in  $\delta$ , using CDCl<sub>3</sub>, as an internal reference (77.0 ppm), unless otherwise noted. <sup>1</sup>H and <sup>13</sup>C NMR spectra were typically acquired in the same number of scans. Optical rotations were measured on either a Jasco DIP-1000 or a Perkin Elmer 241 spectrometer. Enantiomeric excess was determined by either Mosher ester derivatization (ME)<sup>18</sup>, or by chiral shift analysis (CSA) with (+)-Eu(hfc)<sub>3</sub><sup>19</sup>. THF and DME were distilled from sodium benzophenone ketyl under a nitrogen atmosphere. Dichloromethane, benzene and pyridine were distilled from calcium hydride before use. Hexanes, ethyl acetate, ether, and anhydrous ether were used from the supplier without further purification. Silica gel (230–400 mesh) was used for flash column chromatography. TLC analysis was performed on Whatman K6F silica gel 60 plates and were stained with the following: anisaldehyde (method A), phosphomolybdic acid (PMA) (method B), KMnO<sub>4</sub> (method C), 2,4-DNP (method D), and UV light. Reactions were performed in flame or oven dried glassware under a nitrogen or argon atmosphere where appropriate. Other reagents were used from suppliers without purification unless otherwise indicated.

HSQC spectra were acquired on a Bruker Avance 500 spectrometer, using the pulse program of Palmer et al.<sup>20</sup> HSQC spectra were collected with



*Figure 3.* <sup>13</sup>C-<sup>13</sup>C Coupling for (a) C1, (b) C3 (left), and C5 (right), and (c) C2. Each panel is 0.35 ppm.

1 scan per increment, into  $2048 \times 512$  points with no sample spinning and no zero-filling. These gradient selected acquisitions typically required > 15min. The sweep width was 3255Hz in F<sub>2</sub> and 12500Hz in F<sub>1</sub>. A squared sine window function was applied in F<sub>2</sub> and F<sub>1</sub>.

Data for coupling constants was acquired on a Bruker Avance 300 with a direct detection probe. The sweep width was 450 Hz, and no window function was applied. The peaks were fitted using the NMR1 software package (Tripos Assoc.) and coupling constants calculated assuming firstorder coupling.

[2-<sup>13</sup>C]-(4-Bromo-benzyloxy)-acetic acid methyl ester (4). To a suspension of KH (722mg, 18mmol), washed with pentane, in 100mL of THF at 0°C was added 2-<sup>13</sup>C-bromoacetic acid (1g, 0.71 mmol), and then 4-bromobenzyl alcohol (1.46g, 0.86mmol) both in 15mL of THF. The reaction mixture was allowed to warm to room temperature and stirred until the effervescence ceased. The solution was then heated to reflux overnight. After cooling to ambient temperature the reaction mixture was quenched by pouring into a stirring mixture of ice/NH<sub>4</sub>Cl (sat.)/ether 30mL each. The quenched reaction mixture was then transferred to a separatory funnel and the organic layer was washed with NaHCO<sub>3</sub> (sat.). The organic layer was then discarded and the aqueous layer acidified with concd. HCl until acidic. The aqueous layer was then extracted with three 50mL portions of ether, which were dried and concentrated to a white solid. The solid was redissolved in ether, and diazomethane was added to the ether solution at 0°C until a yellow color persisted. Excess diazomethane was quenched with MgSO<sub>4</sub>, and then filtered and concentrated to yield a slightly yellow oil (1.73g, 95% over 2 steps).  $R_{\rm f} = 0.64$ , 1:1 hexanes: EtOAc, method D; IR (neat): 3000.3, 1759.6; <sup>1</sup>H NMR: 3.69 (s, 3H), 4.05 (d, J = 143.8, 2H), 4.51 (d, J = 4.4, 2H), 7.18 (m, 2H), 7.41 (m, 2H); <sup>13</sup>C NMR: 67.10.

[2-<sup>13</sup>C]-(*E*)-(4-Methoxy-benzyloxy)-acetic acid methyl ester (8). The procedure for 4 was followed except 4-methoxybenzyl alcohol was substituted for 4-bromobenzyl alcohol. The procedure yielded a slightly yellow solid (1.33 g, 90% over 2 steps).  $R_{\rm f}$ =0.60, 1:1 hexanes:EtOAc, method D; IR (neat): 3005.4, 1760.3; <sup>1</sup>H NMR: 3.69 (s, 3H), 4.05 (*d*, *J*=143.8, 2H), 4.51 (*d*, *J*=4.4, 2H), 7.18 (m, 2H), 7.41 (m, 2H); <sup>13</sup>C NMR: 67.10.

[2,4-<sup>13</sup>C<sub>2</sub>]-(*E*)-4-(4-Bromo-benzyloxy)-but-2-enoic acid methyl ester. Methyl ester 4 (1.73 g, 0.66 mmol) was dissolved in 25 mL of THF and added to a round bottomed flask. The resulting homogenous solution was cooled to  $-78^{\circ}$ C and DIBAL (1.2eq, 8mL, 1M in toluene) was added dropwise to the solution The reaction was monitored by TLC, and

when the starting material had been consumed the resulting mixture was quenched with MeOH. The quenched solution was allowed to warm to ambient temperature, and was further diluted with ether and Rochelle salt. After stirring for 2h the diluted solution was transferred to a separatory funnel and extracted with two 100mL portions of ether, which were dried over MgSO<sub>4</sub>, and conc. to yield a yellow oil. The aldehyde (5) was flashed in 1:1 hexane:EtOAc. The aldehyde should be used immediately in the next step.  $R_f$ =0.27, 1:1 hexane:EtOAc, method D; <sup>1</sup>H NMR: 4.12 (*d*, *J*=145.4, 2H), 4.58 (*d*, *J*=6.9, 2H), 7.23 (*d*, *J*=8.3, 2H), 7.50 (*d*, *J*=8.5, 2H), 9.72 (*d*, *J*=29.7, 1H); <sup>13</sup>C NMR: 75.4.

To a suspension of NaH (155mg, 0.65mmol) (washed with pentane) in 25mL DME was added [2-<sup>13</sup>C]-(diethoxy-phosphoryl)-acetic acid methyl ester (6) (1.37g, 0.65mmol). The resulting reaction mixture was equilibrated for 30min. After equilibration, the solution was cooled to  $-78^{\circ}$ C and the aldehyde (1.26g, 0.54mmol) was added dropwise in 5mL DME. The reaction mixture was stirred for 30min and then quenched by pouring into a stirring solution of ice/NH<sub>4</sub>Cl (sat)/ether 10mL each. Separation of the layers and extraction of the organic layer, followed by drying over MgSO<sub>4</sub> and conc yielded a clear oil (1.12g, 59% over 2 steps).  $R_f$ =0.57, 1:1 hexanes: EtOAc, method D; IR (neat): 1719.4; <sup>1</sup>H NMR: 3.71 (s, 3H), 4.13 (dm, *J*=141.61, 2H), 4.46 (*d*, *J*=4.12, 2H), 6.09 (dm, *J*=165.21, 1H), 6.94 (m, 1H), 7.18 (*d*, *J*=8.23, 2H), 7.44 (*d*, *J*=8.23, 2H); <sup>13</sup>C NMR: 68.49, 120.79.

[2,4-<sup>13</sup>C<sub>2</sub>]-(*E*)-4-(4-Methoxy-benzyloxy)-but-2-enoic acid methyl ester. The above procedure, applied to **8**, yielded a yellow oil. Aldehyde:  $R_f$ =0.24, 1:1 hexane:EtOAc, method D; IR (neat): 1723; <sup>1</sup>H NMR: 3.81 (s, 3H), 4.06 (dd, *J*=0.5, 141.3, 2H), 4.56 (*d*, *J*=4.39, 2H), 6.89 (*d*, *J*=8.8, 2H), 7.29 (*d*, *J*=8.8, 2H), 9.69 (*d*, *J*=25.8, 1H); <sup>13</sup>C NMR: 75.0. The above procedure, applied to **9**, yielded a yellow oil (975mg, 65% over 2 steps). Ester:  $R_f$ =0.61, 1:1 hexanes:EtOAc, method D; IR (neat): 1720.2; <sup>1</sup>H NMR: 3.74 (s, 3H), 3.81 (s, 3H), 4.14 (dm, *J*=142.43, 2H), 4.49 (*d*, *J*=4.39, 2H), 6.11 (dm, *J*=163.57, 1H), 6.97 (m, 1H), 6.88 (m, 2H), 7.27 (m, 2H); <sup>13</sup>C NMR: 68.25, 120.86.

[2,4-<sup>13</sup>C<sub>2</sub>]-(*E*)-4-(4-Bromo-benzyloxy)-but-2-en-1-ol (2,4-<sup>13</sup>C<sub>2</sub>-7). The unsaturated ester (1.12g, 0.39 mmol) was dissolved in THF and cooled to 0°C in ice. DIBAL (2.5eq, 13.5mL, 1M in toluene) was added to the stirring solution and the resulting mixture was allowed to warm to ambient temperature. After stirring for 2h the mixture was again cooled to 0°C and was quenched with MeOH and then diluted with ether and Rochelle salt and stirred overnight. Extraction of the quenched reaction mixture followed by drying over MgSO<sub>4</sub>, and conc to yield a clear oil (900mg, 89%).  $R_f$ =0.35, 1:1 hexanes:EtOAc, method A; IR (neat): 3388.4, 2853.2; <sup>1</sup>H NMR: 1.77 (s, 1H), 4.02 (dt, J=5.49, 141.34, 2H), 4.16 (m, 2H), 4.46

(d, J = 3.57, 2H), 5.83 (m, 1H), 5.90 (dm, J = 151.49, 1H), 7.21 (d, J = 8.51, 2H), 7.47 (d, J = 8.23, 2H); <sup>13</sup>C NMR: 70.14, 132.38.

[2,4-<sup>13</sup>C<sub>2</sub>]-(*E*)-4-(4-Methoxy-benzyloxy)-but-2-en-1-ol (10). The above procedure, applied to the unsaturated ester from 9, yielded a clear oil (800 mg, 93%).  $R_{\rm f}$ =0.29, 1:1 hexanes:EtOAc, method A; IR (neat): 3388.8, 2838.2; <sup>1</sup>H NMR: 1.63 (bs, 1H), 3.80 (s, 3H), 4.00 (dm, *J*=139.96, 2H), 4.15 (m, 2H), 4.45 (*d*, *J*=4.12, 2H), 5.83 (m, 1H), 5.89 (dm, *J*=152.31, 1H), 6.88 (m, 2H), 7.26 (m, 2H); <sup>13</sup>C NMR: 69.8, 132.1.

[1,3-<sup>13</sup>C<sub>2</sub>]-(*E*)-1-(4-Bromo-benzyloxy)-4-(4-Methoxy-benzyloxy)-but-2ene (11). To a round bottom flask was added NaH (137 mg, 0.57 mmol) and 5mL of THF. The resulting solution was cooled to 0°C in ice. Compound 10 was then added followed by bromobenzyl bromide (1.40g, 0.57 mmol) in 2 portions. The reaction was then allowed to warm to room temperature and monitored by TLC. After the starting material was consumed the reaction was quenched with sat. NH<sub>4</sub>Cl. Typical workup provided an oil (1.25g, 87%).  $R_{\rm f}$ =0.84, 1:1 hexane:EtOAc; IR (neat): 2854.0; <sup>1</sup>H NMR: 3.80 (s, 3H), 4.01 (dt, *J*=4.7, 141.3, 2H), 4.03 (*t*, *J*=4.9, 2H), 4.45 (m, 2H), 5.85 (dm, *J*=152.3, 1H), 5.86 (m, 1H), 6.88 (*d*, *J*=8.5, 2H), 7.24 (m, 4H), 7.46 (*d*, *J*=8.5, 2H); <sup>13</sup>C NMR: 69.8, 129.1.

[1,3-<sup>13</sup>C<sub>2</sub>]-(*E*)-4-(4-Bromo-benzyloxy)-but-2-en-1-ol (1,3-<sup>13</sup>C<sub>2</sub>-7). 11 was dissolved in 9:1 DCM:H<sub>2</sub>O, and cooled to 0°C in ice. DDQ was then added all at once and the reaction was stirred for 1 h. Extraction, drying, and conc yielded an oil (820mg, 96%). IR (neat): 3388.4, 2853.2; <sup>1</sup>H NMR: 1.48 (br s, 1H), 4.03 (t, J=4.9, 2H), 4.16 (dt, J=5.5, 142.2, 2H), 4.47 (s, 2H), 5.84 (dm, J=153.4, 1H), 5.91 (m, 1H), 7.21 (d, J=8.0, 2H), 7.47 (d, J=8.2, 2H); <sup>13</sup>C NMR: 63.0, 128.3.

[2,4-<sup>13</sup>C<sub>2</sub>]-(2R,3R)-[3-(4-Bromo-benzyloxymethyl)-oxiranyl]-methanol (2,4-<sup>13</sup>C<sub>2</sub>-3). To a round bottom flask was added powdered, activated 4Å molecular sieves, and a stir bar. The round bottom flask was then flame dried and allowed to cool to ambient temperature under nitrogen. Methylene chloride, D-(-)-diisopropyl tartrate, and Ti(O*i*Pr)<sub>4</sub> were added sequentially, and the resulting heterogeneous solution was cooled to  $-25^{\circ}$ C (CO<sub>2</sub> (s), CCl<sub>4</sub>) and stirred for 30min. TBHP was then added at  $-25^{\circ}$ C, and the solution was aged for 30min. Meanwhile, the allylic alcohol 2,4-<sup>13</sup>C<sub>2</sub>-7 was diluted in DCM and stirred over 4Å molecular sieves while the catalyst was aging. After the aging period the allylic alcohol was added to the catalyst and the reaction mixture was stirred at  $-25^{\circ}$ C for 1h before being placed in a  $-20^{\circ}$ C freezer overnight. The reaction mixture was quenched by addition of dimethyl sulfide at  $-20^{\circ}$ C and allowing the solution to warm to ambient temperature over 2h. The quenched mixture was filtered through a frit and concentrated to yield a yellow crude product that was purified by flash chromatography to afford as a white solid. Recrystallization was conducted as follows: the solid obtained after flash chromatography was suspended in pentane (1–2mL), and then ether was added until the solution became homogeneous at room temperature. The solution was then placed in a  $-25^{\circ}$ C freezer overnight to allow crystal formation. White needles (826mg, 86%). ee = 100% (ME);  $R_{\rm f}$ =0.39, 2:1 hexane:EtOAc, method A; IR (thin film): 3356.5; <sup>1</sup>H NMR: 1.77 (br s, 1H), 3.09 (d, J=173.7, 1H), 3.24 (m, 1H), 3.71 (m, 2H), 3.78 (dm, J=141.9, 2H), 3.94 (d, J=12.6, 1H), 4.52 (ABq, J=13.7, 2H), 7.21 (d, J=8.2, 2H), 7.47 (d, J=8.5, 2H); <sup>13</sup>C NMR: 55.5, 69.7.

[1,3-<sup>13</sup>C<sub>2</sub>]-(2R,3R)-[3-(4-Bromo-benzyloxymethyl)-oxiranyl]-methanol (1,3-<sup>13</sup>C<sub>2</sub>-3). The above procedure applied to 1,3-<sup>13</sup>C<sub>2</sub>-7 yielded white needles (704 mg, 80%). ee = 100% (ME); IR (thin film): 4305; <sup>1</sup>H NMR: 2.12 (br s, 1H), 3.08 (m, 1H), 3.20 (d, J=158.1, 1H), 3.63 (dd, J=13.2, 142.7, 2H), 3.77 (d, J=10.7, 1H), 4.16 (d, J=12.9, 1H), 4.51 (ABq, J=11.5, 2H), 7.20 (d, J=7.4, 2H), 7.46 (d, J=7.4, 2H); <sup>13</sup>C NMR: 54.1, 61.0.

[1,3,5-<sup>13</sup>C<sub>3</sub>]-(4S,5R)-5-(4-Bromo-benzyloxymethyl)-4-hydroxy-dihydrofuran-2-one (1,3,5-<sup>13</sup>C<sub>3</sub>-2). To a solution of epoxide 2,4-<sup>13</sup>C<sub>2</sub>-3 in 120 mL of 2:1 EtOH:H<sub>2</sub>O, was added K<sup>13</sup>CN and KI. The reaction mixture was heated at reflux for 48 hr, cooled to ambient temperature and carefully acidified to pH 2 with 10% HCl. The resulting mixture was warmed to 60°C and stirred for 16h, at which time the reaction mixture was cooled and transferred to a separatory funnel and the aqueous layer was extracted with three 50mL portions of Et<sub>2</sub>O. The combined ether extracts were dried over MgSO<sub>4</sub>, filtered, and concentrated. The crude product was purified by flash chromatography (2:1 hexanes:EtOAc) to give lactone 1,3,5-<sup>13</sup>C<sub>3</sub>-2 as a slightly yellow solid (593mg, 65%).  $R_{\rm f}$ =0.31, 2:1 hexanes:EtOAc, method A; IR (thin film): 3387, 1780; <sup>1</sup>H NMR: 2.70 (br s, 1H), 2.45 (dm, J=2.7, 18.0, 1H), 2.91 (dt, J=5.8, 18.3, 1H), 3.66 (dm, J=144.0, 2H), 4.46 (m, 3H), 4.51 (dm, J=153.5, 1H), 7.14 (d, J=8.2, 2H), 7.47 (d, J=8.5, 2H); <sup>13</sup>C NMR: 69.5, 69.6, 175.9.

[2,4-<sup>13</sup>C<sub>2</sub>]-(4S,5R)-5-(4-Bromo-benzyloxymethyl)-4-hydroxy-dihydrofuran-2-one (2,4-<sup>13</sup>C<sub>2</sub>-2). The above procedure applied to 1,3-<sup>13</sup>C<sub>2</sub>-7 yielded a yellow solid (479 mg, 60%). IR (thin film): 3365, 1772; <sup>1</sup>H NMR: 2.71 (br s, 1H), 2.45 (ddd, J = 1.8, 18.3, 136.8, 1H), 2.92 (ddd, J = 6.7, 18.3, 133.7, 1H), 3.67 (m, 2H), 4.46 (ABq, J = 11.0, 2H), 4.48 (d, J = 150.5, 1H), 4.53 (m, 1H), 7.14 (d, J = 8.2, 2H), 7.47 (d, J = 8.2, 2H); <sup>13</sup>C NMR: 38.4, 86.2.

#### REFERENCES

- (a) LaPlante, S.R.; Borer, P.N. Changes in <sup>13</sup>C NMR Chemical Shift of DNA as a Tool for Monitoring Drug Interaction. Biophys. Chem. in press. (b) Borer, P.N.; LaPlante, S.R.; Hakkinen, A.; Kumar, A.; Levy, G.C.; Martin, A.; Zanatta, N. <sup>13</sup>C NMR Relaxation in Three DNA Oligonucleotide Duplexes: Model-Free Analysis of Internal and Overall Motion. Biochemistry 1994, 33, 2441-2450. (c) LaPlante, S.R.; Hakkinen, A.; Levy, G.C.; van der Marel, J.; Wang, A.H.-J.; Zanatta, N.; Borer, P.N.<sup>13</sup>C NMR of the Deoxyribose Sugars in Four DNA Oligonucleotide Duplexes: Assignment and Structural Features. Biochemistry 1994, 33, 2430–2440. (d) Ashcroft, J.; LaPlante, S.R.; Borer, P.N.; Cowburn, D. Sequence specific carbon-13 NMR assignment of non-protonated carbons in [d(TAGCGC-TA)]<sub>2</sub> using proton detection. J. Am. Chem. Soc. 1989, 111, 363-365. (e) La-Plante, S.R.; Ashcroft, J.; Cowburn, D.; Levy, G.C.; Borer, P.N.; Carbon-13 NMR assignments of the protonated carbons of [d(TAGCGCTA)]<sub>2</sub> by twodimensional proton-detected heteronuclear correlation. J. Biomol. Struct. Dyn. 1988, 5, 1089–1099. (f) Borer, P.N.; LaPlante, S.R.; Zanatta, N.; Levy, G.C. Hydrogen-bonding effects and carbon-13 NMR of the DNA double helix. Nucleic Acids Res. 1988, 16, 2323-2332. (g) Stone, M.P.; Winkle, S.A.; Borer, P.N. Carbon-13 NMR of ribosyl ApApA, ApApG and ApUpG. J. Biomol. Struct. Dyn. 1986, 3, 767-781. (h) Borer, P.N.; Zanatta, N.; Holak, T.A.; Levy, G.C.; van Boom, J.H.; Wang, A.H.-J. Conformation and Dynamics of short DNA duplexes:  $(dC-dG)_3$  and  $(dC-dG)_4$ . J. Biomol. Struct. Dyn. 1984, 1, 1373–1386.
- Hayes, M.P.; Hatala, P.J.; Sherer, B.A.; Tong, X.; Zanatta, N.; Borer, P.N.; Kallmerten, J. Regioselective synthesis of the <sup>13</sup>C<sub>1</sub>-labeled 2-deoxyribonolactones. Tetrahedron 2001, *57*, 1515–1524.
- Kaptein, R.; McConnell, B.; Serianni, A.S.; Silks, L.A. In Stable Isotope Applications in Biomolecular Structure and Mechanisms, Proceedings of the Conference, Santa Fe, New Mexico, March 27-30, 1994; Trewhella, J., Cross, T.A., Unkefer, C.J., Eds.; Los Alamos National Laboratory: Los Alamos, N.M., 1994; Conference LA-12893-C, p 333.
- Tjandra, N.; Bax, A. Direct measurement of distances and angles in biomolecules by NMR in a dilute liquid crystalline medium. Science 1997, 278, 1111–1114. Hansen, M.R.; Hanson, P.; Pardi, A. Filamentous bacteriophage for aligning RNA, DNA, and proteins for measurement of nuclear magnetic resonance dipolar coupling interactions. Methods Enzymol. 2000, 317, 220– 240.
- Eleuteri, A.; Capaldi, D.C.; Krotz, A.H.; Cole, D.L.; Ravikumar, V.R. Pyridinium Trifluoroacetate/*N*-Methylimidazole as an Efficient Activator for Oligonucleotide Synthesis via the Phosphoramidite Method. Org. Process Res. Dev. 2000, 4, 182–189.
- Caruthers, M.H.; Barone, A.D.; Beaucage, S.L.; Dodds, D.R.; Fisher, E.F.; McBride, L.J.; Matteucci, M.; Stabinsky, Z.; Tang, J.Y. Chemical Synthesis of Deoxyoligonucleotides by the Phosphoramidite Method. Meth. Enzymol. 1987, 154, 287–313.

#### 2-DEOXYRIBONOLACTONES

- Serianni, A.S.; Bondo, P.B. <sup>13</sup>C-Labeled D-Ribose: Chemi-Enzymatic Synthesis of Various Isotopomers. Journal of Biomolecular Structure & Dynamics 1994, 11, 1133–1148.
- Agrofoglio, L.A.; Jacquinet, J.-C.; Lancelot, G. A Multigram, Stereoselective Synthesis of D-[<sup>13</sup>C<sub>5</sub>]Ribose from D-[<sup>13</sup>C<sub>6</sub>]Glucose and Its Conversion into [<sup>13</sup>C<sub>5</sub>]Nucleosides. Tetrahedron Lett. **1997**, *38*, 1411–1412.
- Enantioselective crystallization of the perdeuterated version of 3, has been reported: see Goering, B.K.; Lee, K.; An, B.; Cha, J.K. A Practical Synthesis of Perdeuterated Deoxyriboses and Deoxyribonucleosides. J. Org. Chem. 1993, 58, 1100–1103.
- Phosphonate 6 can be synthesized in 2 steps from 2-bromoacetic acid: see Samuel, S.P.; Niu, T.-Q.; Erickson, K.L. Carbanionic rearrangements of halomethylenecycloalkanes. J. Am. Chem. Soc. 1989, 111, 1429–1436.
- (a) Katsuki, T.; Lee, A.W.M.; Ma, P.; Martin, V.S.; Masamune, S.; Sharpless, K.B.; Tuddenham, D.; Walker, F.J. Synthesis of Saccharides and Related Polyhydroxylated Natural Products. 1. Simple Alditols. J. Org. Chem. **1982**, *47*, 1373–1378. (b) Ma, P.; Martin, V.S.; Masamune, S.; Sharpless, K.B.; Viti, S.M. Synthesis of Saccharides and Related Polyhydroxylated Natural Products. 2. Simple Deoxyalditols. J. Org. Chem. **1982**, *47*, 1378–1380.
- 12. Enantiomeric excess was determined by Mosher ester analysis of the epoxy alcohols. See experimental for details.
- 13. Wrobel, J.E.; Ganem, B. Total Synthesis of (-)-Vertinolide. A General Approach to Chiral Tetronic Acids and Butenolides from Allylic Alcohols. J. Org. Chem. **1983**, *48*, 3761–3764.
- 14. Oikawa, Y.; Yoshioka, T.; Yonemitsu, O. Specific removal of O-methoxybenzyl protection by DDQ oxidation. Tetrahedron Lett. **1982**, *23*, 885–888.
- 15. KCN was substituted for K<sup>13</sup>CN in the homologation/lactonizaiton step.
- 16. Coupling constants have been determined to  $\pm 0.1$  Hz.
- 17. Hatala, P.J., Kallmerten, J.; Borer, P.N. Syracuse University, Syracuse, NY. Unpublished work, 1999.
- (a) Dale, J.A.; Mosher, H.S. Nuclear Magnetic Resonance Enantiomer Reagents. Configurational Correlations via Nuclear Magnetice Resonance Chemical Shifts of Diastereomeric Mandelate, O-Methylmandelate, and α-Methoxy-α-trifluoromethylphenylacetate (MTPA) esters. J. Am. Chem. Soc. 1973, 95, 512–519. (b) Sullivan, G.R.; Dale, J.A.; Mosher, H.S. Correlation of Configuration and <sup>19</sup>F Chemical Shifts of α-Methoxy-α-trifluoromethylphenylacetate derivatives. J. Org. Chem. 1973, 38, 2143–2147.
- 19. Fraser, R.R.; Petit, M.A.; Saunders, J.K. Determination of enantiomeric purity by an optically active nuclear magnetic resonance shift reagent of wide applicability. J. Chem. Soc., Chem. Comm. **1971**, 1450–1451.
- Palmer, A.G. III; Cavanagh, J.; Wright, P.E.; Rance, M. Sensitivity improvement in proton-dectected two-dimensional heteronuclear correlation NMR spectroscopy. J. Magn. Reson. 1991, 93, 151–170.

Received April 10, 2001 Accepted September 26, 2001