

Discovery of a New and Efficient Small Molecule for Neuronal Differentiation from Mesenchymal Stem Cell

Na Ri Kim,[†] Seung Kyu Kang,[†] Hyun Hee Ahn,[†]
Sung Wook Kwon,[†] Woul Seong Park,[†] Ki Suk Kim,[‡]
Sung Soo Kim,[†] Hee Jung Jung,[†] Sang Un Choi,[†]
Jin Hee Ahn,^{*,†} and Kwang Rok Kim^{*,†}

[†]*Drug Discovery Division, Korea Research Institute of Chemical Technology, Daejeon, 305-600, Korea and* [‡]*Korea Institute of Toxicology, Daejeon, 305-600, Korea*

Received October 20, 2009


Abstract: A new synthetic small molecule, compound **1**, which induced a neuronal differentiation from mesenchymal stem cells (MSCs) with an excellent efficiency, was identified. Furthermore the differentiated cell by **1** showed the neural electrophysiological and cholinergic neuron properties.

Neuronal differentiation of stem cells has received much attention because of its potential for the treatment of neurodegenerative diseases such as Parkinson's and Alzheimer's diseases.¹ Moreover, small molecule inducers that can direct differentiation of stem cells are useful tools to facilitate therapeutic applications.² Recently, several small molecules have been reported to have neuronal differentiation ability,^{3–9} but there is no report of single small molecule that can induce functional activity such as electrophysiological properties, so far.

To identify effective neuronal inducers, we screened more than 20 000 small molecules using the neuronal precursor-like PC12¹⁰ cell lines as a primary screening and identified several novel small molecules. Among them, **1**, a quinoxaline based small molecule, revealed the most effective activity to show neurite outgrowth and increase neuronal markers (β -III tubulin and NSE^a) in the PC12 cells (Supporting Information Figure S1). Thus, we considered **1** as a neuronal inducer prototype. Compound **1** and its derivatives were synthesized according to Scheme 1.

The 1-bromo-3-phenylpropane **2** was converted to Grignard reagent in the presence of I₂, followed by coupling with dimethyl oxalate to produce **3**. Keto ester **3** was cyclized with methyl 3,4-diaminobenzoate **4** in EtOH to give a mixture of **5** and **6** (~7:3 ratio), which were separated by crystallization. After ester hydrolysis under basic condition, NH₂OTHP was introduced to yield **7** and **8**, and finally

Scheme 1. Synthesis of Compound **1** and Its Derivatives^a



^a Reagents and conditions: (a) Mg, I₂, THF, room temperature, 0.5 h, and then dimethyl oxalate, Et₂O, –78 to 0 °C 3 h; (b) EtOH, room temperature, 12 h; (c) LiOH, THF/MeOH/H₂O, 50 °C, 10 h; (d) EDCI, HOBT, NH₂OTHP, DMF, room temp, 3 h; (e) trifluoroacetic acid, MeOH, 35 °C, 12 h.

deprotection using trifluoroacetic acid produced **1** and **9**, respectively.

The possibility of neuronal differentiation in rat mesenchymal stem cells (MSCs) (Figure S2, MSCs characterization) with **1** and its derivatives (**5**, **6**, **7**, **8**, and **9**) was explored. The MSCs are highly important and easily accessible adult stem cells that could be differentiated into many cell types including osteoblasts, adipocytes, fibroblasts, myocytes, and neurons. The ability of MSCs to differentiate into neurons holds considerable promise because of their potential to serve as a possible unlimited source of stem cells to treat neurodegenerative diseases. Interestingly, **1** showed a significant phenotype change of MSCs, whereas the others including **9** did not induce any morphological change. After treatment of **1** (20 μ M) for 2 days, **1** induced morphological change and increased the β -III tubulin expression without cytotoxicity (Figure 1a,b and Figure S3).

We also tested the neuron inducing properties of nerve growth factor (NGF),¹¹ representative neuronal differentiation inducers, and the neurophathiazole,⁶ a small molecule that induces neuronal differentiation of neural progenitor cells; however, these agents did not induce any change in the MSCs system (Figures 1b and S4). Compound **1** dose- and time-dependently increased neuronal markers (β -III tubulin and NSE) (Figure 1c). On the basis of Nissl staining,¹² which is a simple and reliable method for neuron detection, it is estimated that >95% of MSCs were converted to a neuronal

^{*}To whom correspondence should be addressed. For J.H.A.: phone, +82-42-860-7076; fax, +82-42-860-7160; e-mail, jhahn@krikt.re.kr. For K.R.K.: phone, +82-42-860-7245; fax, +82-42-860-7160; e-mail, kkrok@krikt.re.kr.

^a Abbreviations: MSCs, mesenchymal stem cells; NGF, nerve growth factor; THF, tetrahydrofuran; EDCI, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide; NH₂OTHP, tetrahydropyran-2-ylamine; HOBT, hydroxybenzotriazole; DMF, dimethylformamide; NSE, neuron-specific enolase; GFAP, glial fibrillary acidic protein; DAPI, 4',6-diamidino-2-phenylindole; Chrna2, cholinergic receptor, nicotinic, α 2; Chrn2, cholinergic receptor, nicotinic, β 2; Chrm4, cholinergic receptor, muscarinic 4; Fbxo2, F-box protein 2; Szn1, Smad-interacting zinc finger protein; RFU, relative fluorescence unit.




Figure 3. Quantitative PCR of some representative genes in **1** treated MSCs. The MSCs were incubated with **1** at 10 and 20 μ M and Ctrl (DMSO) for 2 days, and the mRNA levels were determined by quantitative PCR analysis. The experiments were performed three times independently.




Figure 4. Membrane potential assay of MSCs with acetylcholine. MSCs were incubated with 20 μ M **1** for 48 h. Membrane potential were detected by FLIPR assay kit. Data were analyzed using Flex station software. Open markers are control. Closed markers are **1** treatment. Arrow indicated treatment of 10 μ M acetylcholine.

In the FLIPR assay, the membrane potential was changed on **1** treated MSCs (Figure 4). This result suggests that **1** can differentiate MSCs into a specific type of cholinergic neuronal cell.

In conclusion, we have developed a new synthetic small molecule (**1**) that can induce neuronal differentiation in MSCs. Compound **1** has a capacity to commit MSCs to neuronal fate, which was confirmed by morphological change and expression of neuronal markers (β -III tubulin and NSE). When treated with **1**, more than 95% of MSCs were converted to neuron-like cells. Moreover, quantitative PCR results showed that the expressions of neuropeptide, cholinergic receptors, and Fbxo2, which are a neuronal cell marker, were up-regulated by **1**. Importantly, **1** treated MSCs showed the neural electrophysiological and cholinergic neuron properties. Taken together, **1** will be a useful tool in studying cell therapy and the molecular mechanisms that determine cell fate, even if further biological evaluations and chemical derivatization to find better small molecules are now in progress.

Acknowledgment. This work was supported by the Creative Research Fund 2008, Korea Research Institute of Chemical Technology.

Supporting Information Available: Experimental procedures, biological assays, and figures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Kim, J. H.; Auerbach, J. M.; Rodriguez-Gomez, J. A.; Velasco, I.; Gavin, D.; Lumelsky, N.; Lee, S.; Nguyen, J.; Sanchez-Pernaute, R.; Bankiewicz, K.; McKay, R. Dopamine neurons derived from embryonic stem cells function in an animal model of Parkinson's disease. *Nature* **2002**, *418*, 50–56.
- (2) Ding, S.; Schultz, P. G. A role for chemistry in stem cell biology. *Nat. Biotechnol.* **2004**, *22*, 833–840.
- (3) Wichterle, H.; Lieberam, I.; Porter, J. A.; Jessell, T. M. Directed differentiation of embryonic stem cell into motor neurons. *Cell* **2002**, *110*, 385–397.
- (4) Ding, S.; Wu, T. Y. H.; Brinker, A.; Peters, E. C.; Hur, W.; Gray, N. S.; Schultz, P. G. Synthetic small molecules that control stem cell fate. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 7632–7637.
- (5) Hwang, K. C.; Kim, J. Y.; Chang, W.; Kim, D. S.; Lim, S.; Kang, S. M.; Song, B. W.; Ha, H. Y.; Huh, Y. J.; Choi, I. G.; Hwang, D. Y.; Song, H.; Jang, Y.; Chung, N.; Kim, S. H.; Kim, D. W. Chemicals that modulate stem cell differentiation. *Proc. Natl. Acad. Sci. U.S.A.* **2008**, *105*, 7467–7471.
- (6) Warashina, M.; Min, K. H.; Kuwabara, T.; Huynh, A.; Gage, F. H.; Schultz, P. G.; Ding, S. A synthetic small molecule that induces neuronal differentiation of adult hippocampal neural progenitor cells. *Angew. Chem., Int. Ed.* **2006**, *45*, 591–593.
- (7) Schneider, J. W.; Gao, Z.; Li, S.; Farooqui, M.; Tang, T.; Bezprozvanny, I.; Frantz, D. E.; Hsieh, J. Small-molecule activation of neuronal cell fate. *Nat. Chem. Biol.* **2008**, *4*, 408–410.
- (8) Balasubramanian, V.; Boddeke, E.; Bakels, R.; Kust, B.; Kooistra, S.; Veneman, A.; Copray, S. Effects of histone deacetylation inhibition on neuronal differentiation of embryonic mouse neural stem cells. *Neuroscience* **2006**, *143*, 939–951.
- (9) Williams, D. R.; Lee, M. R.; Song, Y. A.; Ko, S. K.; Kim, G. H.; Shin, I. Synthetic small molecules that induce neurogenesis in skeletal muscle. *J. Am. Chem. Soc.* **2007**, *129*, 9258–9259.
- (10) Hall, F. L.; Fernyhough, P.; Ishii, D. N.; Vulliamt, P. R. Suppression of nerve growth factor-directed neurite outgrowth in PC12 cells by sphingosine, an inhibitor of protein kinase C. *J. Biol. Chem.* **1988**, *263*, 4460–4466.
- (11) Scheibe, R. J.; Ginty, D. D.; Wagner, J. A. Retinoic acid stimulates the differentiation of PC12 cells that are deficient in cAMP-dependent protein kinase. *J. Cell Biol.* **1991**, *113*, 1173–1182.
- (12) Korzhnevskii, D. E.; Otellin, V. A. Immunocytochemical detection of astrocytes in brain slices in combination with Nissl staining. *Neurosci. Behav. Physiol.* **2005**, *35*, 639–641.
- (13) Yang, H.; Xia, Y.; Lu, S. Q.; Soong, T. W.; Feng, Z. W. Basic fibroblast growth factor-induced neuronal differentiation of mouse bone marrow stromal cells requires FGFR-1, MAPK/ERK, and transcription factor AP-1. *J. Biol. Chem.* **2008**, *283*, 5287–5295.
- (14) Woodbury, D.; Schwarz, E. J.; Prockop, D. J.; Black, I. B. Adult rat and human bone marrow stromal cells differentiate into neurons. *J. Neurosci. Res.* **2000**, *61*, 364–370.
- (15) Sanchez-Ramos, J. R. Neural cells derived from adult bone marrow and umbilical cord blood. *J. Neurosci. Res.* **2002**, *69*, 880–893.
- (16) Dezawa, M.; Kanno, H.; Hoshino, M.; Cho, H.; Matsumoto, N.; Itokazu, Y.; Tajima, N.; Yamada, H.; Sawada, H.; Ishikawa, H.; Mimura, T.; Kitada, M.; Suzuki, Y.; Ide, C. Specific induction of neuronal cells from bone marrow stromal cells and application for autologous transplantation. *J. Clin. Invest.* **2004**, *113*, 1701–1710.
- (17) Barnabe, G. F.; Schwandt, T. T.; Calcagnotto, M. E.; Motta, F. L.; Martinez, G.; Oliveira, A. D.; Keim, L. M. N.; D'Almeida, V.; Mendez-Otero, R.; Mello, L. E. Chemically-induced rat mesenchymal stem cells adopt molecular properties of neuronal-like cells but do not have basic neuronal functional properties. *PLoS One* **2009**, *4*, e5222.
- (18) Cho, M. H.; Lee, J. H.; Ahn, H. H.; Lee, J. Y.; Kim, E. S.; Kang, Y. M.; Min, B. H.; Kim, J. H.; Lee, H. B.; Kim, M. S. Induction of neurogenesis in rat bone marrow mesenchymal stem cells using purine structure-based compounds. *Mol. Biosyst.* **2009**, *5*, 609–611.
- (19) Erhardt, J. A.; Hynicka, W.; DiBenedetto, A.; Shen, N.; Stone, N.; Paulson, H.; Pittman, R. N. A novel F Box protein, NFB42, is highly enriched in neurons and induces growth arrest. *J. Biol. Chem.* **1998**, *273*, 35222–35227.
- (20) Cho, G.; Lim, Y.; Zand, D.; Golden, J. A. Sizz1 is a novel protein that functions as a transcriptional coactivator of bone morphogenic protein signalling. *Mol. Cell. Biol.* **2008**, *28*, 1565–1572.