Copper(II) Carboxylate-Promoted Intramolecular Carboamination of Alkenes for the Synthesis of Polycyclic Lactams

LETTERS 2007 Vol. 9, No. 26 5477-5480

ORGANIC

Peter H. Fuller and Sherry R. Chemler*

Department of Chemistry, Natural Science Complex, University at Buffalo, The State University of New York, Buffalo, New York 14260

schemler@buffalo.edu

Received October 2, 2007



ABSTRAC1



The copper(II) carboxylate-promoted intramolecular carboamination reactions of variously substituted γ -alkenyl amides have been investigated. These oxidative cyclization reactions efficiently provide polycyclic lactams, useful intermediates in nitrogen heterocycle synthesis, in good to excellent yields. The efficiency of the carboamination process is dependent upon the structure of the amide backbone as well as the nitrogen substituent.

Nitrogen heterocycles are useful intermediates in complex molecule synthesis, as well as an abundant class of biologically active molecules. The rapid assembly of such molecules and the search for new molecular scaffolds provides a constant challenge to the synthetic and medicinal chemist; thus, new methods of entry into these systems are especially useful.¹ One particularly attractive method is the intramolecular carboamination reaction of alkenes.²

10.1021/ol702401w CCC: \$37.00 © 2007 American Chemical Society Published on Web 11/29/2007

In 2004, our lab disclosed the first copper(II) carboxylatepromoted intramolecular carboamination reaction of Narylsulfonyl 2-allyl anilines (eq 1).^{2c}



More recently we reported further substrate versatility and diastereoselectivity, and provided an experimental analysis of the reaction mechanism.^{2f}

In both of these accounts, a variety of γ - and δ -alkenyl *N*-aryl sulfonamides were shown to undergo the oxidative cyclization in the presence of copper(II) carboxylates to produce the corresponding polycyclic sultams in modest to good yield.^{2c,2f} In an effort to expand the utility of the method,

⁽¹⁾ Zeni, G.; Larock, R. C. Chem. Rev. 2004, 104, 2285.

^{(2) (}a) Larock, R. C.; Yang, H.; Weinreb, S. M.; Herr, R. J. J. Org. Chem. 1994, 59, 4172. (b) Harayama, H.; Abe, A.; Sakado, T.; Kimura, M.; Fugami, K.; Tanaka, S.; Tamaru, Y. J. Org. Chem. 1997, 62, 2113. (c) Sherman, E. S.; Chemler, S. R.; Tan, T. B.; Gerlitis, O. Org. Lett. 2004, 6, 1573. (d) Yip, K.-T.; Yang, M.; Law, K.-L.; Zhu, N.-Y.; Yang, D. J. Am. Chem. Soc. 2006, 8, 3130. (e) Scarborough, C. C.; Stahl, S. S. Org. Lett. 2006, 8, 3251. (f) Sherman, E. S.; Fuller, P. H.; Kasi, D.; Chemler, S. R. J. Org. Chem. 2007, 72, 3896. (g) Peng, J.; Lin, W.; Yuan, S.; Chen, Y. J. Org. Chem. 2007, 72, 3145. (h) Nakhla, J. S.; Wolfe, J. P. Org. Lett. 2007, 9, 3279. (i) Bertrand, M. B.; Leathen, M. L.; Wolfe, J. P. Org. Lett. 2007, 9, 457. (j) Wolfe, J. P. Eur. J. Org. Chem. 2007, 571 and references therein.

a variety of aryl-, vinyl-, and aliphatic γ -alkenyl amides were investigated in the copper(II)-promoted carboamination reaction. The results of our findings are described in this report.

Under our original reaction conditions (e.g., eq 1), 2-allyl aniline amides were poor substrates for the copper(II) carboxylate-promoted intramolecular carboamination reaction, significantly less reactive in comparison to their aryl sulfonamide counterparts (Table 1, entries 1 and 2).^{2f} Success

Table 1.	1. Reaction Optimization ^{<i>a</i>} $\downarrow \downarrow $		
entry	copper salt	$temperature (^{\circ}C)$	yield ^b (%)
1°	$Cu(OAc)_2$	170	16
2^{c}	$Cu(OAc)_2$	190	39
3	$Cu(ND)_2$	190	59
4	$Cu(EH)_2$	190	61

^{*a*} Conditions: Substrate in DMF (0.1 M) was treated with Cu(OR)₂ (3 equiv) and Cs₂CO₃ (1 equiv). The mixture was heated to the indicated temperature for 24 h in a pressure tube. ^{*b*} Yields refer to product isolated by chromatography on SiO₂. ^{*c*} DMSO (4 equiv) was used. The remainder of the material was either starting olefin or olefin-isomerized starting material. Cu(EH)₂ = copper(II) 2-ethylhexanoate, Cu(ND)₂ = copper(II) neodecanoate.

in this matter was subsequently realized when *N*-benzoyl-2-allyl aniline was treated with more organic soluble copper-(II) salts and slightly higher reaction temperatures, providing polycyclic lactam **2a** in moderate yield (Table 1). The organic soluble copper salts, Cu(II) neodecanoate [Cu(ND)₂] and Cu-(II) 2-ethylhexanoate [Cu(EH)₂], were shown to be more reactive than Cu(OAc)₂ (Table 1, entries 3 and 4).^{2f,3} Cu-(EH)₂ was subsequently used throughout the substrate screening because of its lower cost and ease of use [Cu-(EH)₂ is purchased as a solid, whereas Cu(ND)₂ is purchased as a solution in toluene].

A variety of 2-allyl aniline-derived aryl amides were oxidatively cyclized in an efficient manner using the optimized reaction conditions (Table 2). The mildly electron-deficient halogenated substrates **1b** and **1c** reacted efficiently (Table 2, entries 2–4 and 7). Worth noting, the flourine was displaced with dimethylamine when the reaction was run in DMF (Table 2, entry 3) (dimethylamine presumably arises from thermally decomposed DMF). When *tert*-butyl benzene was used as solvent, the carboamination adduct was obtained in good yield with the fluorine intact (Table 2, entry 4). 4-Cyano and 4-methoxy arylamides displayed comparatively lower reactivity (Table 2, entries 5 and 6). Meta-substituted aryl amides demonstrated a preference (ca. 1.8:1) for the ortho addition product over the para (Table 2, entries 7 and 8). This ortho preference is consistent with that of meta-



^{*a*} Conditions: Substrate in DMF (0.1 M) was treated with Cu(EH)₂ (3 equiv) and Cs₂CO₃ (1 equiv). The mixture was heated to 190 °C for 24 h in a pressure tube. ^{*b*} Yields refer to the sum of products isolated by chromatography on SiO₂. The remainder of the material was either starting olefin or olefin-isomerized starting material. ^{*c*} *tert*-Butyl benzene was used as solvent. The structures of the products (e.g., regioisomer) were assigned by analysis of the aromatic region of the ¹H NMR spectra.

substituted aryl sulfonamides^{2c,f} and provides evidence for C–C bond formation via the addition of a carbon radical to an aromatic ring.^{2f,4}

2-Allyl aniline-derived vinyl amides are reactive carboamination substrates as well (Table 3). Interestingly, the unsubstituted vinyl amides **9a**–**c** cyclized in 6-*endo* fashion at 140 °C to provide the polycyclic α,β -unsaturated lactones **10a**–**c** in good yield (Table 3, entries 1–3). These observations are in contrast to similar palladium-catalyzed processes where the 5-*exo* cyclization product (similar to **11**) was the only observed regioisomer.^{2d,5} Increasing the reaction temperature to 190 °C, however, resulted in a 1.2:1 mixture of the 6-*endo* to 5-*exo* carboamination adducts (Table 3, entry 4). The 5-*exo* product **11a** terminated in hydrogen-atom capture rather than olefin formation. This observation presents an intriguing example of temperature control of regiochemistry.

Disappointingly, the more substituted vinyl amides 12 and 14 were less reactive (Table 3, entries 5 and 6). These substrates required higher reaction temperature (190 °C) in order to consume most of the substrate, but in doing so, resulted in the concomitant formation of a mixture of 5-*exo* cyclization products and isomerized olefin starting material.

^{(3) (}a) Antilla, J. C.; Buchwald, S. L. *Org. Lett.* **2001**, *3*, 2077. (b) Baran, P. S.; Richter, J. M. *J. Am. Chem. Soc.* **2004**, *126*, 7450. (c) For a review of copper-facilitated C–N and C–C bond formation, see Chemler, S. R.; Fuller, P. H. *Chem. Soc. Rev.* **2007**, *36*, 1153 and references therein.

^{(4) (}a) Ito, R.; Migita, T.; Morikawa, N.; Simamura, O. *Tetrahedron* **1965**, *21*, 955. (b) Pryor, W. A.; Davis, W. H.; Gleaton, J. H. *J. Org. Chem.* **1975**, *40*, 2099.

^{(5) (}a) Hegedus, L. S.; McKearin, J. M. J. Am. Chem. Soc. **1982**, 104, 2444. (b) Danishefsky, S.; Taniyama, E. Tetrahedron Lett. **1983**, 24, 15.





^{*a*} Conditions: Substrate in DMF (0.1 M) was treated with Cu(EH)₂ (3 equiv) and Cs₂CO₃ (1 equiv). The mixture was heated to 140 °C for 24 h in a pressure tube. ^{*b*} Yields refer to the sum of products isolated by chromatography on SiO₂. The remainder of the material was either starting olefin or olefin-isomerized starting material. ^{*c*} Heated to 190 °C for 24 h.

Our previous studies showed that the copper(II) carboxylate-promoted intramolecular carboamination reaction is effective in cyclizing a variety of aliphatic *N*-aryl sulfonamides (e.g., eq 2).^{2f}



This prompted us to investigate the cyclization reaction of a corresponding *N*-benzoyl aliphatic amide (eq 3). Unfortunately, this substrate was shown to be unreactive upon heating at 190 $^{\circ}$ C for 24 h.



This result led us to examine a variety of structurally different imides and amides as shown Table 4. Imide **17** cyclized in high yield while the amide substrate in eq 3 did not react. This indicates that an sp² hybridized carbon in the substrate's backbone decreases the activation energy for cyclization. (This could be due to lower relative torsional strain in the five-membered ring of the transition state and product for substrate **17**.) In addition, imide **17** cyclized at lower temperature and in higher yield and selectivity than



^{*a*} Conditions: Substrate in DMF (0.1 M) was treated with Cu(EH)₂ (3 equiv) and Cs₂CO₃ (1 equiv). The mixture was heated to 120 °C for 24 h in a pressure tube. ^{*b*} Yields refer to the sum of products isolated by chromatography on SiO₂. The remainder of the material was either starting olefin or olefin-isomerized starting material. ^{*c*} Heated to 190 °C. ^{*d*} Reaction run for 72 h. ^{*e*} *tert*-Butyl benzene was used as solvent.

its benzyl amide counterpart **19** (Table 4, entries 1 and 2). Thus, the imide is more reactive than the alkyl amide. Substrates with geminally disubstituted carbon backbones cyclized more efficiently due to the Thorpe–Ingold effect (compare entries 6 and 7, Table 4).⁶

The aryl amides 22 and 25 were able to undergo the cyclization in an efficient manner albeit with the formation of the hydroamination byproducts (Table 4, entries 3 and 4). In order to reduce hydroamination and maximize carboamination, a less hydrogen-atom-donating solvent, i.e., tert-butyl benzene, was used. Although increased selectivity for carboamination was observed, formation of the hydroamination byproduct still occurred. Regardless, these are the first examples where two fused five-membered rings are formed in the copper(II) carboxylate-promoted intramolecular carboamination reaction. Substrate 28, which lacks the geminal methyl groups in the backbone, only produced hydroamination adduct 29 (Table 4, entry 5). Because of the apparent difficulty in forming a five-membered ring with the aryl substituent7 (vide supra), naphthalene substrates 30 and 32 were investigated. Gratifyingly, both substrates

⁽⁶⁾ Beesley, R. M.; Ingold, C. K.; Thorpe, J. F. J. Am. Chem. Soc. 1915, 107, 1080.

underwent the oxidative cyclization in good yield without the simultaneous formation of the hydroamination byproduct (Table 4, entries 6 and 7).

During the synthesis of naphthalene substrate **30**, an interesting observation was made. When the corresponding amide was treated under the coupling reaction conditions developed by Buchwald et al.⁸ (5 mol % CuI, DMDA, iodonaphthalene, K₃PO₄, in DMF at 120 °C), trace amounts of carboamination (ca. 2-3%) product (**31**) are observed in the ¹H NMR. This occurrence is easily explained by the ability of Cu(I) to disproportionate into Cu(II) and Cu(0).⁹ Thus, in an effort to exploit this reactivity, a one-pot arylation/carboamination reaction was performed. Upon completion of the coupling reaction, 3 equiv of Cu(EH)₂ was added to the reaction mixture. After 24 h the carboamination adduct was obtained in 71% yield (eq 4).



Another interesting observation was made when performing the carboamination reaction on heteroaromatic ring amides. Both the amido furan **34a** and the amido pyrrole **34b** derivatives of 2-allyl aniline were unreactive substrates. The amido thiophene **34c** cyclized in 43% yielding *two* carboamination products **35c** and **36c** in a 1:1 ratio (eq 5).

This observation can be rationalized on the basis of the



mechanism shown in Scheme 1. Thus, syn aminocupration followed by homolysis of the resulting unstable organocopper intermediate gives primary radical **A**.^{2f} This radical can either





add 1,6 to give the carboamination product 35c after oxidation of the aryl radical intermediate **B**, or 1,5 (*ipso*), giving rise to carboamination product 36c via intermediate **C**, radical migration, and oxidative re-aromatization as shown in Scheme 1.¹⁰ Rearrangement of intermediate **C** via a carbocation is also possible.

In conclusion, the intramolecular copper(II) carboxylatepromoted carboamination reaction is an efficient method to form a variety of polycyclic lactams. The substrate scope has been expanded from sulfonamides, to include not only aryl amides, but vinyl amides and alkyl imides as well. Furthermore, a simple one-pot arylation/carboamination reaction sequence was achieved. The synthetic utility of this methodolgy and its application in natural product as well as biologically active molecule synthesis is currently ongoing in our lab and will be reported in due course.

Acknowledgment. We thank Mr. Dov Shalman (Northwestern University), Ms. Fatima Sequiera (University at Buffalo, ACS administered PRF Summer Scholar 40968-G1), and Ms. Melantha Jackson (from Utica College, NSF REU Fellowship at SUNY, Buffalo, CHE-0453206) for preliminary studies. This work was supported by the National Institutes of Health (NIGMS R01-GM078383).

Supporting Information Available: Procedures and characterization data and NMR spectra for all new products. This material is available free of charge via the Internet at http://pubs.acs.org.

⁽⁷⁾ Stevens, C. V.; Meenen, E. V.; Masschelein, K. G. R.; Eeckhout, Y.; Hooghe, W.; D'hondte, B.; Nemkyin, V. N.; Zhdankin, V. V. *Tetrahedron Lett.* **2007**, *48*, 7108.

⁽⁸⁾ Klapars, A.; Antilla, J. C.; Huang, X.; Buchwald, S. L. J. Am. Chem. Soc. 2001, 123, 7727.

⁽⁹⁾ Cotton, F. A.; Wilkinson, G.; Murillo, C. A.; Bochmann, M. Advanced Inorganic Chemistry; John Wiley & Sons: New York, 1999.

OL702401W

^{(10) (}a) Studer, A.; Bossart, M. *Tetrahedron* **2001**, *57*, 9649. (b) Guindeuil, S.; Zard, S. Z. *Chem. Commun.* **2006**, 665. (c) Gagosz, F.; Moutrille, C.; Zard, S. Z. *Org. Lett.* **2002**, *4*, 2707. (d) Kyei, A. S.; Tchabanenko, K.; Baldwin, J. E.; Adlington, R. M. *Tetrahedron Lett.* **2004**, *45*, 8931.