This article was downloaded by: [Nova Southeastern University] On: 31 December 2014, At: 19:12 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/lncn20</u>

Synthesis of Carbocyclic Analogues of Thymidine

O. R. Ludek ^a & C. Meier ^{a b}

^a Institute of Organic Chemistry, University of Hamburg, Hamburg, Germany

^b Universität Hamburg, Institut für Organische Chemie, Martin-Luther-King-Platz 6, D-20146, Hamburg, Germany

Published online: 31 Aug 2006.

To cite this article: O. R. Ludek & C. Meier (2003) Synthesis of Carbocyclic Analogues of Thymidine, Nucleosides, Nucleotides and Nucleic Acids, 22:5-8, 683-685, DOI: <u>10.1081/NCN-120022700</u>

To link to this article: <u>http://dx.doi.org/10.1081/NCN-120022700</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

NUCLEOSIDES, NUCLEOTIDES & NUCLEIC ACIDS Vol. 22, Nos. 5–8, pp. 683–685, 2003

Synthesis of Carbocyclic Analogues of Thymidine

O. R. Ludek and C. Meier*

Institute of Organic Chemistry, University of Hamburg, Hamburg, Germany

ABSTRACT

A new route towards an enantiomerically pure carbocyclic 2'-deoxyribonucleoside precursor was developed. After coupling with a nucleobase the product is easily accessible for further modifications at the 3'-hydroxy group.

INTRODUCTION

Nucleoside-based inhibitors of reverse transcriptase (RT) were the first drugs to be used in the chemotherapy of AIDS. After entering the cell, these substances should be activated to their triphosphates by cellular kinases. Thus, for the most extensively used drug, 3'-azido-3'-deoxythymidine (AZT), whereas phosphorylation is facile, the product (AZTMP) is a very poor substrate for the second kinase, thymidylate kinase (TmpK). Due to the steric demand of the azido group in the 3'-position of AZTMP compared to the 3'-hydroxyl group of the natural substrate, 2'-deoxythymidine monophosphate (dTMP), the structure of the enzyme changes, which causes a 200-fold decrease in the rate of phosphorylation.^[1] Docking experiments, performed on the basis of molecular modelling, led to the assumption that carbocyclic nucleotide analogues of AZTMP should still bind well to the active

683

DOI: 10.1081/NCN-120022700 Copyright © 2003 by Marcel Dekker, Inc.

Downloaded by [Nova Southeastern University] at 19:12 31 December 2014

1525-7770 (Print); 1532-2335 (Online) www.dekker.com

^{*}Correspondence: C. Meier, Universität Hamburg, Institut für Organische Chemie, Martin-Luther-King-Platz 6, D-20146 Hamburg, Germany; Fax: +49 404 2838 2495; E-mail: chris.meier@chemie.uni-hamburg.de.

site of TmpK. Due to the increased flexibility these nucleotides should not cause a structural change of the enzyme. Here, we will present the enantioselective synthesis of carbocyclic analogues of thymidine as well as the corresponding monophosphates by using a convergent synthetic strategy. The carbocyclic moiety is coupled to the heterocyclic base to obtain carbocyclic thymidine (*carba*-dT) **6** as a key intermediate for further variations in the 3'-position.

RESULTS



In our search for a short and efficient convergent synthesis towards carbocyclic 2'-deoxythymidine **6**, we were attracted by the simplicity of forming the carbocyclic moiety in two stages using the method of Biggadike et al.^[2]. The two stereogenic centers in **1** are introduced by an asymmetric hydroboration of a substituted cyclopentadiene, prepared from cyclopentadiene. After protecting the free hydroxygroup in **1** with the benzyl group, the chiral olefin **2** can be hydroborated regio- and stereoselective using 9-BBN in THF affording the β -alcohol **3** in 87% yield. The β -alcohol **3** can easily be transferred into the desired α -alcohol **4** by a Mitsunobu reaction. The coupling of the carbocyclic precursor **4** and the nucleobase was also achieved by a Mitsunobu reaction, using *N3*-benzoylthymidine as the nucleophile.^[3] Removal of the protection groups afforded enantiomerically pure carbocyclic thymidine **6** in 15% overall yield.



Copyright @ 2003 by Marcel Dekker, Inc. All rights reserved

Carbocyclic Analogues of Thymidine

The free hydroxy groups in *carba*-dT $\mathbf{6}$ can now be replaced by groups isoster to the azido group in order to investigate the sterical and/or electronical influence of the substituent in this position on the substrate properties towards thymidylate kinase.

REFERENCES

- 1. Lavie, A.; Schlichting, I.; Vetter, I.R.; Konrad, M.; Reinstein, J.; Goody, R.S. Nature Med. **1997**, *3*, 922.
- 2. Biggadike, K.; Borthwick, A.D.; Evans, D.; Exall, A.M.; Kirk, B.E.; Roberts, S.M.; Stephenson, L.; Youds, P. J. Chem. Soc., Perkin Trans. 1 **1988**, 549.
- 3. Bonnal, C.; Chavis, C.C.; Lucas, M. J. Chem. Soc., Perkin Trans. 1 1994, 1401.

685



