## Stable spirocyclic neutral radicals: aluminium and gallium boraamidinates<sup>†</sup>

Tristram Chivers,<sup>\**a*</sup> Dana J. Eisler,<sup>*a*</sup> Chantall Fedorchuk,<sup>*a*</sup> Gabriele Schatte,<sup>*a*</sup> Heikki M. Tuononen<sup>*a*</sup> and René T. Boeré<sup>*b*</sup>

Received (in Berkeley, CA, USA) 5th May 2005, Accepted 2nd June 2005 First published as an Advance Article on the web 11th July 2005 DOI: 10.1039/b506253e

Stable dark red (M = Al) or dark green (M = Ga) neutral radicals {[PhB( $\mu$ -N<sup>t</sup>Bu)<sub>2</sub>]<sub>2</sub>M} are obtained by the oxidation of their corresponding anions with iodine, and EPR spectra supported by DFT calculations show that the spin density is equally delocalized over all four nitrogen atoms in these spiroconjugated systems.

The study of radicals of the heavier main-group elements is a fascinating undertaking that, through a combination of X-ray structural studies, EPR spectroscopy and theoretical calculations, provides informative insights into the bonding arrangements in odd-electron systems.<sup>1</sup> In Group 13 chemistry, many of the so-called stable radicals<sup>2</sup> are obtained as charged species, that is, in salts as either anion or cation radicals. The neutral paramagnetic complexes of the type [M(dbdab)<sub>2</sub>]<sup>•</sup> (M = Al, Ga; dbdab = 1,4-di-*tert*-butyl-1,4-diazabutadiene) have been formulated as (dbdab<sup>-•</sup>)M(dbdab<sup>2–</sup>)M(III) complexes, in which the spin is located on one of the dbdab ligands, on the basis of EPR<sup>3a</sup> and UV-PES spectra.<sup>3b</sup>



The boraamidinate ligand  $[RB(NR')_2]^{2-}$  (A)<sup>4</sup> is isoelectronic with the extensively studied amidinate anions  $[RC(NR')_2]^-$  (B).<sup>5</sup> Early work on ligands of the type A was limited primarily to complexes of Group 4 or Group 14 metals.<sup>4</sup> Interest in these dianionic ligands has been rekindled recently through reports of (a) complexes in which the ligand A bridges very short M=M triple bonds (M = Mo, W),<sup>6a</sup> (b) trisubstituted octahedral Group 4 dianions,<sup>6b</sup> and (c) the spirocyclic Group 13 anions {[PhB( $\mu$ -N'Bu)\_2]<sub>2</sub>M}<sup>-</sup> (1b, M = Ga; 1c, M = In).<sup>7</sup> The first

Fax: +1 (403) 289-9488; Tel: +1 (403) 220-5741

<sup>b</sup>Department of Chemistry and Biochemistry, University of Lethbridge, Lethbridge, Alberta, Canada T1K 3M4. E-mail: boere@uleth.ca; Fax: +1 (403) 329-2057; Tel: +1 (403) 329-2045 structural characterizations of lithium derivatives of **A**, which form dimeric or trimeric clusters, have also been described.<sup>8</sup> An intriguing feature of the latter reagents is the formation of coloured solutions upon oxidation, which are thought to contain anion radicals of the type  $[RB(NR')_2]^{-.9}$  We report here the first examples of the stabilization of these paramagnetic chelating ligands, which is achieved *via* coordination to Group 13 metal centres in the neutral spirocyclic radicals { $[PhB(\mu-N'Bu)_2]_2M$ } (2a, M = Al; 2b, M = Ga).

Treatment of one equivalent of the dilithiated boraamidinate Li<sub>2</sub>[PhB( $\mu$ -N<sup>t</sup>Bu)<sub>2</sub>] with one equivalent of {[PhB( $\mu$ -N<sup>t</sup>Bu)<sub>2</sub>]-AlCl(OEt<sub>2</sub>)} (**3**)<sup>10</sup> in a boiling mixture of benzene and diethyl ether produces the aluminium boraamidinate complex { $\mu$ -Li(OEt<sub>2</sub>)[PhB( $\mu$ -N<sup>t</sup>Bu)<sub>2</sub>]<sub>2</sub>Al} (**4a**).† Subsequent reaction of either **4a** or { $\mu$ -Li(OEt<sub>2</sub>)[PhB( $\mu$ -N<sup>t</sup>Bu)<sub>2</sub>]<sub>2</sub>Ga}<sup>7</sup> (**4b**) with iodine immediately generates dark red or dark green solutions, respectively (Scheme 1).† Dark red crystals of **2a** and dark green crystals of **2b** suitable for X-ray structural determinations‡ were grown from concentrated diethyl ether solutions at 258 K. These radicals are stable in the solid state under an inert atmosphere at room temperature for weeks.

X-Ray structural determinations revealed that complexes 2a and 2b are isostructural. The molecular structures and pertinent structural parameters for 2a and 2b are depicted in Fig. 1. In each case, the molecule lies on a crystallographic two-fold axis which imposes crystallographic equivalence on the two boraamidinate ligands. The two planes defining these ligands are mutually orthogonal [2a:  $\angle$  89.7(2)°; 2b:  $\angle$  89.2(2)°]. The four-membered BN<sub>2</sub>M rings in 2a and 2b are essentially planar. The metrical parameters of the boraamidinate ligands in 2a and 2b are identical within experimental error. For example, the boron and nitrogen atoms all exhibit distorted trigonal planar geometries; the sum of the bond angles is close to  $360^{\circ}$  in each case [2a: B(1),  $359.9(2)^{\circ}$ ; N(1), 359.9(2)°; N(2), 360.0(2)°. **2b**: B(1), 360.1(2)°; N(1), 360.0(2)°; N(2), 359.8(2)<sup>°</sup>]. The geometry about the central metal atom is distorted tetrahedral (bond angle range 76.37(9)-128.5(1)°, 2a;  $72.81(9)-130.9(1)^{\circ}$ , **2b**). In each of the structures, the two M–N and all B-N bond distances are equal within the limit of experimental



Scheme 1 Oxidation of 4a and 4b with iodine.

<sup>&</sup>lt;sup>a</sup>Department of Chemistry, University of Calgary, Calgary, Alberta, Canada T2N 1N4. E-mail: chivers@ucalgary.ca;

<sup>†</sup> Electronic supplementary information (ESI) available: Details of the syntheses of **2a**, **2b**, **3** and **4a**, computational details and Figure S1. See http://dx.doi.org/10.1039/b506253e



Fig. 1 X-Ray crystal structure of **2a** (M = Al) and **2b** (M = Ga) (Hatoms are omitted): Selected bond lengths [Å] and bond angles [°] [PBE0/ TZVP optimized parameters are in square brackets]. **2a**: Al(1)–N(1) 1.842(2) [1.859], Al(1)–N(2) 1.852(2), N(1)–B(1) 1.445(3) [1.442], N(2)–B(1) 1.449(3), N(1)–Al(1)–N(1)\* 128.5(1) [128.2], N(1)–Al(1)–N(2) 76.37(9) [76.3], N(1)\*–Al(1)–N(2) 127.99(9), N(2)–Al(1)–N(2)\* 128.2(1), N(1)– B(1)–N(2) 104.2(2) [105.6]; **2b**: Ga(1)–N(1) 1.923(2) [1.928], Ga(1)–N(2) 1.922(2), N(1)–B(1) 1.445(3) [1.437], N(2)–B(1) 1.438(3), N(1)–Ga(1)– N(1)\* 130.9(1) [130.4], N(1)–Ga(1)–N(2) 72.81(9) [72.8], N(1)\*–Ga(1)– N(2) 130.02(9), N(2)–Ga(1)–N(2)\* 130.6(1), N(1)–B(1)–N(2) 104.7(2) [105.5]. Symmetry transformations used to generate equivalent atoms: \* -x, y,  $-z + \frac{1}{2}$ .

error. As expected from the relative sizes of  $M^{3+}$  ions,<sup>11</sup> the Al–N distances in **2a** are *ca.* 0.08 Å shorter than the corresponding Ga–N distances in **2b**.

Theoretical calculations<sup>†</sup> were first carried out on the  $D_{2d}$  symmetric diamagnetic model systems {[PhB( $\mu$ -NMe)\_2]\_2M}<sup>-</sup> (M = Al, Ga) to evaluate their frontier orbitals. For both M = Al and Ga, the HOMO transforms as the  $a_2$  irreducible representation in the  $D_{2d}$  point group (Figure S1). Thus, a oneelectron oxidation will yield the corresponding neutral radicals {[PhB( $\mu$ -NMe)\_2]\_2M} with an <sup>2</sup>A<sub>2</sub> ground state and  $D_{2d}$  symmetry.

Similar to the model systems, the spirocyclic radicals 2a and 2b are also expected to retain the  $D_{2d}$  symmetry of their parent diamagnetic anions { $[PhB(\mu-N'Bu)_2]_2M$ }<sup>-</sup> (M = Al, Ga). This is confirmed by X-ray structural analyses of 2a and 2b (vide supra) that reveal only slight deviations from idealized structures (Fig. 1). The retention of the  $D_{2d}$  symmetry in **2a** and **2b** is in sharp contrast to the observations for the related diazabutadiene radicals  $[M(dbdab)_2]$  (M = Al, Ga) (vide supra)<sup>3</sup> in which the two dbdab ligands exhibit distinctly different metrical parameters. This structural feature was explained recently by using DFT calculations,<sup>12</sup> which revealed that the HOMOs of the  $D_{2d}$  symmetric diamagnetic anions  $[M(dbdab)_2]^-$  (M = Al, Ga) are doubly degenerate orbitals of e-symmetry. Upon oxidation, an unstable partially filled degenerate set of orbitals is formed that incurs a Jahn-Teller distortion. Hence, the resulting radicals adopt  $C_{2\nu}$ symmetry with mixed-valent dbdab ligands and localized spin density.<sup>12</sup> In comparison, the diamagnetic anions in **4** are also  $D_{2d}$ symmetric, however, the HOMO of these systems is not comprised of a degenerate set of orbitals. Thus, neither Jahn-Teller distortion nor spin localization are of consequence upon oxidation.

Single point calculations for the model radicals  $\{[PhB(\mu-NMe)_2]_2M\}$  (M = Al, Ga) in their optimized geometries show that their SOMOs consist solely of nitrogen p-orbitals and are equally delocalized over all four nitrogen atoms (Fig. 2a). Thus the conditions for spiroconjugation are fulfilled.<sup>13</sup> Similarly, Mulliken population analysis reveals the spin density to be equally



Fig. 2 (a) The SOMO of {[PhB( $\mu$ -NMe)<sub>2</sub>]<sub>2</sub>Al} drawn at isosurface level  $\pm$  0.05. (b) Spin density map of {[PhB( $\mu$ -NMe)<sub>2</sub>]<sub>2</sub>Al} drawn at isosurface levels 0.02 ( $\alpha$ -spin density, red) and -0.002 ( $\beta$ -spin density, blue).

distributed among all nitrogen centres. Although the SOMOs do not include contributions from boron or from the central metal, these atoms also have non-zero spin density values arising from the spin polarization effects (Fig. 2b). In consequence, the EPR spectra of both spirocyclic radicals are expected to show hyperfine structure due to the central metal (**2a**: <sup>27</sup>Al, I = 5/2, 100%; **2b**: <sup>69</sup>Ga, I = 3/2, 60.4%; <sup>71</sup>Ga, I = 3/2, 39.6%), two equivalent boron atoms (<sup>10</sup>B, I = 3, 19.6%; <sup>11</sup>B, I = 3/2, 80.4%) and four equivalent nitrogen centres (<sup>14</sup>N, I = 1, 99.6%).

Figs. 3a and 4a show the experimental EPR spectra of **2a** and **2b**, respectively. Both radicals give intensely coloured solutions (*vide supra*) in diethyl ether or benzene that persist for days (M = AI) or several hours (M = Ga) at room temperature. Excellent simulations of the spectra were obtained by using the hyperfine coupling (hfc) constants given in Table 1. In general, there is a very good agreement between the experimental and the calculated hfc values (Table 1).<sup>14</sup> Hence, the spectral simulations



Fig. 3 Experimental (a) and simulated (b) X-band EPR spectra of a diethyl ether solution of **2a** at 295 K.



**Fig. 4** Experimental (a) and simulated (b) X-band EPR spectra of a diethyl ether solution of **2b** at 295 K.

 Table 1
 Experimental<sup>a</sup> and calculated hyperfine coupling constants
 [G]

$\{[PhB(\mu\text{-}NR)_2]_2M\}$	Isotope	n <sup>b</sup>	Exptl. $(R = {}^{t}Bu)$	Calc.c(R = Me)
$\mathbf{M} = \mathbf{A}\mathbf{l}^d$	<sup>27</sup> A1	1	11.47	16.86
	<sup>14</sup> N	4	4.66	4.17
	${}^{10}B$	2	1.72	2.01
	${}^{11}B$		5.15	6.03
$M = Ga^{e}$	<sup>69</sup> Ga	1	28.00	45.61
	<sup>71</sup> Ga		35.40	57.67
	$^{14}N$	4	4.69	4.27
	${}^{10}B$	2	1.73	2.03
	${}^{11}B$		5.20	6.10

<sup>*a*</sup> Parameters were determined from simulations optimized to match the experimental spectra; very close agreement between experiment and simulation was obtained (Figs. 3 and 4). <sup>*b*</sup> Number of nuclei. <sup>*c*</sup> Details of computational methods used are given in the supplementary information. <sup>*d*</sup> 100% Lorentzian line shapes with a LW of 1.05 G. <sup>*e*</sup> 100% Lorentzian line shapes with a LW of 1.30 G.

and DFT calculations indicate uniform spin delocalization throughout both boraamidinate ligands in **2a** and **2b** and confirm the retention of spirocyclic structures in solution.

In summary, the neutral radicals 2a and 2b are the first examples of complexes in which a boraamidinate anion radical  $[PhB(N'Bu)_2]^{-}$  is stabilized by coordination to a metal centre. They also provide an unique example of spiroconjugation in a

purely inorganic framework.<sup>16</sup> The isolation of these paramagnetic complexes suggests that redox behaviour may be a more important feature of boraamidinate complexes than heretofore recognized.

## Notes and references

‡ Crystal data for **2a**: C<sub>28</sub>H<sub>46</sub>AlB<sub>2</sub>N<sub>4</sub>,  $M_r = 487.29$ , orthorhombic, *Pbcn*, a = 17.252(4), b = 8.911(2), c = 20.212(4) Å, V = 3107.1(1) Å<sup>3</sup>, Z = 4, T = 173(2) K,  $\mu$ (Mo-K $\alpha$ ) = 0.086 mm<sup>-1</sup>, 15287 reflections collected, 2712 unique ( $R_{int} = 0.068$ ),  $R_1 = 0.0577$  [ $I > 2\sigma(I$ ]],  $wR_2 = 0.1468$  (all data). **2b**: C<sub>28</sub>H<sub>46</sub>GaB<sub>2</sub>N<sub>4</sub>,  $M_r = 530.03$ , orthorhombic, *Pbcn*, a = 17.173(3), b = 8.945(2), c = 20.337(4) Å, V = 3123.8(11) Å<sup>3</sup>, Z = 4, T = 173(2) K,  $\mu$ (Mo-K $\alpha$ ) = 0.902 mm<sup>-1</sup>, 49170 reflections collected, 2756 unique ( $R_{int} = 0.056$ ),  $R_1 = 0.0395$  [ $I > 2\sigma(I$ ]],  $wR_2 = 0.0926$  (all data). CCDC 267896 and 267897. See http://dx.doi.org/10.1039/b506253e for crystallographic data in CIF or other electronic format.

- 1 For a recent review, see: P. P. Power, Chem. Rev., 2003, 103, 789-809.
- 2 As defined in ref. 1, the term "stable", as applied to radicals, is taken to indicate "a species that can be isolated and shows no sign of decomposition under an inert atmosphere at room temperature", whereas a "persistent" radical has "a relatively long lifetime under the conditions it is generated".
- 3 (a) W. Kaim and W. Matheis, *Chem. Commun.*, 1991, 597–598 and references therein; (b) F. G. N. Cloke, C. I. Dalby, P. J. Daff and J. C. Green, *J. Chem. Soc., Dalton Trans.*, 1991, 181–184.
- 4 For a short review, see: P. Blais, J. K. Brask, T. Chivers, C. Fedorchuk and G. Schatte, *ACS Symp. Ser.*, ed. P. J. Shapiro and D. A. Atwood, 2002, vol. 822, pp. 195–207.
- 5 For reviews, see: (a) F. T. Edelmann, Coord. Chem. Rev., 1994, 137, 403–481; (b) J. Barker and M. Kilner, Coord. Chem. Rev., 1994, 133, 219–300.
- 6 (a) D. R. Manke, Z.-H. Loh and D. G. Nocera, *Inorg. Chem.*, 2004, 43, 3618–3624; (b) D. R. Manke and D. G. Nocera, *Inorg. Chem.*, 2003, 42, 4431–4436.
- 7 T. Chivers, C. Fedorchuk, G. Schatte and M. Parvez, *Inorg. Chem.*, 2003, 42, 2084–2093.
- 8 J. K. Brask, T. Chivers and G. Schatte, Chem. Commun., 2000, 1805–1806.
- 9 T. Chivers, C. Fedorchuk, G. Schatte and J. K. Brask, *Can. J. Chem.*, 2002, 80, 821–831.
- 10 The monocyclic structure of **3** was confirmed by X-ray crystallography. Details will be given in a full account of this work.
- 11 R. D. Shannon, Acta Crystallogr., Sect. A, 1976, A32, 751-767.
- 12 W. W. Schoeller and S. Grigoleit, J. Chem. Soc., Dalton Trans., 2002, 405–409.
- 13 H. E. Simmons and T. Fukunaga, J. Am. Chem. Soc., 1967, 89, 5208–5215.
- 14 The largest deviations occur for the metal centres and arise from two primary sources. Firstly, the hfc values of metal centres arise purely from spin polarization effects, which are known to be extremely difficult to model accurately.<sup>15</sup> Secondly, the basis set used for gallium is energy-optimized, not property-optimized, and does not include any enhanced s-part, a requirement for obtaining more than semi-quantitative accuracy<sup>15</sup>.
- 15 M. L. Munzarová, in *Calculation of NMR and EPR parameters: Theory and Applications*, ed. M. Kaupp, M. Bühl and V G. Malkin, VCH, Weinheim, 2004, pp. 463–482.
- 16 A spirophenalenyl-based radical with a central tetrahedral BO<sub>2</sub>N<sub>2</sub> unit has been reported, but the spin density is highly delocalized onto the phenalene rings. X. Chi, M. E. Itkis, B. O. Patrick, T. M. Barclay, R. W. Reed, R. T. Oakley, A. W. Cordes and R. C. Haddon, *J. Am. Chem. Soc.*, 1999, **121**, 10395–10402.