

Desymmetrization of Metallated Cyclohexadienes with Chiral *N*-*tert*-Butanesulfinyl Imines


Modhu Sudan Maji, Roland Fröhlich, and Armido Studer*

NRW Graduate School of Chemistry and Fachbereich Chemie, Organisch-Chemisches Institut, Westfälische Wilhelms-Universität, Corrensstrasse 40, 48149 Münster, Germany

studer@uni-muenster.de

Received March 3, 2008

ABSTRACT



This communication describes the desymmetrization of various achiral metallated cyclohexadienes with a series of chiral *N*-*tert*-butanesulfinyl imines. Depending on the metal used, either the symmetrical diene (dicyclohexadienyl-zinc) or the desymmetrized diene (cyclohexadienyl-MgCl) is obtained in a good regioselectivity with excellent diastereoselectivity. The products formed should be useful building blocks for natural product synthesis. The symmetrical 1,4-dienes are readily oxidized to the corresponding diarylmethylamine derivatives.

The desymmetrization of 1,4-cyclohexadienes is a highly useful route to interesting chiral building blocks for natural product synthesis. Ionic and radical C–C bond forming reactions, cycloadditions, and transition-metal mediated processes have been successfully used to achieve that goal.¹

We have recently shown that the chiral cyclohexadienyl-Ti complex **1** reacts highly stereoselectively with various aldehydes to the desymmetrized cyclohexadienes **2**.² More recently, a catalytic variant was developed using silylated cyclohexadiene **3**³ in combination with a chiral Cu-catalyst.⁴ So far, our research in this area focused on the study of chiral cyclohexadienyl nucleophiles. In the present communication, we describe the desymmetrization of achiral metallated cyclohexadienes with the aid of a chiral electrophile. As

promising electrophiles, we decided to study chiral *N*-*tert*-butanesulfinyl imines **4**, which have often been used in stereoselective synthesis^{5,6} to give dienes of type **5** (Scheme 1).

The influence of the metal M on the desymmetrization reaction was studied first. To this end, various cyclohexadienyl metal compounds, which were readily prepared from the corresponding Li-derivative via transmetalation, were reacted with chiral sulfonyl imine **4a** (R = Ph). Along with the desired desymmetrization product **5a**, the symmetrical

(3) For the use of silylated cyclohexadienes in radical chemistry, see: (a) Studer, A.; Amrein, S. *Angew. Chem., Int. Ed.* **2000**, *39*, 3080. (b) Studer, A.; Amrein, S.; Schleth, F.; Schulte, T.; Walton, J. C. *J. Am. Chem. Soc.* **2003**, *125*, 5726. (c) Walton, J. C.; Studer, A. *Acc. Chem. Res.* **2005**, *38*, 794.

(4) Umeda, R.; Studer, A. *Org. Lett.* **2007**, *9*, 2175.


(5) Reviews: (a) Ellman, J. A.; Owens, T. D.; Tang, T. P. *Acc. Chem. Res.* **2002**, *35*, 984. (b) Ellman, J. A. *Pure Appl. Chem.* **2003**, *75*, 39. (c) Zhou, P.; Chen, B.-C.; Davis, F. A. *Tetrahedron* **2004**, *60*, 8003. (d) Morton, D.; Stockman, R. A. *Tetrahedron* **2006**, *62*, 8869.

(6) Some recent examples on allyl metal additions: (a) Li, S.-W.; Batey, R. A. *Chem. Commun.* **2004**, 1382. (b) Foubelo, F.; Yus, M. *Tetrahedron: Asymmetry* **2004**, *15*, 3823. (c) Kolodney, G.; Sklute, G.; Perrone, S.; Knochel, P.; Marek, I. *Angew. Chem., Int. Ed.* **2007**, *46*, 9291.


(1) Reviews: (a) Willis, M. C. *J. Chem. Soc., Perkin Trans. 1* **1999**, 1765. (b) Rahman, A. N.; Landais, Y. *Curr. Org. Chem.* **2002**, *6*, 1369. (c) Hoffmann, R. W. *Synthesis* **2004**, 2075. (d) Schleth, F.; Studer, A. *Synlett* **2005**, 3033.

(2) (a) Schleth, F.; Studer, A. *Angew. Chem., Int. Ed.* **2004**, *43*, 313. (b) Schleth, F.; Vogler, T.; Harms, K.; Studer, A. *Chem.–Eur. J.* **2004**, *10*, 4171.


Scheme 1. Desymmetrization of Cyclohexadienes



diene **6a** was formed in larger amounts depending on the metal used. The relative configuration of the major isomer **5a** was unambiguously assigned by X-ray analysis (see Figure 1). For the symmetrical diene **6a** the relative configuration was assigned in analogy to the X-ray structure of its derivative **6b**.



5a



6b

Figure 1. X-ray structures of compounds **5a** and **6b**.

The reactive cyclohexadienyl-Li compound provided the desired desymmetrization product **5a** (50%) along with an inseparable mixture of an unidentified isomer and the symmetrical diene **6a** (Table 1, entry 1). A reversal of the regioselectivity was achieved with the dicyclohexadienyl

Table 1. Reaction of Various Metallated Cyclohexadienes with **4a**

entry	M	temp [°C]	5a [%]	6a [%]
1	Li	-50	50	11 ^a
2	ZnC ₆ H ₇	-50	14 ^a	62
3	ZnC ₆ H ₇	-30 ^b	24 ^a	50
4	ZnC ₆ H ₇	-78 ^c	18 ^a	29 ^a
5	ZnC ₆ H ₇	-50 ^d	58	9
6	ZnCl	-50	13 ^a	55
7	MgCl ^e	-78	69	6
8	MgCl ^f	-78	77	8

^a Isolated as an inseparable mixture with other isomers. ^b Reaction was run for 18 h. ^c Reaction was run for 64 h. ^d Reaction was conducted in Et₂O. ^e Transmetalation with 0.55 equiv MgCl₂. ^f Transmetalation with 1.0 equiv MgCl₂.

zinc derivative.⁷ The symmetrical isomer **6a** was isolated as major compound in 62% yield with a high diastereoselectivity (entry 2).⁸ The diene **5a** was isolated as an inseparable mixture containing the diastereoisomer of **6a** and another unidentified isomer in 14% combined yield. The same reaction at -30 °C afforded a similar result (entry 3). At -78 °C, reaction with **4a** was far slower and a lower selectivity resulted (entry 4). Interestingly, the selectivity was reversed upon switching to Et₂O as a solvent (entry 5). We assume that transmetalation to zinc did not occur under these conditions since the selectivity was similar to the selectivity obtained for the Li-experiment. Reaction of the more Lewis-acidic cyclohexadienyl-ZnCl-derivative with **4a** gave a slightly lower selectivity as compared with the result obtained using the dicyclohexadienyl zinc derivative (entry 6). The addition of cyclohexadienyl-triisopropoxy titanate (M = Ti(O*i*-Pr)₃) to **4a** at -78 °C occurred with low selectivity (6 isomers were identified by GC analysis) and with a low combined yield (32%, not shown in Table 1). At higher temperature, the Ti-cyclohexadienyl derivative was not stable.⁹ Pleasingly, with the cyclohexadienyl-MgCl compound the desired desymmetrized product **5a** was obtained with excellent diastereoselectivity (69% yield, ds > 99%, entry 7). The diene **6a** was isolated in 6% yield as the only detectable side product. The other possible three isomers of the product 1,3-diene were not identified. A slightly lower regioselectivity but higher yield was obtained upon using 1.0 equiv of MgCl₂ in the transmetalation step (entry 8). To


(7) Allyl-zinc compounds have successfully been used in stereoselective allylations: (a) Ren, H.; Dunet, G.; Mayer, P.; Knochel, P *J. Am. Chem. Soc.* **2007**, *129*, 5376. (b) Dunet, G.; Mayer, P.; Knochel, P *Org. Lett.* **2008**, *10*, 117. See also ref. 6c.

(8) An authentic sample of the diastereoisomer of symmetrical diene **6a** was prepared for comparison. In the fraction of the inseparable isomers (14% combined yield), the diastereoisomer of **6a** was identified and the diastereoselectivity was calculated to be 50:1.

(9) Knoop, C. A.; Studer, A. *Adv. Synth. Catal.* **2005**, *347*, 1542.

summarize the screening studies we can state that depending on the metal used, either the symmetrical diene **6a** (cyclohexadienyl-ZnC₆H₇) or the diene **5a** (cyclohexadienyl-MgCl) could be obtained in a good yield with good regioselectivity and excellent diastereoselectivity.

Table 2. Reaction of Cyclohexadienyl-MgCl with Various Imines^a



entry	R	compd	5 [%]	ds (5) [%] ^b	6 [%]
1	4-MeC ₆ H ₄	b	74	>99	4
2	2-MeC ₆ H ₄	c	69 ^c	>99	13
3	4-MeOC ₆ H ₄	d	86	>99	5
4	2-MeOC ₆ H ₄	e	72	>99	8
5	2-furyl	f	77	>99	5
6	<i>trans</i> -PhCH=CH	g	75	>99	5
7	<i>i</i> -Pr	h	79 ^d	>99	4
8	C ₆ H ₁₁	i	74 ^d	>99	9
9	<i>t</i> -Bu	j	58 ^{d,e}	>99	1
10	PhCH ₂	k	48 ^{c,d}	>99	3

^a Conditions: 1,4-cyclohexadiene (1.0 mmol), *s*-BuLi (1.1 mmol), TMEDA (1.1 mmol) in THF at $-78\text{ }^{\circ}\text{C}$ for 1.5 h, then MgCl₂ (0.55 mmol) at $-78\text{ }^{\circ}\text{C}$ for 3 h, imines **4** for 30 min at $-78\text{ }^{\circ}\text{C}$. ^b Diastereoselectivity with respect to the other three possible 1,3-diene isomers. ^c Containing 2% of the diastereoisomer of **6**. ^d The symmetrical diene **6** could not be separated from **5**. Yield was calculated based on the ratio **5** to **6** measured by GC-analysis. ^e The diastereoisomer of **6j** was detected by GC analysis (<1%).

To study the scope and limitations of our new method, various imines were reacted under the optimized conditions with cyclohexadienyl-MgCl (Table 2) and with the dicyclohexadienyl zinc derivative (Table 3). The relative configuration of the products **5b–k** was assigned in analogy to compound **5a**, and the assignment of the relative configuration of products **6a** and **6c–i** is based on the X-ray structure of **6b**.

Table 3. Reaction of Cyclohexadienyl-ZnC₆H₇ with Various Imines^a

entry	R	compd	6 [%]	dr (6)	other isomers [%] ^b
1	4-MeC ₆ H ₄	b	64	30:1	16
2	2-MeC ₆ H ₄	c	49	9:1	24
3	4-MeOC ₆ H ₄	d	59 ^c	11:1	10
4	2-MeOC ₆ H ₄	e	49 ^c	n.d.	18
5	2-furyl	f	32	n.d.	22
6	<i>i</i> -Pr	h	57 ^c	>99:1	9
7	C ₆ H ₁₁	i	58 ^c	>99:1	4


^a Conditions: 1,4-cyclohexadiene (2.0 mmol), *s*-BuLi (2.2 mmol), TMEDA (2.2 mmol) in THF at $-78\text{ }^{\circ}\text{C}$ for 1.5 h, then ZnCl₂ (1.1 mmol) at $-78\text{ }^{\circ}\text{C}$ for 3 h, then imines **4**, at $-50\text{ }^{\circ}\text{C}$ for 18 h to 2 d. ^b Combined yield of other isomers formed. ^c The other isomer(s) could not be separated. Yield calculated based on the ratio determined by GC- or ¹H NMR analysis.

Excellent diastereoselectivities were obtained for the reaction with the *p*-substituted sulfinyl aryl amines **4b** and **4d** to give **5b** and **5d** as major compounds with good yields and high regioselectivities (entries 1,3). The ortho-substituted congeners delivered similar results (entries 2,4). A high regioselectivity was also achieved with the furyl derivative **4f** (entry 5). Nonaromatic sulfinyl imines **4g–k** reacted with cyclohexadienyl-MgCl with excellent diastereoselectivities and good to excellent regioselectivities (entries 6–10).

Compared to the Mg-dienyl complex, the dicyclohexadienyl zinc derivative provided slightly lower selectivities favoring the symmetrical dienes **6** (Table 3). Aromatic *N*-*tert*-butanesulfinyl imines **4b–e** provided the dienes **6b–e** in moderate to good yields with high diastereoselectivities (49–64%, entries 1–4).¹⁰ The 1,3-dienes were formed as a mixture of isomers in 10–24% combined yield. A lower regioselectivity was obtained with the furyl derivative **4f** (entry 5). High regioselectivities, excellent diastereoselectivities and good yields were achieved with the alkylsulfinyl imines **4h** and **4i** (entries 6,7).

As a mechanism to explain the regiodivergent outcome and the stereoselectivities of the reactions, we suggest the following two models. The Grignard reagent probably reacts from its symmetrical cyclohexadienyl-MgCl isomer via a six-membered chair transition state according to model **A** (Scheme 2). The Mg-metal acts as Lewis acid to complex

Scheme 2. Models to Explain the Selectivity




the nitrogen lone pair and the oxygen atom of the sulfinyl group.¹¹ The sulfinyl imine reacts via its *cis* isomer¹² to **5**. Reaction of the dicyclohexadienyl-zinc compound probably occurs via the non symmetrical dienyl metal structure as depicted in **B** to provide **6**. A similar transition state model was recently suggested by Marek for diastereoselective allylations of allyl zinc derivatives to chiral *N*-*tert*-butanesulfinyl imines.^{6c}

(10) An authentic sample of the diastereoisomer of symmetrical diene **6a** was prepared for comparison. The ¹H NMR showed characteristic signals which were used to identify the presence of the diastereoisomers of **6b–d**. Due to signal overlap, the diastereoisomer ratio could not be determined for **6e** and **f**. By GC analysis, the diastereoisomer of **6i** was not identified. Selectivity of **6h** was assigned in analogy.

(11) Tang, T. P.; Ellman, J. A. *J. Org. Chem.* **1999**, *64*, 12.

(12) Stereoselective additions onto *cis*-imines: (a) Corey, E. J.; Decicco, C. P.; Newbold, R. C. *Tetrahedron Lett.* **1991**, *32*, 5287. (b) Roy, U. K.; Roy, S. *Tetrahedron Lett.* **2007**, *48*, 7177.

Scheme 3. DDQ Mediated Oxidation



Finally, we decided to oxidize the product dienes to the corresponding chiral diarylmethylamine derivatives. This important structural unit can be found in biologically active compounds.¹³ In agreement with our previous studies,¹⁴ the 1,3-dienes **5** could not be cleanly oxidized. However, smooth oxidation of the symmetrical 1,4-dienes could be achieved.

(13) Stereoselective syntheses: (a) Hayashi, T.; Ishigedani, M. *J. Am. Chem. Soc.* **2000**, *122*, 976. (b) Plobeck, N.; Powell, D. *Tetrahedron: Asymmetry* **2002**, *13*, 303. (c) Tokunaga, N.; Otomaru, Y.; Okamoto, K.; Ueyama, K.; Shintani, R.; Hayashi, T. *J. Am. Chem. Soc.* **2004**, *126*, 13584. (d) Castagnolo, D.; Giorgi, G.; Spinosa, R.; Corelli, F.; Botta, M. *Eur. J. Org. Chem.* **2007**, 3676.

(14) Umeda, R.; Studer, A. *Org. Lett.* **2008**, *10*, 993.

Hence, treatment of **6b** or **6c** with DDQ in benzene at room temperature afforded the corresponding protected diarylmethylamines **7b** and **7c** in good yields as diastereoisomerically pure compounds (Scheme 3).

In summary, we present a novel approach for the desymmetrization of achiral metallated cyclohexadienes with chiral *N*-tert-butanesulfinyl imines. The reaction of readily generated cyclohexadienyl-MgCl with various *N*-tert-butanesulfinyl imines provided the corresponding desymmetrized 1,3-dienes with excellent diastereoselectivities and high regioselectivities. The regioselectivity could be reversed by using dicyclohexadienyl-Zn. The 1,4-dienes obtained with the Zn-chemistry were readily oxidized to the corresponding diarylmethylamine derivatives.

Acknowledgment. We thank the NRW Graduate School of Chemistry (stipend to M.S.M.), Novartis Pharma AG (Young Investigator Award to A.S.), and the Fonds der Chemischen Industrie for supporting our work.

Supporting Information Available: Experimental details and characterization data for the products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL800478Q