

A $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ Catalyzed Sakurai–Hosomi Allylation of Isatins and Isatin Ketoimines Using Allyltrimethylsilane


Zhong-Yan Cao, Yan Zhang, Cong-Bin Ji, and Jian Zhou*

Shanghai Key Laboratory of Green Chemistry and Chemical Processes, East China Normal University, 3663 N. Zhongshan Road, Shanghai, 200062, P. R. China

jzhou@chem.ecnu.edu.cn

Received October 9, 2011

ABSTRACT



It is reported that $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ could efficiently activate the cheap but less reactive allyltrimethylsilane for the allylation of isatins or isatin ketoimines, with catalyst loading down to 0.1 mol %. This is the first example of Sakurai–Hosomi allylation of ketoimines using allyltrimethylsilane. A rare example of chiral mercury catalysis is also reported.

The exploration of new catalytic properties of cheap and easily available metal salts such as iron, bismuth, and copper salts is of current interest in organic synthesis. In contrast, limited attention was paid to mercury catalysis, mainly because of the worry about its toxicity.¹ This is to some extent perplexing because mercury is still widely used for gauges, barometers, high-intensity discharge lamps, electrodes, electrical switches, and extraction of gold.^{1a} In addition, mercury amalgam is a common tooth filling, and thimerosal is still used in vaccines in many countries.^{1a} No convincing proof revealed that mercury-containing vaccines and mercury amalgams lead to adverse health effects.¹

Recent studies have shown that mercury compounds have interesting properties worthwhile to explore.² For example, Barron reported that a stabilized arene–mercury complex could efficiently catalyze H/D exchange of C_6D_6 with arenes.^{2a,b} Nishizawa had identified $\text{Hg}(\text{OTf})_2$ as a

powerful catalyst for a series of transformations,^{2c,d} which had been applied to natural product synthesis. A recent example was the synthesis of hippuristanol via $\text{Hg}(\text{OTf})_2$ -catalyzed spiroketalization by Deslongchamps.^{2f}

In light of these facts, we believe the discovery of new catalytic properties of mercury, an “element of mystery”,^{1a} is still needed. During our efforts in the synthesis of 3,3-disubstituted oxindoles for biological evaluation,³ we had developed a highly efficient Friedel–Crafts reaction of 3-substituted 3-hydroxyoxindoles and aromatic compounds catalyzed by $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$.^{3d} Its high efficiency results from aromatic mercuration that produces a strong acid to facilitate the generation of carbocationic intermediate and simultaneously forms the more reactive nucleophile aryl mercural. Here, we wish to report that $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ could activate allyltrimethylsilane for a highly efficient allylation of isatins and isatin derived ketoimines.


The Sakurai–Hosomi allylation reaction is very useful for the synthesis of homoallylic alcohols or amines from inexpensive and nontoxic allylsilanes.⁴ Although significant achievements have been made in allylation reactions

(1) (a) Clarkson, T. W.; Magos, L. *Crit. Rev. Toxicol.* **2006**, 36, 609. (b) Doja, A.; Roberts, W. *Can. J. Neurol. Sci.* **2006**, 33, 341.

(2) (a) Borovik, A. S.; Barron, A. R. *J. Am. Chem. Soc.* **2002**, 124, 3743. (b) Branch, C. S.; Barron, A. R. *J. Am. Chem. Soc.* **2002**, 124, 14156. (c) Yamamoto, H.; Sasaki, I.; Hirai, Y.; Namba, K.; Imagawa, H.; Nishizawa, M. *Angew. Chem., Int. Ed.* **2009**, 48, 1244. (d) Namba, K.; Kaihara, Y.; Yamamoto, H.; Imagawa, H.; Tanino, K.; Williams, R. M.; Nishizawa, M. *Chem.—Eur. J.* **2009**, 15, 6560. (e) Ravindar, K.; Reddy, M. S.; Deslongchamps, P. *Org. Lett.* **2011**, 13, 3178. (f) Ravindar, K.; Reddy, M. S.; Lindqvist, L.; Pelletier, J.; Deslongchamps, P. *Org. Lett.* **2010**, 12, 4420. For a review, see: (g) Nishizawa, M.; Imagawa, H.; Yamamoto, H. *Org. Biomol. Chem.* **2010**, 8, 511.

(3) (a) Qian, Z.-Q.; Zhou, F.; Du, T.-P.; Wang, B.-L.; Ding, M.; Zhao, X.-L.; Zhou, J. *Chem. Commun.* **2009**, 6753. (b) Liu, Y.-L.; Wang, B.-L.; Cao, J.-J.; Chen, L.; Zhang, Y.-X.; Wang, C.; Zhou, J. *J. Am. Chem. Soc.* **2010**, 132, 15176. (c) Ding, M.; Zhou, F.; Liu, Y.-L.; Wang, C.-H.; Zhao, X.-L.; Zhou, J. *Chem. Sci.* **2011**, 2, 2035. (d) Zhou, F.; Cao, Z.-Y.; Zhang, J.; Yang, H.-B.; Zhou, J. *Chem.—Asian J.* **2011**, in press.

by Leighton, using strain-release-activated silane reagents, the activation of less reactive allylsilanes for reaction design is still needed.^{4k,l} In this context, two activation models have been adopted: one is to activate the electrophile by an acid to react with allylsilane (eq 1),^{4b} and the other is to activate allylsilanes by a Lewis base to directly react with electrophiles^{4c} or to help the formation of reactive allyl metallic species (eq 2).^{4h–j} Despite achievements, efficient allylation of ketones or ketoimines is very limited, which often relies on the dual activation of both reaction partners and using allyltrimethoxysilane or specially designed allylsilanes that can be easily activated by Lewis base to form reactive allylmetallic species.^{4h–j} As far as we knew, the catalytic allylation of ketoimines using cheaper but less reactive allyltrimethylsilane **1** was not reported yet,⁴ and no catalytic asymmetric allylation of ketones using allyltrimethylsilane was reported.⁴



To activate the less reactive allyltrimethylsilane **1** for reaction development, we considered a new approach: the direct activation of silane **1** by a Lewis acid. It was postulated that the interaction of a carbophilic Lewis acid with the double bond of **1** might form an intermediate carbocation, which was stabilized by hyper-conjugative overlap with the adjacent carbon–silicon bond, followed by the cleavage of the silyl electrofuge to afford a reactive allylic metallic species for further transformations (eq 3). To test this hypothesis, we first analyzed the interaction between silane **1** and Lewis acids (10 mol %) by NMR analysis under N₂. We found that hard Lewis acids such as Sc(OTf)₃ and In(ClO₄)₃·8H₂O have no obvious interaction




Figure 1. NMR analysis of the interaction between selected Lewis acids and silane **1**.

Table 1. Comparing Different Lewis Acids

entry ^a	cat.	×	time (h)	yield ^b (%)
1	Sc(OTf) ₃	2	48	31
2	In(ClO ₄) ₃ ·8H ₂ O	2	48	75
3	Ph ₃ PAuCl + AgOTf	2	48	10
4	Bi(OTf) ₃	2	48	50
5	Hg(OTf) ₂	2	1	99
6	Hg(ClO ₄) ₂ ·3H ₂ O	2	2	95
7	Hg(ClO ₄) ₂ ·3H ₂ O	1	4	98
8 ^c	Hg(ClO ₄) ₂ ·3H ₂ O	0.3	12	93
9 ^d	Hg(ClO ₄) ₂ ·3H ₂ O	0.1	144	71


^a On a 0.40 mmol scale. ^b Isolated yield. ^c On a 7.0 mmol scale. ^d On a 20.0 mmol scale, with 23% of **2a** recovered.

with silane **1** after several hours (Figure 1). Soft Lewis acids such as Ph₃PAuOTf and AgOTf could interact with **1** slowly. However, borderline Lewis acid Bi(OTf)₃ and soft Lewis acid Hg(ClO₄)₂·3H₂O or Hg(OTf)₂ could rapidly interact with silane **1** within 10 min, and the generation of propylene was observed by NMR analysis. Next, we examined the performance of these Lewis acids in the reaction of silane **1** and isatin **2a** in CH₂Cl₂ at 25 °C, with 2 mol % catalyst. Some typical results were shown in Table 1 (for details about NMR studies, screened Lewis acids, and optimization, see the Supporting Information). Of hard Lewis acids screened, In(ClO₄)₃·8H₂O and Sc(OTf)₃, which failed to activate silane **1**, catalyze this reaction slowly (Table 1, entries 1 and 2), possibly through the activation of isatin **2a**. Of Lewis acids interacting with silane **1**, Ph₃PAuOTf and Bi(OTf)₃ catalyze the reaction slowly, giving product **3a** in 10 and 50% yield after 2 days, respectively (Table 1, entries 3 and 4).

In contrast, Hg(OTf)₂ and Hg(ClO₄)₂·3H₂O enable the completion of reaction within 2 h to afford **3a** in 99 and 95% yield, respectively (Table 1, entries 5 and 6). Although Hg(OTf)₂ was more reactive, cheap and easy to handle, Hg(ClO₄)₂·3H₂O⁵ was chosen as the optimum. The catalyst loading could be reduced to 0.3 mol % for a 7.0 mmol scale reaction, which finished within 12 h to give **3a** in 93% yield

(4) For reviews, see: (a) Hosomi, A. *Acc. Chem. Res.* **1988**, *21*, 200. (b) Denmark, S. E.; Fu, J. *Chem. Rev.* **2003**, *103*, 2763. (c) Denmark, S. E.; Beutner, G. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 1560. (d) Shibasaki, M.; Kanai, M. *Chem. Rev.* **2008**, *108*, 2853. Also see: (e) Hosomi, A.; Shirahata, A.; Sakurai, H. *Tetrahedron Lett.* **1978**, *33*, 3043. (f) Wang, D.-K.; Zhou, Y.-G.; Tang, Y.; Hou, X.-L.; Dai, L.-X. *J. Org. Chem.* **1999**, *64*, 4233. (g) Ishihara, K.; Hiraiwa, Y.; Yamamoto, H. *Synlett.* **2001**, 1851. (h) Yamasaki, S.; Fujii, K.; Wada, R.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2002**, *124*, 6536. (i) Wadamoto, M.; Yamamoto, H. *J. Am. Chem. Soc.* **2005**, *127*, 14556. (j) Kamei, T.; Fujita, K.; Itami, K.; Yoshida, J.-I. *Org. Lett.* **2005**, *7*, 4725. For the use of strained silacycles, see: (k) Kinnaird, J. W. A.; Ng, P. Y.; Kubota, K.; Wang, X.; Leighton, J. L. *J. Am. Chem. Soc.* **2002**, *124*, 7920. (l) Berger, R.; Rabbat, P. M. A.; Leighton, J. L. *J. Am. Chem. Soc.* **2003**, *125*, 9596.


(5) For a review about metal perchlorate hydrates, see: Dalpozzo, R.; Bartoli, G.; Sambri, L.; Melchiorre, P. *Chem. Rev.* **2010**, *110*, 3501.

Scheme 1. Allylation of Isatins **2** Using Allyltrimethylsilane **1**^{a,b}^a On a 0.40 mmol scale. ^b Isolated yield.

(Table 1, entry 8). Further lowering the catalyst loading to 0.1 mol % for a 20.0 mmol scale reaction, product **3a** could still be obtained in 71% yield after 6 days, with 23% of **2a** recovered. As far as we know, this is the lowest catalyst loading for the allylation of ketones using allylsilane.

We further examined the substrate scope by running the reaction in CH_2Cl_2 with 1 mol % of $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ (THF was used when unprotected isatins dissolved poorly in CH_2Cl_2). The nature and the position of the substituents on isatin had no obvious influence, and all the desired products **3a–l** were obtained in high yield (Scheme 1). These products are very useful.^{6k} For example, product **3a** can be used for the synthesis of CPC-1,^{6b} donaxaridine, and dioxibrassinine,⁶ⁱ and product **3h** for convolutamydine A, B, and E.⁶ⁱ In the total synthesis of perophoramidine,^{6f} Qin used 6.0 equiv of allylmagnesium bromide to react with 6-bromoisatin to ensure the high yield of product **3d**; our method is obviously a good alternative.

We next tried the $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ catalyzed allylation of isatin derived ketomines **4** using silane **1**. As far as we knew, the use of allyltrimethylsilane **1** for allylation of ketomines was unknown, and the limited reports on Sakurai–Hosomi allylation of ketomines were based on the more reactive but expensive allyltrimethoxysilane or special allylsilanes.^{4h,j} We were pleased to find that different substituted isatin ketimine **4** worked well in MeOH at 40 °C, using 5 mol % of

Scheme 2. Allylation of Isatin Derived Ketimine **4**^{a,b}^a On a 0.40 mmol scale. ^b Isolated yield.

$\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ as the catalyst (Scheme 2). It should also be noted that aminooxindoles **5** were very useful building blocks for a number of bioactive compounds,^{6a} but synthetic methods to 3-allyl-3-aminoxindoles were very limited.⁷

In the NMR studies, we observed that $\text{Bi}(\text{OTf})_3$ could interact with allyltrimethylsilane **1** as rapidly as $\text{Hg}(\text{OTf})_2$, but it catalyzed the allylation of isatin **2** very slowly (entry 4 vs 5, Table 1). During our work, Meshram also reported that the use of 5 mol % $\text{Bi}(\text{OTf})_3 \cdot 4\text{H}_2\text{O}$ was necessary to catalyze the allylation of isatin **2**, with slow addition at -78 °C and then warmed to 20 °C.^{6j} Although less reactive than $\text{Hg}(\text{OTf})_2$, $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ is obviously more efficient than $\text{Bi}(\text{OTf})_3 \cdot 4\text{H}_2\text{O}$ for this reaction, and the catalyst loading could be down to 0.1 mol %.

We further compared both catalysts (Scheme 3). First, $\text{Bi}(\text{OTf})_3$ failed to catalyze the allylation of ketimine **4a** under the same conditions. Although Ollevier reported that the allylation of compound **6** only gave the desired product **7** in 16% yield with carbamate **8** in 36% yield,⁸ even using 5 mol % of $\text{Bi}(\text{OTf})_3 \cdot 4\text{H}_2\text{O}$ and 5.0 equiv of silane **1** in refluxing CH_2Cl_2 , we found that the use of 5 mol % of $\text{Hg}(\text{ClO}_4)_2 \cdot 3\text{H}_2\text{O}$ and only 2.0 equiv of **1** could afford the desired product **7** in 70% yield at room temperature.


The catalytic asymmetric allylation of isatins is of great interest because enantioenriched 3-allyl-3-hydroxyoxindoles are very useful.^{6a} Although Krische have already developed a highly enantioselective allylation of *N*-benzyl isatins using allylacetate,^{6d} the allylation of *unprotected* isatins waited for development,⁶ and only 42% ee was obtained when using tetraallylstannane.^{6b} In the Pd-catalyzed asymmetric allylation of unprotected isatin **2a** using allyl alcohol and Et_3B , Zhou found the nitrogen of the

(6) For a comprehensive review, see: (a) Zhou, F.; Liu, Y.-L.; Zhou, J. *Adv. Synth. Catal.* **2010**, 352, 1381. For methods to 3-allyl 3-hydroxyoxindoles, see: (b) Kitajima, M.; Mori, I.; Arai, K.; Kogure, N.; Takayama, H. *Tetrahedron Lett.* **2006**, 47, 3199. (c) Hanhan, N. V.; Sahin, A. H.; Chang, T. W.; Fetting, J. C.; Franz, A. K. *Angew. Chem., Int. Ed.* **2010**, 49, 744. (d) Itoh, J.; Han, S. B.; Krische, M. J. *Angew. Chem., Int. Ed.* **2009**, 48, 6313. (e) Qiao, X.-C.; Zhu, S.-F.; Zhou, Q.-L. *Tetrahedron: Asymmetry* **2009**, 20, 1254. (f) Wu, H.; Xue, F.; Xiao, X.; Qin, Y. *J. Am. Chem. Soc.* **2010**, 132, 14052. (g) Nair, V.; Jayan, C. N.; Ros, S. *Tetrahedron* **2001**, 57, 9453. (h) Sano, D.; Nagata, K.; Itoh, T. *Org. Lett.* **2008**, 10, 1593. (i) Kawasaki, T.; Nagaoka, M.; Satoh, T.; Okamoto, A.; Ukon, R.; Ogawa, A. *Tetrahedron* **2004**, 60, 3493. (j) Meshram, H. M.; Ramesh, P.; Reddy, C. B.; Kumar, G. S. *Chem. Lett.* **2011**, 40, 357. (k) Alcaide, B.; Almendros, P.; Rodriguez-Acebes, R. *J. Org. Chem.* **2006**, 71, 2346. (l) Sano, D.; Nagata, K.; Itoh, T. *Org. Lett.* **2008**, 10, 1593. (m) Bui, T.; Candeias, N. R.; Barbas, C. F., III *J. Am. Chem. Soc.* **2010**, 132, 5574. (n) Zheng, K.; Yin, C.; Liu, X.; Lin, L.; Feng, X. *Angew. Chem., Int. Ed.* **2011**, 50, 2573.


(7) (a) Lesma, G.; Landoni, N.; Pilati, T.; Sacchetti, A.; Silvani, A. *J. Org. Chem.* **2009**, 74, 4537. (b) Alcaide, B.; Almendros, P.; Aragoncillo, C. *Eur. J. Org. Chem.* **2010**, 2845. (c) Mouri, S.; Chen, Z.; Mitsunuma, H.; Furutachi, M.; Matsunaga, S.; Shibasaki, M. *J. Am. Chem. Soc.* **2010**, 132, 1255. For reviews: (d) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. *Chem. Rev.* **2011**, 111, 2626. (e) Shen, K.; Liu, X.; Lin, L.; Feng, X. *Chem. Sci.* **2011**, 2, No. DOI:10.1039/C1SC00544H.

(8) Ollevier, T.; Li, Z. *Adv. Synth. Catal.* **2009**, 351, 3251.


Scheme 3. Comparing Bi(OTf)₃ with Hg(ClO₄)₂·3H₂O



product was also masked with an allyl group.^{6c} Although chiral mercury catalysis was rare,⁹ we found that at 0 °C in THF, 1.0 mol % of (*S*)-BINAP/Hg(ClO₄)₂·3H₂O could give the *S*-enantiomer of product **3** in excellent yield with up to 63% ee. Although there has much room for improvement, this is the first example of catalytic asymmetric allylation of ketones using allyltrimethylsilane.⁴



To gain more insight into the mechanism, we first tried NMR studies and observed little interaction between Hg²⁺ and *N*-methylisatin, suggesting Hg(ClO₄)₂·3H₂O might not act as a Lewis acid to activate isatin (see the Supporting Information). Since diallylmercury **9** could react with aldehydes in water,¹⁰ we tried the reaction of **9** and **2a**, which worked slowly, and found 1.7 equiv of **9** was needed to ensure 90% yield of product **3a** after 24 h. We also prepared allylmercury triflate **10** or perchlorate **11**, from anion exchange of allylmercury chloride **13** using AgOTf or AgClO₄·H₂O, to react with **2a** but found that no reaction took place.



To probe if diallylmercury **9** was the key intermediate for allylation of isatin, we tried using different Hg(II) sources to mediate the reaction of isatin **2a** with silane **1**. Although the use of 1 mol % of **9** or 2 mol % of TMSOTf failed to initiate the reaction (Table 2, entries 1 and 2), the combination of both resulted in high reactivity (Table 2, entry 3). The possible role of TMSOTf was to cleave the Hg–O bond of adduct **10** to release O-TMS protected product **15** and allylmercury triflate **11**, which further interact with silane **1** to give diallylmercury **9**. Indeed, the in situ prepared triflate **11** or perchlorate **12** initiated the reaction well (Table 2, entries 6 and 7), though allylmercury chloride **13** failed to initiate the reaction (Table 2, entry 5). These results were also in

(9) Only two examples were reported by now: (a) Yamamoto, H.; Ho, E.; Namba, K.; Imagawa, H.; Nishizawa, M. *Chem.—Eur. J.* **2010**, *16*, 11271. (b) Oh, T.; Lopez, P.; Reilly, M. *Eur. J. Org. Chem.* **2000**, 2901.


(10) Hashmi, A. S. K.; Schwarz, L.; Bolte, M. *Tetrahedron Lett.* **1998**, *39*, 8969.

Table 2. Control Experiments

entry	Hg source	additive	time (h)	yield ^a (%)	ee ^b (%)
1	9	no	48	trace	
2	no	TMSOTf ^c	48	trace	
3	9	TMSOTf ^c	2	91	
4	9	TMSOTf ^c and (<i>S</i>)-BINAP ^d	72	84	57
5	13	no	24	trace	
6	13	AgOTf ^e	4	90	
7	13	AgClO ₄ ·H ₂ O ^e	4	91	

^a Isolated yield. ^b By HPLC analysis. ^c 2.0 mol %. ^d 1.0 mol %, in THF at 0 °C. ^e 1.1 mol %; both silver salts could not catalyze the reaction (**13**: allylmercury chloride).

accordance with NMR studies that the in situ generated **11** or **12** could react with silane **1** as quickly as Hg(OTf)₂ or Hg(ClO₄)₂·3H₂O. These results also confirmed that weak-coordinating counteranions such as OTf[−] and ClO₄[−] were key to ensure the high Lewis acidity of Hg(II) to activate silane **1**. A possible catalytic cycle was then proposed as below (for more details and discussion, please see the Supporting Information).



In conclusion, we have demonstrated that Hg(OTf)₂ and Hg(ClO₄)₂·3H₂O could activate allyltrimethylsilane for a highly efficient allylation of isatins and isatin ketoimines. The catalyst loading could be down to 0.1 mol %, the lowest known for a Sakurai–Hosomi allylation reaction. Our results further demonstrated that mercury salts had some interesting catalytic properties worthwhile to explore.^{2,11} Efforts to develop a highly enantioselective version are now in progress in our laboratory.

Acknowledgment. We thank the National Natural Science Foundation of China (20902025, 21172075), Shanghai Pujiang Program (10PJ1403100), and 973 Program (2011CB-808600) for the financial support, which is highly appreciated.

Supporting Information Available. Experimental procedures and characterizations, copies of ¹H NMR and ¹³C NMR of new compounds, and HPLC profiles. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(11) As suggested by one reviewer, we checked the Hg contamination level of product **3a** by both CVAFS and ICP-MS analysis, which gave consistent results that ca. 3.0% of Hg of catalyst remained in product **3a**. For details, please see the Supporting Information.