



J. Serb. Chem. Soc. 77 (6) 733–739 (2012) JSCS–4304 JSCS-info@shd.org.rs • www.shd.org.rs/JSCS UDC 547.78+66.095.252+544.354.3+ 547.233:57–188 Short communication

SHORT COMMUNICATION Synthesis of some 3,5-diaryl-2-isoxazoline derivatives in ionic liquids media

JAVAD SAFAEI-GHOMI* and MOHAMMAD ALI GHASEMZADEH

Department of Chemistry, Qom Branch, Islamic Azad University, Qom, I. R. Iran

(Received 31 August, revised 19 December 2011)

Abstract: Biologically active isoxazoline derivatives were efficiently synthesized in excellent yields and in smaller reaction times using mild, effective and environmentally friendly butylmethylimidazolium bromide as the solvent and catalyst. By use of this catalyst, isoxazoline derivatives were produced *via* cyclization reaction of a chalcone and hydroxylamine hydrochloride in the ionic liquid media. The separation of the product was facile and the catalyst could be separated and recycled. The method is very rapid, safe and avoids the use of hazardous and expensive reagents and solvents.

Keywords: isoxazoline; cyclization reaction; ionic liquids; butylmethylimidazolium bromide; multi-component reactions; biological activities.

INTRODUCTION

Compounds incorporating heterocyclic ring systems continue to attract considerable interest due to the wide range of their biological activities. Amongst them, five-membered heterocyclic compounds occupy a unique place in the realm of natural and synthetic organic chemistry. Isoxazolines as heterocyclic compounds have found wide application as pharmaceutical and agrochemical agents. For instance, isoxazolines possess biological activities,^{1–8} such as insecticidal, antibacterial, antibiotic, antitumour, antifungal, antimicrobial activity, and antiinflammatory and analgesic. In addition, isoxazoline derivatives have played a crucial role in the theoretical development of heterocyclic chemistry and are also used extensively in organic synthesis.^{9,10} Consequently, much attention has been paid to the development of new methodologies for their preparation. Several methods have been reported for the synthesis of isoxazoline derivatives. The synthetic routes for the preparation of isoxazoline derivatives mainly are cyclization reaction of chalcones,^{11–14} 1,3-dipolar cycloaddition of oximes to alkenes,^{15,16}



^{*} Corresponding author. E-mail: safaei@kashanu.ac.ir doi: 10.2298/JSC110831001S

direct transformation of 3-arylpropargyl hydroxylamines hydrochlorides,¹⁷ reaction of alkenes or alkynes with ketones,¹⁸ 1,3-dipolar cycloaddition of nitrile oxides to vinylic compounds¹⁹ and cyclization of *O*-propargylic hydroxylamines.²⁰

In accordance with the significance of the application these compounds, a mild and efficient route to the synthesis of isoxazoline derivatives in ionic liquid media is reported herein.

In the past decade, ionic liquids have received substantial attention in organic synthesis because of their environmentally benign nature, high polarity, and good thermal stability. Often shorter reaction times, high yields, cleaner reaction products and high selectivity are obtained from ionic-liquid reaction media.^{21,22} Ionic liquids (ILs) have attracted increasing interest recently in the context of green organic synthesis. Although ionic liquids were initially introduced as alternative green reaction media because of their unique chemical and physical properties of non-volatility, non-inflammability, negligible vapor pressure, reusability and high thermal stability, today they have marched far beyond this boundary, showing their significant role in controlling reactions as solvents or catalysts.^{23,24}

There are many reports concerning the applications of ionic liquids as solvents and catalysts in organic reactions, such as Friedel–Crafts reaction,²⁵ Diels– –Alder reaction,^{26,27} Biginelli reaction,²⁸ Beckmann rearrangement,²⁹ Michael reaction,³⁰ reduction of aldehydes,³¹ electophilic substitution of aromatic rings,^{32,33} 1,3-dipolar cycloaddition^{34,35} and other reactions.^{36–38}

In the continuation of ongoing studies on the application of IL media, it was found that dialkylimidazolium halide derivatives have many advantages over conventional solvents.^{39–42} These reagents are safe, easy to handle, environment-tally benign and present fewer disposal problems. Therefore, butylmethylimidazolium bromide ([bmim]Br) (Scheme 1) could be an excellent candidate for employment in organic reactions. The advantage of these methods over conventional methods is that they provide greater selectivity, enhanced reaction rates, cleaner products and manipulative simplicity. According to the development of methods that consist of using environmentally friendly reagents and in continuation of our ongoing program to develop environmentally benign methods, the use of an ionic liquid in the synthesis of isoxazoline derivatives in a mild, facile and clean manner is reported herein.



Scheme 1. 1-Butyl-3-methylimidazolium bromide ([bmim]Br).

EXPERIMENTAL

Butylmethylimidazolium bromide ([bmim]Br) was purchased from Fluka. All melting points are uncorrected and were determined in a capillary tube using a Boetius melting point

Available online at www.shd.org.rs/JSCS/



734

microscope. The Fourier transform infrared (FTIR) spectra were recorded on a Nicolet Magna 550 spectrometer (KBr). The ¹H- and ¹³C-NMR spectra were obtained on a Bruker 400 MHz spectrometer in CDCl₃ as solvent using tetramethylsilane (TMS) as an internal standard. All reactions were monitored and checked by thin layer chromatography (TLC) using hexane//ethyl acetate (8:2) and the spots were examined using a UV lamp. The elemental analyses (C, H, N) were obtained from a Carlo ERBA Model EA 1108 analyzer. The mass spectra were recorded on a Joel D-30 instrument at an ionization potential of 70 eV.

General procedures for the synthesis of chalcones 1a-j

Chalcone derivatives have been prepared by condensation of various aromatic aldehydes and acetophenones in alkaline ethanol according to the procedure reported by Nielsen *et al.*⁴³

To a mixture of NaOH (2.5 g) in 20 ml water and 15 ml EtOH, was added 0.050 mol of the required acetophenone derivative and the solution was stirred at 0-5 °C for 10 min, then 0.050 mol of the required aromatic aldehyde was added and the reaction mixture was refluxed for an appropriate time. After completion of reaction, as indicated by TLC using hexane/ethyl acetate (7:3) the residue was filtered, washed with water and recrystallyzed from ethanol.

Typical procedures for the synthesis of isoxazolines **3a-j**

Conventional heating. Hydroxylamine hydrochloride (0.010 mol) was dissolved in water (5 ml) and added to a solution of chalcone (0.01 mol) in pyridine (10 ml). The reaction mixture was refluxed (magnetic stirring) on oil bath for 3 h. After completion of the reaction, observed by TLC using hexane/ethyl acetate (8:2), the mixture was cooled to room temperature, poured into ice-cold water and then acidified with dilute acetic acid. The obtained solid was filtered, washed with water and recrystallyzed from ethanol.

[Bmim]Br method. A mixture of chalcone (0.010 mol), hydroxylamine hydrochloride (0.010 mol) and [bmim]Br (0.76 g, 0.0035 mol) was stirred at room temperature for the appropriate period After completion of the reaction, as indicated by TLC using hexane/ethyl acetate (8:2), the reaction mixture was extracted with diethyl ether (3×10 ml). The organic layer was evaporated under reduced pressure and the solid residue was recrystallyzed from ethanol. The immiscible ionic liquid phase was recovered and heated for 3 h under vacuum for further use. The best results were obtained with a molar ratio of 1:1:0.35 of chalcone, hydroxylamine hydrochloride and [bmim]Br.

Data of synthesized compounds are are given in Supplementary material.

RESULTS AND DISCUSSION

In this research, the ring closure reaction of chalcone **1** and hydroxylamine hydrochloride **2** occurred under I) conventional and II) in the presence of [bmim]Br to afford the isoxazoline derivatives **3** (Scheme 2). The reactions were performed using α,β -unsaturated carbonyl compounds with diverse substituents. The results of the experiments are summarized in Table I.

A possible mechanism for this reaction, proposed based on our experimental results together with some literature data for the cyclization reaction of chalcone, is that it is realized in two steps; first nucleophilic attack of the carbonyl group by the NH_2 moiety occurs, which is followed by oxime formation and then intramolecular cyclization leads to the five-membered ring products.^{12,13}



SAFAEI-GHOMI and GHASEMZADEH



Scheme 2. Preparation of isoxazolines 3 in [bmim]Br.

Entry	R	R'	Product	M.p., °C	
				Found	Reported
1	Н	Н	3 a	150-152	151-15211
2	CH ₃	Н	3 b	167–168	168^{11}
3	Cl	Н	3c	174–175	175^{11}
4	OCH ₃	Н	3d	149–150	148–151 ¹²
5	NO_2	Br	3e	150-151	150–151 ¹⁵
6	OH	Н	3f	149-150	_a
7	OH	CH ₃	3g	140-142	_a
8	NO_2	CH ₃	3h	121-123	122-123 ¹⁵
9	Br	CH ₃	3i	142-143	142-14315
10	OCH ₃	Br	3ј	168–170	_a

TABLE I. One-pot synthesis of substituted isoxazolines catalyzed by ([bmim]Br)

^aCompounds **3f**, **3g** and **3j** are new products

736

The one-pot condensation of α , β -unsaturated carbonyl compounds and hydroxylamine hydrochloride proceeded in the presence of the ionic liquid smoothly to give the corresponding products in high yields and shorter reaction times in comparison to thermal heating method; the results are summarized in Table II. It was found that [bmim]Br is a fairly good catalytic medium for this reaction. Compared with traditional catalysts, such as AcOH, KOH and NaOH, [bmim]Br not only gives higher yields but also takes much less time and can be easily recycled. Compared with traditional solvents, ionic liquids can be easily reused.

Product	Conventional conditions		Stirred in [bmim]Br		
FIGURE	Time, h	Yield, %	Time, min	Yield, %	
3a	3.5	35	52	63	
3b	3.5	45	50	70	
3c	3	50	48	78	
3d	3	52	46	80	
3e	3	60	45	87	
3f	3	60	44	82	
3g	3	55	52	75	
3h	3	60	42	84	
3i	3	52	44	82	
3ј	3	50	45	80	

TABLE II. Comparison of the two methods for the synthesis of 3,5-diaryl-2-isoxazolines



As mentioned earlier, one of the goals in the use of ionic liquids is to identify and exploit the advantages these compounds can offer over conventional organic solvents besides their greener nature. It has been documented that, compared with classical organic solvents, reactions carried out in ionic liquids often offer enhanced reactivity, better yields and simpler operational process. These advantages were confirmed in the present paper for the synthesis of 3,5-diaryl-2--isoxazoline. The obtained results can be explained by the better catalytic and coordinative properties^{44,45} of this kind of ionic liquids in comparison to other methods.

CONCLUSIONS

In conclusion, the ionic liquid promoted one-pot annulation reactions of chalcones and hydroxylamine hydrochloride and was proved an efficient medium for the synthesis of isoxazoline derivatives. Compared with conventional organic solvents, the use of [bmim]Br had several advantages, *i.e.*, high yields, mild conditions and simpler procedure in shorter reaction times. In addition, the ionic liquid used can be easily recovered and effectively reused at least 5 times more without any significant loss in efficiency.

SUPPLEMENTARY MATERIAL

Spectral data of the compounds are available electronically from http:////www.shd.org.rs/JSCS/, or from the corresponding author on request.

Acknowledgement. The authors gratefully acknowledge the financial support of this work by the Research Affairs Office of the Islamic Azad University, Qom Branch, Qom, I. R. Iran.

ИЗВОД

СИНТЕЗА НОВИХ ДЕРИВАТА 3,5-ДИАРИЛ-2-ИЗОКСАЗОЛИНА У ЈОНСКИМ ТЕЧНОСТИМА

JAVAD SAFAEI-GHOMI и MOHAMMAD ALI GHASEMZADEH

Department of Chemistry, Qom Branch, Islamic Azad University, Qom, I. R. Iran

Биолошки активни деривати изоксазолина синтетисани су у одличном приносу, у присуству бутилметилимидазолијум-бромида као растварача и катализатора. Производи су добијени реакцијом циклизације халкона и хидроксиламин-хидрохлорида. Производи су лако раздвојени, а катализатор је могуће одвојити и поновно користити. Описани поступак је кратак и омогућава да се избегне употреба опасних и скупих реагенаса и растварача.

(Примљено 31. августа, ревидирано 19. децембра 2011)

REFERENCES

- 1. R. Huisgen, Angew. Chem. Int. Ed. Engl. 2 (1963) 565
- 2. P. Caramella, P. Grunanger, 1,3-Dipolar Cycloaddition Chemistry, Vol. 1, Wiley, New York, USA, 1984
- 3. V. M. Barot, Oriental J. Chem. 16 (2000) 177

SAFAEI-GHOMI and GHASEMZADEH

- 4. M. D. Ankhiwala, M. V. Hathi, J. Indian Chem. Soc. 71 (1994) 587
- C. J. Shishoo, M. B. Devani, G. V. Ullas, S. Ananhan, V. S. Bhadti, J. Heterocycl. Chem. 18 (1981) 43
- 6. B. L. Varma, Indian. J. Heterocycl. Chem. 13 (2003) 111
- R. P. Tangallapally, S. D. Rakesh, N. Budha, B. Meibohm, *Bioorg. Med. Chem. Lett.* 17 (2007) 6638
- 8. A. G. Habeeb, P. N. Praveen, R. E. Knaus, J. Med. Chem. 44 (2001) 2921
- Y. V. Tomilovi, G. P. Okonnishnikova, E. V. Shulishov, O. M. Nefedov, *Russ. Chem. Bull.* 44 (1995) 2114
- E. I. Klimova, M. Marcos, T. B. Klimova, A. T. Cecilio, A. T. Ruben, R. R. Lena, J. Organomet. Chem. 106 (1999) 585
- 11. M. Venugopal, P. T. Perumal, Proc. Indian Acad. Sci. (Chem. Sci.). 105 (1993) 19
- 12. M. Kidwai, P. Misra, Synth. Commun. 29 (1999) 3237
- 13. A. Voskienė, V. Mickevičius, Chem. Heterocycl. Compd. 45 (2009) 1485
- 14. T. Shah, V. Desai, J. Serb. Chem. Soc. 72 (2007) 443
- A. Tavares, O. M. S. Ritter, U. B. Vasconcelos, B. C. Arruda, A. Schrader, P. H. Schneider, A. A. Merlo, *Liquid Cryst.* 37 (2010) 159
- R. Maurya, A. Ahmad, P. Gupta, K. Chand, M. Kumar, P. Rawat, N. Rasheed, G. Palit, *Med. Chem. Res.* 20 (2011) 139
- 17. L. Pennicott, S. Lindell, Synlett 3 (2006) 463
- K. I. Itoh, H. Sakamaki, N. Nakazato, A. Horiuchi, E. Horn, C. A. Horiuchi, Synthesis 20 (2005) 3541
- 19. I. N. N. Namboothiri, N. Rastogi, Top. Heterocycl. Chem. 12 (2008) 1
- 20. D. W. Knight, A. J. Proctor, J. M. Clough, Synlett 4 (2010) 628
- 21. H. Olivier-Bourbigou, L. Magna, J. Mol. Catal. 419 (2002) 182
- 22. D. Zhao, M. Wu, Y. Kou, E. Min, Catal. Today 74 (2002) 157
- 23. P. Wasserscheid, T. Welton, *Ionic Liquids in Synthesis*, Wiley-VCH Verlag, Stuttgart, Germany, 2002
- A. P. M. Martins, P. F. Clarissa, N. M. Dayse, N. Zanatta, G. H. Bonacorso, *Chem. Rev.* 108 (2008) 2015
- 25. A. Stark, B. L. MacLean, R. D. Singer, J. Chem. Soc., Dalton Trans. 1 (1999) 63
- 26. T. Fischer, A. Sethi, T. Welton, J. Woolf, Tetrahedron Lett. 40 (1999) 793
- 27. P. Ludley, N. Karodia, Tetrahedron Lett. 42 (2001) 2011
- 28. J. J. Peng, Y. Q. Deng, Tetrahedron Lett. 42 (2001) 5917
- 29. R. X. Ren, L. D. Zueva, W. Ou, Tetrahedron Lett. 42 (2001) 8441
- 30. B. C. Ranu, S. S. Dey, Tetrahedron 60 (2004) 4183
- 31. G. W. Kabalka, R. R. Malladi, Chem. Commun. 22 (2000) 2191
- 32. K. K. Laali, V. J. Gettwert, J. Org. Chem. 66 (2001) 35
- 33. F. Zulfiqar, T. Kitazume, Green Chem. 2 (2000) 296
- 34. J. F. Dubreuil, J. P. Bazureau, Tetrahedron Lett. 41 (2000) 7351
- 35. D. Conti, M. Rodriquez, A. Sega, M. Taddei, Tetrahedron Lett. 44 (2003) 5327
- 36. A. R. Hajipour, Z. Nasreesfahani, A. E. Ruoho, Org. Prep. Proc. Int. 40 (2008) 385
- 37. A. Zare, A. Hasaninejad, M. Shekouhy, A. R. M. Zare, Org. Prep. Proc. Int. 40 (2008) 457
- A. R. Williams, A. J. Angel, K. L. French, D. R. Hurst, D. D. Beckman, C. F. Beam, Synth. Commun. 29 (1999) 1977
- 39. J. Safaei-Ghomi, A. R. Hajipour, J. Chin. Chem. Soc. 56 (2009) 416

Available online at www.shd.org.rs/JSCS/



738

- 40. J. Safaei-Ghomi, M. Taheri, M. A. Ghasemzadeh, Org. Prep. Proc. Int. 42 (2010) 485
- 41. J. Safaei-Ghomi, A. R. Hajipour, M. Esmaeili, Dig. J. Nanomater. Bios. 5 (2010) 865
- 42. J. Safaei-Ghomi, A. R. Hajipour, Org. Prep. Proc. Int. 43 (2011) 372
- 43. A. T. Nielsen, W. J. Houlihan, Org. React. 16 (1968) 1
- 44. Y. Chen, Y. Zu, Y. Fu, X. Zhang, P. Yu, G. Sun, T. Efferth, Molecules 15 (2010) 9486
- 45. S. Kantevari, M. V. Chary, A. P. Rudra Das, S. V. N. Vuppalapati, N. Lingaiah, *Catal. Commun.* 9 (2008) 1575.



Copyright of Journal of the Serbian Chemical Society is the property of National Library of Serbia and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.