# Organic & Biomolecular Chemistry

Cite this: Org. Biomol. Chem., 2011, 9, 7997

www.rsc.org/obc

# Organocatalytic asymmetric Michael-type reaction between $\beta$ , $\gamma$ -unsaturated $\alpha$ -keto ester and $\alpha$ -nitro ketone $\dagger$

Pengfei Li,<sup>*a,b*</sup> Sau Hing Chan,<sup>*b*</sup> Albert S. C. Chan<sup>\**a*</sup> and Fuk Yee Kwong<sup>\**b*</sup>

Received 18th July 2011, Accepted 15th September 2011 DOI: 10.1039/c1ob06191g

A Michael-type reaction of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester and  $\alpha$ nitro ketone was established. With a thiourea catalyst derived from cinchona alkaloid, the reactions afford products in 47– 94% yields with 68–96% ee.

Organocatalysis has been an alternative and/or complementary approach to currently developed organometallic and enzymatic catalysis. Indeed, the emerging of an organocatalytic protocol has displayed great success in modular organic molecule assembly/transformations in the last decade.<sup>1</sup> In particular, organocatalytic asymmetric Michael addition reactions have received widespread attention for accessing a variety of optically active adducts which are synthetically useful building blocks in versatile organic synthesis.<sup>2</sup> Notably, the Michael-type domino reactions efficiently furnish complex chiral natural molecules and diverse intermediates in a one-pot process.<sup>3</sup>

 $\beta$ , $\gamma$ -Unsaturated  $\alpha$ -keto esters represent a resourceful moiety in synthetically useful components. For instance, the asymmetric aldol reactions of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto esters furnish  $\beta$ -hydroxy carbonyl compounds bearing a tertiary alcohol motif.<sup>4</sup> Enantioselective Diels–Alder reactions provide a beneficial route for assembling cyclic products.<sup>5</sup> Asymmetric Michael additions<sup>6</sup> and Michael-type tandem reactions<sup>7</sup> result in  $\gamma$ -functional carbonyl compounds, which can generally be further transformed to obtain the stereospecific molecules.

Recently, we successfully developed an organocatalytic asymmetric aldol reaction of a  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester with a ketone.<sup>8</sup> We found that the aldol reaction, rather than Michael addition or Diels–Alder reaction, was occurred preferentially between the ketone and  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester. Moreover, the size of the ketone substrate affected the product yield and stereo-induction greatly. These outcomes underlined the fact that a sterically more congested ketone could efficiently prevent the aldol reaction of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -ketoesters. Based on the aforementioned findings, we proposed a more active nucleophile with proper steric hindrance that might lead to a Michael-type reaction of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto esters. To examine this idea, a  $\alpha$ -nitro ketone was chosen as the nucleophile and the model reactions of this ketone with a  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester were then investigated.

Very recently, Wang<sup>9a</sup> and Yan<sup>9b</sup> reported the asymmetric organocatalytic Michael/hemiketalization/retro-Henry reaction between a  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester and a  $\alpha$ -nitro ketone, giving more than 90% ee and yield in the presence of the bifunctional Brønsted-base thiourea catalyst.<sup>10</sup> However, the reactions of aliphatic  $\alpha$ -nitro ketones afforded relatively poor enantioselectivities (no more than 79% ee). Herein, we present this transformation in good yield with 68–96% ee using a thiourea catalyst derived from cinchona alkaloid. Notably up to 96% ee was obtained from the reaction of an aliphatic  $\alpha$ -nitro ketone and a  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester.

At the outset of our initial investigation, we focused on the reaction between methyl 2-oxo-4-phenylbut-3-enoate (1a) and 2-nitro-1-phenylethanone (2a). Interestingly, the <sup>1</sup>H NMR spectrum of the product showed that a Michael-type reaction occurred (Scheme 1). There were two groups of doublet peaks for two hydrogens of the C=C double bond conjugated with the aromatic ring at the range from 6.00 ppm to 7.00 ppm (J = 16.0 Hz) in the <sup>1</sup>H NMR spectrum of the aldol adduct. In contrast, only one group of doublet peaks near 6.8 ppm was found for the Michael-type product and the coupling constant had changed to 9.6 Hz. This information indicates that the conjugate system between the C=C double bond and the aromatic ring is broken.



Scheme 1  $^{1}$ H NMR spectra of products from 2-oxo-4-phenylbut-3-enoate.

Having confirmed the reaction type, we next carried out the screening of organocatalysts (Table 1). Catalyst I and II have

<sup>&</sup>lt;sup>a</sup>Department of Chemistry, Faculty of Science, Hong Kong Baptist University, Hong Kong, China. E-mail: ascchan@hkbu.edu.hk

<sup>&</sup>lt;sup>b</sup>State Key Laboratory for Chirosciences and Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hong Kong, China. E-mail: bcfyk@inet.polyu.edu.hk; Fax: (+852)-2364-9932; Tel: (+852)-3400-8682

<sup>†</sup> Electronic supplementary information (ESI) available. See DOI: 10.1039/c1ob06191g

#### Table 1 Screening of organocatalysts<sup>a</sup>



<sup>*a*</sup> Reaction conditions: **1a** (0.13 mmol), **2a** (0.1 mmol), catalyst (10 mol%) in CH<sub>2</sub>Cl<sub>2</sub> (0.3 mL) at 20 °C for 24 h. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> ee was determined by chiral HPLC.

been successfully employed in the asymmetric aldol reactions of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto esters with nitromethane<sup>4b</sup> and ketone,<sup>8</sup> respectively. However, unsatisfactory results were obtained from both I- and II-catalyzed reactions (Table 1, entries 1–3). When thiourea catalysts III and IV were used, remarkable enhancements of both yield and enantioselectivity were achieved. The best enantioselectivity was obtained with the aid of IIIb catalyst (77%, Table 1, entry 5). Particularly noteworthy was that this methodology could constitute an access to both enantiomers of **3aa**, taking advantage of the pseudoenantiomeric effect of thiourea catalyst derived from cinchona alkaloid.

Further optimization of reaction conditions was investigated and representative results are listed in Table 2. Commonly used organic solvents were examined and Et<sub>2</sub>O was found to be more suitable for this transformation (Table 2, entries 1–7). Systematic screening studies including the effect of mixed solvents, reaction temperature, the molar ratio of reactants, various added additives and catalyst loading revealed that **IIIb** enabled the formation of **3aa** in 56% yield with 92% ee at 20 °C in 12 h (Table 2, entry 8, see ESI for more details†).

Under the optimized reaction conditions, **IIIb**-catalyzed Michael-type reactions between a series of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto esters **1** and **2a** were surveyed (Table 3). In the presence of **IIIb**, Michael-type products **3aa–ra** were isolated in 47–74% yield with 72–94% ee. The ester functionality of the  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester had a large influence on the enantioselectivity: increasing the size of ester group resulted in lowering product ee (Table 3, entries 1–3). Various substituents, either electron-withdrawing (Table 3, entries 4–9, 12–15) or electron-donating groups (Table 3, entries 10–11, 16–17), could be introduced into the aromatic ring of  $\beta$ , $\gamma$ -

 Table 2
 Optimization of reaction conditions<sup>a</sup>

Ph CO <sub>2</sub> Me		111b <sup>0</sup> 2 (10 mol%		h
1a	2a		Ph <sup>wi</sup> 3aa CO <sub>2</sub> M	Иe
Solvent	$T/^{\circ}C$	t/h	Yield (%) <sup>b</sup>	ee (%) <sup>c</sup>
$CH_2Cl_2$	20	24	69	77
EtOAc	20	24	70	79
Toluene	20	24	62	85
THF	20	24	71	82
1,4-dioxane	20	24	79	80
t-BuOMe	20	24	44	93
$Et_2O$	20	24	48	93
$Et_2O$	20	12	56	94
	$\begin{array}{c} & & \\ & & \\ & & \\ & & \\ \hline \\ & \\ &$	$\begin{array}{c} \begin{array}{c} & & & \\ & & \\ & & \\ & & \\ \hline \\ h \\ & \\ h \\ h$	$\begin{array}{c cccc} & & & & & & \\ & & & & & \\ & & & & & \\ \hline & & & &$	$\begin{array}{c cccc} & & & & & & & & & & & & & & & & & $

<sup>*a*</sup> Unless noted, the following reaction conditions were used: **1a** (0.13 mmol), **2a** (0.1 mmol), **IIIb** (10 mol%) in solvent (0.3 mL) at temperature indicated for the time given in Table 2. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> ee was determined by chiral HPLC. <sup>*d*</sup> **IIIb** (15 mol%) and Et<sub>2</sub>O (0.5 mL) were used.

 Table 3
 Substrate scope<sup>a</sup>

-								
$R^{1} \xrightarrow{O}_{CO_{2}R^{2}}^{+} R^{3} \xrightarrow{O}_{2} XO_{2} \underbrace{(15 \text{ mol}\%)}_{Et_{2}O(0.5 \text{ mL})}^{O}_{2N} \xrightarrow{O}_{R^{3}}_{CO_{2}R^{2}} CO_{2}R^{2}$								
$\mathbf{R}^1$	$\mathbb{R}^2$	<b>R</b> <sup>3</sup>	Product	Yield (%) <sup>b</sup>	ee (%)			
Ph	Me	Ph	3aa	56	94			
Ph	Et	Ph	3ba	50	84			
Ph	<i>i</i> -Pr	Ph	3ca	61	72			
2-ClPh	Et	Ph	3da	50	82			
2-BrPh	Et	Ph	3ea	71	86			
3-ClPh	Et	Ph	3fa	62	84			
3-BrPh	Et	Ph	3ga	58	91			
3-NO <sub>2</sub> Ph	Me	Ph	3ha	56	86			
3-NO <sub>2</sub> Ph	Et	Ph	3ia	49	90			
3-MePh	Et	Ph	3ja	50	82			
3-MeOPh	Et	Ph	3ka	47	86			
4-FPh	Et	Ph	3la	56	88			
4-ClPh	Et	Ph	3ma	47	88			
4-BrPh	Et	Ph	3na	70	92			
4-NO <sub>2</sub> Ph	Et	Ph	3oa	58	84			
4-MePh	Et	Ph	3pa	56	82			
4-MeOPh	Et	Ph	3qa	63	73			
5-Me-2-thienyl	Et	Ph	3ra	94	68			
Ph	Me	<i>n</i> -Pr	3ab	51	96			
Ph	Et	<i>n</i> -Pr	3bb	50	83			
3-ClPh	Et	<i>n</i> -Pr	3fb	45	89			
3-BrPh	Et	<i>n</i> -Pr	3gb	43	90			
3-NO <sub>2</sub> Ph	Et	<i>n</i> -Pr	3ib	61	82			
3-MeOPh	Et	<i>n</i> -Pr	3kb	63	73			
4-FPh	Et	<i>n</i> -Pr	3lb	69	84			
4-ClPh	Et	<i>n</i> -Pr	3mb	55	88			
4-BrPh	Et	<i>n</i> -Pr	3nb	50	94			
4-MePh	Et	<i>n</i> -Pr	3pb	54	78			
	$R^1$ $R^1$	$\begin{array}{c} & & & \\ & & \\ R^1 & & & \\ & & & \\ & & & \\ & & & \\ R^1 & & & \\ & & & \\ & & & \\ R^1 & & & \\ & & & \\ R^2 & & \\ \end{array}$ $\begin{array}{c} & & & \\ & & & \\ R^3 & & & \\ & & & \\ R^3 & & & \\ & & & \\ R^3 & $	$\begin{array}{c c} & & & & \\ & & & \\ R^1 & & & R^2 & R^3 \\ \hline \\ & & & & \\ R^1 & & R^2 & R^3 \\ \hline \\ R^1 & & & R^3 & R^3 \\ \hline \\ R^1 & & & R^3 \\ \hline \\ R^1 & & & R^3 \\$	Ib       Ib $O_2N$ $R^1                                     $	IIIb $R^1 + \begin{pmatrix} 0 \\ 1 \end{pmatrix} \\ R^1 \end{pmatrix} \\ R^1 \end{pmatrix} \\ R^2$ R R R R R R R R R R R R R R R R R R R			

<sup>*a*</sup> Unless noted, the following reaction conditions were used: 1 (0.13 mmol), **2** (0.1 mmol), **IIIb** (15 mol%) in  $Et_2O$  (0.5 mL) at 20 °C for 12 h. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> ee was determined by chiral HPLC.

unsaturated  $\alpha$ -keto ester. No significant electronic effect on the aromatic moiety was observed. A heteroaromatic  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester afforded **3ra** in up to 94% yield with 68% ee (Table 3, entry 18).

The reaction of challenging aliphatic  $\alpha$ -nitro ketone with **1** was also probed. 1-Nitropentan-2-one (**2b**) reacted with **1a** to afford

**3ab** in moderate yield with up to 96% ee (Table 3, entry 19). Product **3bb** was formed in similar yield with 83% ee from ethyl 2-oxo-4phenylbut-3-enoate (Table 3, entry 20). The substituted aromatic  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester could also be employed to afford **3ab– 3pb** in medium yields with good to excellent enantioselectivities (Table 3, entries 21–28). It should be noted that more than 90% ee were obtained not only from the aromatic  $\alpha$ -nitro ketone but also from the aliphatic  $\alpha$ -nitro ketone.

The absolute configuration of the products was R, which was determined by comparison of optical rotations with those in the literature.<sup>9</sup>

In conclusion, we have developed a new Michael-type reaction of  $\beta$ , $\gamma$ -unsaturated  $\alpha$ -keto ester with both aromatic and aliphatic  $\alpha$ -nitro ketone to afford products in 47–94% yields with 68–96% ee. Compared with Wang's and Yan's reports,<sup>9</sup> higher ee values were obtained from aliphatic  $\alpha$ -nitro ketone. It should be noted that this methodology offered an enantioselective pathway for the synthesis of chiral 5-nitro-pent-2-enoates, a precursor to chiral  $\alpha$ ketolactam. Development of new type organocatalysts and further applications of these reactions are currently under investigation.

## Acknowledgements

We gratefully thank the European Commission FP7-201431 (CATAFLU.OR) and PolyU Internal Grant DA (A-PD0X) for financial support.

## Notes and references

- For selected books on organocatalysis, see: (a) E. N. Jacobsen, A. Pfaltz and H. Yamamoto, *Comprehensive Asymmetric Catalysis*, Springer, 1999; (b) A. Berkessel and H. Gröger, *Asymmetric Organocatalysis*, Wiley, Weinheim, 2005; (c) P. I. Dalko, *Enantioselective Organocatalysis*, Wiley, Weinheim, 2007; (d) M. T. Reetz, B. List, S. Jaroch and H. Weinmann, *Organocatalysis*, Springer, 2007; (e) B. List, *Asymmetric Organocatalysis*, Springer, 2009.
- 2 For selected reviews on organocatalytic asymmetric Michael addition, see: (a) O. M. Berner, L. Tedeschi and D. Enders, *Eur. J. Org. Chem.*, 2002, 1877; (b) R. Ballini, G. Bosica, D. Fiorini, A. Palmieri and M. Petrini, *Chem. Rev.*, 2005, **105**, 933; (c) D. Almaşi, D. A. Alonso and C. Nájera, *Tetrahedron: Asymmetry*, 2007, **18**, 299; (d) S. B. Tsogoeva, *Eur. J. Org. Chem.*, 2007, 1701.
- 3 For selected references, see: (a) T. Bui and C. F. Barbas III, Tetrahedron Lett., 2000, 41, 6951; (b) Y. Yamamoto, N. Momiyama and H. Yamamoto, J. Am. Chem. Soc., 2004, 126, 5962; (c) J. W. Yang, M. T. Hechavarria Fonseca and B. List, J. Am. Chem. Soc., 2005, 127, 15036; (d) M. Marigo, T. Schulte, J. Franzen and K. A. Jørgensen, J. Am. Chem. Soc., 2005, 127, 15710; (e) D. Enders, M. R. M. Huttl, C. Grondal and G. Raabe, Nature, 2006, 441, 861; (f) Y. Wang, X. Liu and L. Deng, J. Am. Chem. Soc., 2006, 128, 3928; (g) H. Sundén, I. Ibrahem, G.-L. Zhao, L. Eriksson and A. Córdova, Chem.-Eur. J., 2007, 13, 574; (h) P. Galzerano, F. Pesciaioli, A. Mazzanti, G. Bartoli and P. Melchiorre, Angew. Chem., Int. Ed., 2009, 48, 7892; (i) L. Albrecht, H. Jiang, G. Dickmeiss, B. Gschwend, S. G. Hansen and K. A. Jørgensen, J. Am. Chem. Soc., 2010, 132, 9188; (j) R. Imashiro, H. Uehara and C. F. Barbas III, Org. Lett., 2010, 12, 5250; (k) H. Uehara, R. Imashiro, G. Hernández-Torres and C. F. Barbas III, Proc. Natl. Acad. Sci. U. S. A., 2010, 107, 20672; (1) B. Tan, N. R. Candeias and C. F. Barbas III, Nat. Chem., 2011, 3, 473.
- 4 For some references, see: (a) C. Christensen, K. Juhl, R. G. Hazell and K. A. Jørgensen, J. Org. Chem., 2002, 67, 4875; (b) H. Li, B. Wang and L. Deng, J. Am. Chem. Soc., 2006, 128, 732; (c) C.-L. Cao, X.-L. Sun,

Y.-B. Kang and Y. Tang, *Org. Lett.*, 2007, **9**, 4151; (*d*) C. Zheng, Y. Wu, X. Wang and G. Zhao, *Adv. Synth. Catal.*, 2008, **350**, 2690; (*e*) G. Blay, V. Hernández-Olmos and J. R. Pedro, *Org. Biomol. Chem.*, 2008, **6**, 468.

- 5 For selected references, see: (a) J. Thorhauge, M. Johannsen and K. A. Jørgensen, Angew. Chem., Int. Ed., 1998, 37, 2404; (b) D. A. Evans, E. J. Olhava, J. S. Johnson and J. M. Janey, Angew. Chem., Int. Ed., 1998, 37, 3372; (c) H. Audrain, J. Thorhauge, R. G. Hazell and K. A. Jørgensen, J. Org. Chem., 2000, 65, 4487; (d) D. A. Evans, J. S. Johnson and E. J. Olhava, J. Am. Chem. Soc., 2000, 122, 1635; (e) R. A. Stavenger and S. L. Schreiber, Angew. Chem., Int. Ed., 2001, 40, 3417; (f) K. Juhl and K. A. Jørgensen, Angew. Chem., Int. Ed., 2003, 42, 1498; (g) F. Gohier, K. Bouhadjera, D. Faye, C. Gaulon, V. Maisonneuve, G. Dujardin and R. Dhal, Org. Lett., 2007, 9, 211; (h) S. Samanta, J. Krause, T. Mandal and C.-G. Zhao, Org. Lett., 2007, 9, 2745; (i) M. He, B. J. Beahm and J. W. Bode, Org. Lett., 2008, 10, 3817; (j) F. Gallier, H. Hussain, A. Martel, A. Kirschning and G. Dujardin, Org. Lett., 2009, 11, 3060; (k) W. Yao, L. Pan, Y. Wu and C. Ma, Org. Lett., 2010, 12, 2422; (l) D. Xu, Y. Zhang and D. Ma, Tetrahedron Lett., 2010, 51, 3827; (m) Y. Ying, Z. Chai, H.-F. Wang, P. Li, C.-W. Zheng, G. Zhao and Y.-P. Cai, Tetrahedron, 2011, 67, 3337; (n) M. Terada and H. Nii, Chem.-Eur. J., 2011, 17, 1760.
- 6 (a) K. B. Jensen, J. Thorhauge, R. G. Hazell and K. A. Jørgensen, Angew. Chem., Int. Ed., 2001, 40, 160; (b) K. A. Jørgensen, Synthesis, 2003, 1117; (c) R. P. Herrera, D. Monge, E. Martín-Zamora, R. Fernández and J. M. Lassaletta, Org. Lett., 2007, 9, 3303; (d) X. Xiong, C. Ovens, A. W. Pilling, J. W. Ward and D. J. Dixon, Org. Lett., 2008, 10, 565; (e) Y. Liu, D. Shang, X. Zhou, Y. Zhu, L. Lin, X. Liu and X. Feng, Org. Lett., 2010, 12, 180.
- 7 (a) N. Halland, T. Velgaard and K. A. Jørgensen, J. Org. Chem., 2003, 68, 5067; (b) H. L. Van Lingen, W. Zhuang, T. Hansen, F. P. J. T. Rutjes and K. A. Jørgensen, Org. Biomol. Chem., 2003, 1, 1953; (c) C.-L. Cao, X.-L. Sun, Y.-B. Kang and Y. Tang, Org. Lett., 2007, 9, 4151; (d) Q.-G. Wang, X.-M. Deng, B.-H. Zhu, L.-W. Ye, X.-L. Sun, C.-Y. Li, C.-Y. Zhu, Q. Shen and Y. Tang, J. Am. Chem. Soc., 2008, 130, 5408; (e) L.-W. Ye, X. Han, X.-L. Sun and Y. Tang, Tetrahedron, 2008, 64, 8149; (f) M. A. Calter and J. Wang, Org. Lett., 2009, 11, 2205; (g) Y. Gao, Q. Ren, L. Wang and J. Wang, Chem.-Eur. J., 2010, 16, 13068; (h) Y. Gao, Q. Ren, S.-M. Ang and J. Wang, Org. Biomol. Chem., 2011, 9, 3691; (i) Z. Dong, J. Feng, X. Fu, X. Liu, L. Lin and X. Feng, Chem.-Eur. J., 2011, 17, 1118.
- 8 (a) P. Li, J. Zhao, F. Li, A. S. C. Chan and F. Y. Kwong, Org. Lett., 2010, 12, 5616; (b) P. Li, S. H. Chan, A. S. C. Chan and F. Y. Kwong, Adv. Synth. Catal., 2011, 353, 1179.
- 9 During the completion of our experimental works, two related, yet different substrate scope papers appeared, see: (a) Y. Gao, Q. Ren, W.-Y. Siau and J. Wang, *Chem. Commun.*, 2011, 47, 5819; (b) R. Lu, Y. Yan, J. Wang, Q. Du, S. Nie and M. Yan, *J. Org. Chem.*, 2011, **76**, 6230.
- 10 For selected references of thiourea or bifunctional Brønsted-base thiourea catalysis, see: (a) D. P. Curran and L. H. Kuo, J. Org. Chem., 1994, 59, 3259; (b) M. S. Sigman and E. N. Jacobsen, J. Am. Chem. Soc., 1998, 120, 4901; (c) T. Okino, Y. Hoashi and Y. Takemoto, J. Am. Chem. Soc., 2003, 125, 12672; (d) A. Berkessel, F. Cleemann, S. Mukherjee, T. N. Müller and J. Lex, Angew. Chem., Int. Ed., 2005, 44, 807; (e) R. P. Herrera, V. Sgarzani, L. Bernardi and A. Ricci, Angew. Chem., Int. Ed., 2005, 44, 6576; (f) J. Ye, D. J. Dixon and P. S. Hynes, Chem. Commun., 2005, 4481; (g) D. R. Li, A. Murugan and J. R. Falck, J. Am. Chem. Soc., 2008, 130, 46; (h) G. Dickmeiss, V. D. Sio, J. Udmark, T. B. Poulsen, V. Marcos and K. A. Jørgensen, Angew. Chem., Int. Ed., 2009, 48, 6650; (i) T. Bui, S. Syed and C. F. Barbas III, J. Am. Chem. Soc., 2009, 131, 8758; (j) C. R. Jones, G. D. Pantos, A. J. Morrison and M. D. Smith, Angew. Chem., Int. Ed., 2009, 48, 7391; (k) S. J. Zuend and E. N. Jacobsen, J. Am. Chem. Soc., 2009, 131, 15358; (1) W. J. Nodes, D. R. Nutt, A. M. Chippindale and A. J. A. Cobb, J. Am. Chem. Soc., 2009, 131, 16016; (m) S. J. Zuend, P. C. Matthew, P. L. Mathieu and E. N. Jacobsen, Nature, 2009, 461, 968; (n) A. Peschiulli, B. Procuranti, C. J. O'Connor and S. J. Connon, Nat. Chem., 2010, 2, 380; (o) Y. Gao, Q. Ren, H. Wu, M. Li and J. Wang, Chem. Commun., 2010, 46, 9232; (p) Y. Gao, Q. Ren, L. Wang and J. Wang, Chem.-Eur. J., 2010, 16, 13068; (q) Q. Ren, Y. Gao and J. Wang, Chem.-Eur. J., 2010, 16, 13594.