Varied Reactivity of the Germene Mes₂Ge=CR₂ toward **Nitriles**

Sakina Ech-Cherif El Kettani,[†] Mohamed Lazraq,[†] Henri Ranaivonjatovo,^{*,‡} Jean Escudié,[‡] Claude Couret,[‡] Heinz Gornitzka,[‡] and Nathalie Merceron[‡]

Laboratoire de Chimie Organique, BP 1796, Faculté des Sciences Dhar El Mehraz, Université Sidi Mohamed Ben Abdellah, Fes-Atlas, Fes, Morocco, and Hétérochimie Fondamentale et Appliquée, UMR 5069, Université Paul Sabatier, 118 Route de Narbonne, 31062, Toulouse Cedex 04, France

Received July 16, 2004

The germene $Mes_2Ge=CR_2$ (1, CR_2 = fluorenylidene) shows varied behavior toward nitriles. It reacts as a 1,2-dipole with t-BuCN giving a four-membered ring heterocycle, 2-aza-3germacyclobut-1-ene, as a 1,4-dipole with PhCN, leading to a six-membered ring heterocycle, 3,4-dihydro-3-germaisoquinoline, and as a base, abstracting the α -H of R'CH₂CN to give α -cyanogermanes (R' = H, 2-thienyl, 2-FC₆H₄, 4-FC₆H₄).

Introduction

The synthesis of the first stable compound containing a Si=C double bond, the silene (Me₃Si)₂Si=C(OSiMe₃)-Ad,¹ was a milestone in organometallic chemistry. Other silenes and some germenes >Ge=C< and stannenes >Sn=C< have been prepared since this date.² Despite the great steric hindrance necessary for their stabilization as monomers, a flourishing chemistry of these organometallic alkenes has been developed. Their reactions with various unsaturated compounds showed them to be, among others, valuable sources of metalated heterocycles difficult to obtain by other routes. To the best of our knowledge, in the case of nitriles, only the [2+4] cycloadduct of the silene Me₂Si=C(SiMe₃)₂ with PhC=N has been reported so far,^{3a} while Me₂Si=CH₂, generated by pyrolysis of the corresponding silacyclobutane, inserted into a CH bond of acetonitrile.^{3b} Silenes $(Me_3Si)_2Si=C(OSiMe_3)R$ gave a [2+2] cycloadduct with the C=C bond of acrylonitrile.^{3c} In the case of germenes, only one recent paper reported the reactions of an acetylene-bridged bis(germene) with dinitrile compounds.⁴

(1) Brook, A. G.; Abdekassen, F.; Gutekunst, B.; Gutekunst, G.; Kallury, R. K. J. Chem. Soc., Chem. Commun. 1981, 191-192.

(2) For reviews, see: (a) Raabe, G.; Michl, J. The Chemistry of Organic Silicon Compounds; Patai, S., Rappoport, Z., Eds.; J. Wiley and Sons: New York, 1989, Chapter 17, pp 1015–1141. (b) Barrau, J.; Escudié, J.; Satgé, J. *Chem. Rev.* **1990**, *90*, 283–319. (c) Brook, A. G.; Brook, M. A. Adv. Organomet. Chem. **1996**, *39*, 71–158. (d) Baines, K. M.; Stibbs, W. G. Adv. Organomet. Chem. **1996**, *39*, 275–324. (e) K. M.; Stubbs, W. G. Adv. Organomet. Chem. 1990, 39, 273–224. (e) Müller, T.; Ziehe, W.; Auner, N. The Chemistry of Organic Silicon Compounds; Patai, S., Rappoport, Z., Eds.; Wiley: New York, 1998; Vol. 2, Part 2, Chapter 16, pp 857–1063. (f) Escudié, J.; Couret, C.; Ranaivonjatovo, H. Coord. Chem. Rev. 1998, 178–180, 565–592. (g)

Katavonjatovo, H. Coord. Chem. Rev. 1996, 178–180, 505–532. (g)
Escudié, J.; Ranaivonjatovo, H. Adv. Organomet. Chem. 1999, 44, 113–174. (h) Weidenbruch, M. J. Organomet. Chem. 2002, 646, 39–52. (3) (a) Wiberg, N.; Preiner, G.; Schieda, O. Chem. Ber. 1981, 114, 3518–3532. (b) Bush, R. D.; Golino, C. M.; Roark, D. N.; Sommer, L. H. J. Organomet. Chem. 1973, 59, C17–C20. (c) Brook, A. G.; Hu, S. S.; Saxena, A. K.; Lough, A. J. Organometallics 1991, 10, 2758–2767.
 (4) Meiners, F.; Haase, D.; Koch, R.; Saak, W.; Weidenbruch, M. Organometallics 2002, 21, 3990–3995.

Thus, we became interested in investigating the reactivity of nitriles with the germene $Mes_2Ge=CR_2$ (1)⁵ (Mes = 2,4,6-trimethylphenyl, CR_2 = fluorenylidene), in which the Ge=C unsaturation is conjugated with the fluorenylidene unit. Interestingly, three kinds of products were obtained in this study, depending on the nitrile used.

Results and Discussion

Germene 1 reacted at 140 °C as a 1,2-dipole with trimethylacetonitrile, giving 2-aza-3-germacyclobut-1ene (2), a four-membered ring derivative with both a heavy group 14 element and a C=N double bond⁶ (eq 1).



The observed regioselectivity (N bonded to germanium) in this [2+2] cycloaddition between the Ge=C and C=N unsaturation was expected from the $Ge^{\delta +}=C^{\delta -}$ polarity of the germanium-carbon double bond. The structure of 2 was evidenced by mass spectroscopy (besides the molecular peak, the main fragment was the starting germene obtained by a $[4 \rightarrow 2+2]$ cycloreversion) and particularly by NMR spectroscopic analysis

^{*} To whom correspondence should be addressed. E-mail: ranaivon@ chimie.ups-tlse.fr.

Université Sidi Mohamed Ben Abdellah.

[‡] Université Paul Sabatier.

^{(5) (}a) Couret, C.; Escudié, J.; Satgé, J.; Lazraq, M. J. Am. Chem. Soc. 1987, 109, 4411-4412. (b) Lazraq, M.; Escudié, J.; Couret, C.; Satgé, J.; Dräger, M.; Dammel, R. Angew. Chem., Int. Ed. Engl. 1988, 27, 828-829.

⁽⁶⁾ Rather similar 1-aza-2,3-disilacyclobut-1-enes have been obtained by thermolysis or photolysis of cyclotrisilanes (R₂Si)₃ in the presence of nitriles. Although in a formal sense they are the product of a [2+2] cycloaddition reaction between the disilene $R_2Si=SiR_2$ and a nitrile, their formation is likely to proceed via initially generated azasilacyclopropenes followed by ring expansion with a second silylene $R_2Si:$ (a) Weidenbruch, M.; Meiners, F.; Saak, W. *Can. J. Chem.* **2000**, 78, 1469–1473. (b) Belzner, J.; Ihmels, H.; Noltemeyer, M. Tetrahedron Lett. 1995, 36, 8187-8190.



(δ ¹³C (C=N) = 198.41 ppm). In the ¹H NMR spectrum, a very broad singlet, characteristic of a cyclic structure,⁷ was observed for the *ortho*-methyl groups of equivalent mesityl groups due to their slow rotation caused by the large steric hindrance, which confers to **2** a surprisingly high thermal stability.

The structure of **2** was confirmed chemically by its hydrolysis, which gave a nearly quantitative yield of tetramesityldigermoxane (**3**)⁸ and fluorenyl(*tert*-butyl)-ketone (**4**).⁹ Their formation resulted from the initial cleavages of the Ge–C and Ge–N bonds, giving first the dihydroxygermane **5**^{8a,10} (which slowly converts to **3**) and the transient imine **6** (δ ¹³C NMR (C=N) = 189.1 ppm), which undergoes further hydrolysis to **4** (Scheme 1).

The behavior of **1** toward benzonitrile is completely different, giving compound **7** (eq 2). The structure of **7** was established by an X-ray diffraction (Figure 1).



The first step of the reaction probably was the formation of **8** by [2+4] cycloaddition between the C=N moiety and the Ge=C-C=C dienic system of **1**, followed by rearomatization via a 1,3-sigmatropic shift of H, leading to 3,4-dihydro-3-germaisoquinoline (7). This a priori surprising diene behavior of **1** can, however, be understood both from its cisoid structure and from the conjugation of the Ge=C double bond with the fluorenylidene unit.¹¹ Usually, one of the features of Diels-Alder reactions with nitriles as dienophiles is the extremely high reaction temperature required, often above 200 °C.¹² Therefore, the formation of **7** in 78% yield at room temperature was rather unexpected and confirmed the high reactivity of **1**.

(9) ¹H NMR data were in accordance with those found in: Meyers, C. Y.; Wahner, A. P.; Manohar, S. K.; Carr, S. E.; Chan-Yu-King, R.; Robinson, P. D. Acta Crystallogr. **1991**, C47, 1236–1239.



Figure 1. Molecular structure of **7** with thermal ellipsoids drawn at the 50% probability level. Hydrogen atoms are omitted for clarity. Selected bond lengths (Å) and angles (deg): Ge1–N1, 1.855 (4); C1–N1, 1.306(6); Ge1–C30, 1.979(4); Ge1–C21, 1.964(5); Ge1–C14, 1.990(5); C13–C14, 1.516(6); C8–C13, 1.395(6); C1–C8, 1.496(7); N1–Ge1–C14, 98.74(19); N1–Ge1–C21, 101.16(19); N1–Ge1–C30, 111.19(17); C21–Ge1–C30, 112.6(2); C1–N1–Ge1, 122.8(3); N1–C1–C8, 124.5(4).

In contrast, with nitriles of type R'CH₂C=N with R' groups of various electronic effects, **1** preferentially reacted at the hydrogen α to the C=N bond. These reactions occurred at room temperature in Et₂O solutions to afford α -cyanogermanes **9a**-**d** in quantitative yield, regioselectively with exclusive formation of the GeCH(R')C=N moiety as expected from the Ge^{δ +}=C^{δ -} polarity (eq 3). Thus germene **1** contains a strongly basic fluorenylidene unit.



Such electrophilic addition of acidic HA compounds to the Ge=C double bond of **1** has previously been reported to occur only with the more "acidic" hydrogens of alcohols, water, thiols, and imines containing an NH bond to give $Mes_2Ge(A)C(H)R_2$ adducts.^{5a,13}

Experimental Section

General Procedures. All manipulations were carried out under N_2 or Ar using standard Schlenk techniques with solvents freshly distilled from sodium benzophenone. ¹H and ¹³C NMR spectra were recorded in CDCl₃ on a Bruker AC 250 instrument at 250.1 MHz and a Bruker Avance 300 instrument at 75.48 MHz, respectively. Mass spectra were measured on a Hewlett-Packard 5989 A spectrometer by EI at 70 eV. Melting points were determined on a Leitz microscope heating stage 250. IR were recorded on a Perkin-Elmer 1600 FT-IR spectrometer. Elemental analyses were performed by the "Service de Microanalyse de l'Ecole de Chimie de Toulouse".

The carbon atoms of the fluorenyl group are numbered as shown in Figure 1.

Synthesis of 2. Germene **1** was prepared as previously described⁵ by addition of 1 molar equiv of *tert*-butyllithium (1.7

⁽⁷⁾ In the four-membered ring 1-oxa-2-germacyclobutane, obtained from **1** and benzophenone, a similar very broad signal was observed for the *ortho*-methyl groups of the mesityl substituents: Lazraq, M.; Couret, C.; Escudié, J.; Satgé, J.; Dräger, M. *Organometallics* **1991**, *10*, 1771–1778.

^{(8) (}a) Duverneuil, G.; Mazerolles, P.; Perrier, E. *Appl. Organomet. Chem.* **1994**, *8*, 119–127. (b) Samuel, M. S.; Jennings, M. C.; Baines, K. M. J. Organomet. Chem. **2001**, *636*, 130–137.

⁽¹⁰⁾ Slow decomposition of **5** to **3** was observed previously: (a) El Baz, F.; Rivière-Baudet, M.; Chazalette, C.; Gornitzka, H. *Phosphorus, Sulfur Silicon* **2000**, *163*, 121–142. (b) Rivière-Baudet, M.; El Baz, F.; Satgé, J.; Khallaayoun, A.; Ahra, M. *Phosphorus, Sulfur Silicon* **1996**, *112*, 203–217.

⁽¹¹⁾ The unique precedent diene behavior of **1** was postulated in its reaction with *t*-BuC≡P at 170 °C: Lazraq, M.; Escudié, J.; Couret, C.; Bergsträsser, U.; Regitz, M. *Chem. Commun.* **1993**, 569–570.

⁽¹²⁾ Hamer, J. 1.4-Cycloaddition Reactions, Academic Press: New York, 1967; pp 97-125.

⁽¹³⁾ Lazraq, M.; Escudié, J.; Couret, C.; Satgé, J.; Soufiaoui, M. Organometallics **1991**, *10*, 1140–1143.

M in pentane) to a solution of the fluorogermane Mes₂Ge(F)-CHR₂ (1.00 g, 2.02 mmol) in Et₂O (20 mL) cooled at -78 °C. Warming to room temperature afforded an orange solution of 1, which is nearly quantitatively produced along with a precipitate of LiF. Crude solutions of 1 were used without further purification. The mixture of Mes₂Ge=CR₂ and t-BuCN (0.17 g, 2.02 mmol) was heated in a sealed tube at 140 °C for 15 h. After filtration to remove LiF and evaporation of solvents in vacuo, crystallization from pentane at -20 °C afforded 0.54 g of 2: white crystals; mp 228 °C; yield 48%; ¹H NMR (CDCl₃) δ 0.83 (s, 9H, t-Bu), 2.01 (br s, 12H, o-Me), 2.24 (s, 6H, p-Me), 6.75 (s, 4H, arom CH of Mes), [7.01 (d, ${}^{3}J_{HH} = 7.6$ Hz, 2H), 7.10 and 7.33 (2t, ${}^{3}J_{\rm HH}$ = 7.6 Hz, 2 × 2H), 7.85 (d, ${}^{3}J_{\rm HH}$ = 7.6 Hz, 2H), H of CR₂]; ¹³C NMR (CDCl₃) δ 20.98 (*p*-Me), 23.20 (o-Me), 28.68 (C(CH₃)₃), 42.79 (C(CH₃)₃), 73.77 (CR₂), 120.03, 123.80, 125.89 and 126.42 (CH of CR2), 126.70 (m-CH of Mes), 132.90 (ipso-C of Mes), 138.30, 139.50, 143.39, and 144.38 (oand p-C of Mes, quat. C of CR₂), 198.41 (C=N); IR (KBr) 1542.0 cm⁻¹ (ν C=N); MS, m/z (%) 559 (M, 4), 502 (M - t-Bu, 2), 476 (Mes₂Ge=CR₂, 42), 440 (M - Mes, 2), 311 (Mes₂Ge - 1, 100), 192 (MesGe - 1, 38). Anal. Calcd for C₃₆H₃₉GeN: C, 77.45; H, 7.04. Found: C, 77.21; H, 7.25.

Hydrolysis of 2. Compound **2** slowly hydrolyzed when left in wet CDCl₃ solution at room temperature. After 3 days, the NMR signals of **2** disappeared and new signals assigned to dihydroxygermane **5** and imine **6** were observed in the ¹H and ¹³C NMR spectra.

5: ¹H NMR (CDCl₃) δ 1.84 (s, 2H, OH), 2.28 (s, 6H, *p*-Me), 2.45 (s, 12H, *o*-Me), 6.83 (s, 4H, arom. H); ¹³C NMR (CDCl₃) δ 21.12 (*p*-Me), 23.14 (*o*-Me), 129.32 (*m*-C), 133.20 (*ipso*-C), 140.08 (*p*-C), 143.10 (*o*-C).

6: ¹H NMR (CDCl₃) δ 1.49 (s, 9H, *t*-Bu), 5.09 (s, 1H, C*H*R₂), [7.26 (t, ³*J*_{HH} = 7.6 Hz, 2H), 7.35 (d, ³*J*_{HH} = 7.6 Hz, 2H), 7.38 (t, ³*J*_{HH} = 7.6 Hz, 2H), 7.78 (d, ³*J*_{HH} = 7.6 Hz, 2H), CR₂]; ¹³C NMR (CDCl₃) δ 27.98 (C(*C*H₃)₃), 40.78 (*C*(CH₃)₃), 52.78 (*C*R₂), 120.08, 124.34, 127.53 and 127.71 (CH of CR₂), 140.10 and 142.00 (quat. C of CR₂), 189.08 (C=N).

 ${\bf 5}$ and ${\bf 6}$ were not isolated. After 2 weeks, NMR analysis of this solution showed the nearly quantitative formation of ${\bf 3}^8$ and ${\bf 4}.$

3: ¹³C NMR (CDCl₃) δ 21.12 (*p*-Me), 23.14 (*o*-Me), 129.32 (*m*-C of Mes), 133.20 (*ipso*-C), 140.08 (*p*-C of Mes), 143.10 (*o*-C of Mes).

4 was also identified by an independent synthesis.⁹ 13 C NMR (CDCl₃) δ 26.57 (CMe₃), 45.64 (CMe₃), 57.11 (HCR₂), 120.36, 124.43, 127.32 and 127.81 (CH of CR₂), 142.38 and 143.60 (quat. C of CR₂), 213.18 (CO); MS (EI, m/z, %) 250 (M, 10), 192 (M - *t*-Bu - 1, 3), 165 (R₂CH, 30), 57 (*t*-Bu, 100).

Synthesis of 7. One equivalent of PhCN (0.21 g, 2.02 mmol) was added to a solution of 1, prepared as previously described,⁵ cooled at -78 °C. The reaction mixture gradually turned from orange to light yellow after overnight stirring at room temperature. LiF was then filtered, and the solvents were removed in vacuo. Recrystallization of the crude material from pentane at -20 °C afforded white crystals of 7 (0.91 g, yield 78%, mp 200–201 °C): ¹H NMR (C₆D₆) δ 1.85 (s, 3H, *p*-Me), 1.97 (br s, 6H, o-Me), 2.15 (s, 3H, p-Me), 2.52 (s, 6H, o-Me), 4.55 (s, 1H, CH–Ge), 6.43 and 6.81 (2s, 2 \times 2H, arom H of Mes), 7.10– 7.29 (m, 7H, m- and p-H of C₆H₅ and 4H of CR₂), 7.51, 7.71, and 7.81 (3d, ${}^{3}J_{\rm HH} =$ 7.6 Hz, 3 \times 1H, 3H of CR₂), 7.88–7.92 (m, 2H, o-H of C₆H₅); ¹³C NMR (C₆D₆) δ 20.42 and 20.75 (p-Me), 22.80 and 24.68 (o-Me), 41.25 (CHR₂), 120.43, 121.20, 124.44, 125.95, 126.37, 126.90, 127.36 (CH of CR₂), 127.68, 128.00, 128.52, and 129.08 (m-CH of Mes, o- and m-CH of Ph), 129.16 (p-CH of Ph), 129.81, 135.47, 136.05, 138.55, 138.75, 141.15, 141.20, 141.96, 142.25, 143.25, 143.99, and 145.47 (ipso-C of Ph, ipso-, o-, and p-C of Mes, C_{12,13,15,16}), 171.35 (C=N); IR (KBr) 1556 cm⁻¹ (ν C=N); MS, m/z (%) 579 (M, 9), 564 (M - Me, 3), 460 (M - Mes, 3), 341 (M - 2Mes, 10), 311 (Mes₂Ge - 1, 100), 267 (M - Mes₂Ge, 18), 190 (M - Mes₂Ge

- Ph, 17). Anal. Calcd for $C_{38}H_{35}GeN:$ C, 78.93; H, 6.10. Found: C, 78.75; H, 6.05.

Crystal data for 7: $C_{38}H_{35}$ GeN, M = 578.26, monoclinic, $P2_1/c$, a = 11.701(3) Å, b = 15.722(4) Å, c = 16.171(4) Å, $\beta = 91.083(4)^\circ$, V = 2974.3(12) Å³, Z = 4, T = 193(2) K. 11 756 reflections (4208 independent, $R_{int} = 0.1054$) were collected at low temperatures using an oil-coated shock-cooled crystal on a Bruker-AXS CCD 1000 diffractometer with Mo K α radiation ($\lambda = 0.71073$ Å). The structure was solved by direct methods (SHELXS-97),¹⁴ and all non hydrogen atoms were refined anisotropically using the least-squares method on $F^{2,15}$. Largest electron density residue: 0.683 e Å⁻³, R_1 (for $I > 2\sigma$ -(I)) = 0.0488 and $wR_2 = 0.0972$ (all data) with $R_1 = \Sigma ||F_0| - |F_c||/\Sigma|F_0|$ and $wR_2 = (\Sigma w(F_0^2 - F_c^2)^2/\Sigma w(F_0^2)^2)^{0.5}$.

Reaction of Mes₂Ge=CR₂ with Nitriles R'CH₂C=N: Formation of α -Cyanogermanes 9. To a crude solution of 1 prepared as previously described⁵ from 1.00 g (2.02 mmol) of Mes₂Ge(F)CHR₂ in Et₂O (20 mL) cooled at -78 °C was added 1 equiv of the nitrile R'CH₂-C=N. The reaction mixture was allowed to warm to room temperature, during which time it slowly turned from orange to pale yellow. After overnight stirring, solvent was removed in vacuo and replaced by 15 mL of pentane. Crystallization at -20 °C afforded pure **9a**-**d** as crystalline compounds.

9a: white crystals; 0.96 g, yield 92%; mp 142–143 °C; ¹H NMR (CDCl₃) δ 1.64 (s, 3H, CH₂), 2.11 (s, 12H, *o*-Me), 2.27 (s, 6H, *p*-Me), 5.10 (s, 1H, *CH*R₂), 6.81 (s, 4H, arom H of Mes), [7.15 (d, ³*J*_{HH} = 7.6 Hz, 2H), 7.18 and 7.35 (2t, ³*J*_{HH} = 7.6 Hz, 2 × 2H), 7.83 (d, ³*J*_{HH} = 7.6 Hz, 2H), CR₂]; ¹³C NMR (CDCl₃) δ 6.19 (CH₂), 20.92 (*p*-Me), 24.20 (*o*-Me), 43.81 (*C*HR₂), 119.43 (C=N), 119.76, 125.12, 126.38, and 126.68 (CH of CR₂), 129.69 (*m*-CH of Mes), 133.35, 139.13, 140.74, 142.64, and 143.55 (*ipso*-, *o* and *p*-C of Mes and quat. C of CR₂); IR (KBr) 2232 cm⁻¹ (*v*C=N); MS, *m*/*z* (%) 517 (M, 1), 477 (M – CH₂CN, 3), 352 (M – CHR₂, 85), 311 (Mes₂Ge – 1, 15), 192 (MesGe – 1, 20), 165 (CHR₂, 100). Anal. Calcd for C₃₃H₃₃GeN: C, 76.78; H, 6.44. Found: C, 77.02; H, 6.55.

9b: white crystals; 1.01 g, yield 82%; mp 145 °C; ¹H NMR (CDCl₃) δ 1.69 and 2.03 (2s, 2 × 6H, *o*-Me), 2.10 and 2.24 (2s, 2 × 3H, p-Me), 4.96 (s, 1H, CH-C=N), 5.11 (s, 1H, CHR₂), 6.39 and 6.68 (2s, 2 × 2H, arom H of Mes), 6.69-8.15 (m, 12H, arom H of C₆H₄F and CR₂); ¹³C NMR (CDCl₃) δ 20.71 and 20.86 (*p*-Me), 23.57 (d, ${}^{3}J_{CF}$ =1.1 Hz, *C*HC=N), 24.38 and 24.90 (*o*-Me), 46.67 (d, ${}^{5}J_{CF} = 1.5$ Hz, CHR₂), 114.90 (d, ${}^{2}J_{CF} = 21.9$ Hz, CH–CF), 119.75 and 119.76 (CH of CR₂), 121.96 (d, ${}^{2}J_{\rm CF}$ = 15.1 Hz, *ipso*-C of C₆H₄F), 122.26 (C=N), 124.27 (d, $J_{CF} = 3.3$ Hz, CH of C₆H₄F), 124.68, 124.81, 124.94, 126.04, 126.21, 126.36, 128.65, and 129.48 (*m*-CH of Mes and CH of CR₂), 128.39 (d, $J_{CF} = 8.3$ Hz, CH of C₆H₄F), 131.00 (d, $J_{CF} = 3.5$ Hz, CH of C₆H₄F), 130.92, 132.81, 138.16, 138.71, 141.29, 141.58, 142.78, 143.42, 143.69, and 144.43 (ipso-, o-, and p-C of Mes and quat. C of CR₂), 159.67 (d, ${}^{1}J_{CF} = 244.5$ Hz, CF); ¹⁹F NMR (CDCl₃) δ –39.0; MS, m/z (%) 477 (Mes₂GeCHR₂, 2), 446 (M - CHR₂, 4), 312 (Mes₂Ge, 10), 192 (MesGe - 1, 41), 165 (CHR₂, 100). Anal. Calcd for C₃₉H₃₆FGeN: C, 76.75; H, 5.95. Found: C, 76.51; H, 6.05.

9c: white crystals; 1.06 g, yield 86%; mp 248 °C; ¹H NMR (CDCl₃) δ 1.78 and 1.85 (2s, 2 × 6H, *o*-Me), 2.15 and 2.20 (2s, 2 × 3H, *p*-Me), 4.70 (s, 1H, CH–C=N), 5.18 (s, 1H, C*H*R₂), 6.48 and 6.60 (2s, 2 × 2H, arom H of Mes), 6.82–8.01 (m, 12H, arom H of C₆H₄F and CR₂); ¹³C NMR (CDCl₃) δ 20.78 and 20.82 (*p*-Me), 24.79 and 24.98 (*o*-Me), 29.2 (*C*HC=N), 45.84 (*C*HR₂), 115.36 (d, ²J_{CF} = 21.1 Hz, *m*-CH of C₆H₄F), 119.97 and 120.00 (CH of CR₂), 122.66 (C=N), 124.32, 124.82, 125.84, 125.91, 126.18, 129.01, 129.37, 129.69, and 129.79 (*m*-CH of Mes, CH of CR₂, *o*-CH of C₆H₄F), 130.03, 131.40, 131.80, 138.54, 138.72, 141.45, 142.98, 143.23, 143.91, and 144.15 (*ipso-*, *o-*, and *p*-C

⁽¹⁴⁾ Sheldrick, G. M. *Acta Crystallogr.* **1990**, *A46*, 467–473. (15) Sheldrick, G. M. *SHELXL-97*, Program for Crystal Structure Refinement; University of Göttingen, 1997.

of Mes, quat. C of CR₂, *ipso*-C of C₆H₄F), 161.78 (d, ${}^{1}J_{CF} = 246.0$ Hz, CF); ${}^{19}F$ NMR (CDCl₃) δ -39.9; MS, m/z (%) 477 (Mes₂-GeCHR₂, 19), 446 (M - CHR₂, 45), 312 (Mes₂Ge, 14), 192 (MesGe - 1, 48), 165 (CHR₂, 100). Anal. Calcd for C₃₉H₃₆-FGeN: C, 76.75; H, 5.95. Found: C, 77.01; H, 5.80.

9d: light pink crystals; 0.95 g, yield 79%; mp 255 °C; ¹H NMR (CDCl₃) δ 1.76 and 1.96 (2s, 2 × 6H, *o*-Me), 2.14 and 2.21 (2s, 2 × 3H, *p*-Me), 4.91 (s, 1H, *CH*R₂), 5.31 (s, 1H, CH–C=N), 6.46 and 6.64 (2s, 2 × 2H, arom H of Mes), 6.50–8.01 (m, 11H, arom H of thienyl and CR₂); ¹³C NMR (CDCl₃) δ 20.85 (*C*HC=N), 20.94 and 24.47 (*p*-Me), 25.06 and 25.32 (*o*-Me), 46.19 (*C*HR₂), 97.54 (C=N), 119.94 and 120.23 (CH of CR₂), 121.83 (*C*SCH) 123.95, 124.45, 124.77, 126.04, 126.12, 126.41, 127.25, and 127.64 (CH of CR₂ and thienyl), 129.01 and 129.50 (*m*-CH of Mes), 131.11, 135.02, 138.47, 138.74, 141.40, 143.13,

143.55, 143.96, and 144.31 (quat. C of CR₂, *ipso-*, *o-* and *p-*C of Mes); IR (KBr) 2222.7 cm⁻¹ (ν C \equiv N); MS, *m*/*z* (%) 477 (Mes₂-GeCHR₂, 30), 434 (M - CHR₂, 22), 311 (Mes₂Ge - 1, 23), 192 (MesGe - 1, 41), 165 (CHR₂, 100). Anal. Calcd for C₃₇H₃₅-GeNS: C, 74.27; H, 5.90. Found: C, 74.50; H, 6.02.

Acknowledgment. Financial support of this work by the CMIFM (AI 00/216/SM) is gratefully acknowledged.

Supporting Information Available: Table of bond lengths and bond angles for **7**. This material is available free of charge via the Internet at http://pubs.acs.org.

OM0494622