

Synthesis of Novel Azaspiro[3.4]octanes as Multifunctional Modules in Drug Discovery


Dong Bo Li,[†] Mark Rogers-Evans,^{*,‡} and Erick M. Carreira^{*,†}

Laboratorium für Organische Chemie, ETH Zürich, CH-8093 Zürich, Switzerland, and
Discovery Research, Chemistry Technologies & Innovation, F. Hoffmann-La Roche AG,
CH-4070 Basel, Switzerland

mark.rogers-evans@roche.com; carreira@org.chem.ethz.ch

Received September 19, 2011

ABSTRACT



Step-economic and scalable syntheses of novel thia-azaspiro[3.4]octanes are reported. These spirocycles and some related intermediates can serve as uncharted multifunctional modules for drug discovery chemistry.

The identification of novel building blocks through design and synthesis constitutes a critical role for organic chemistry in the drug discovery process. We have recently described a collection of spirocyclic building blocks,^{1–3} which provide entry into novel chemical, pharmacological, and proprietary space. In line with these interests we continue to expand the collection of spirocycles by exploring new structures. Characteristics of these “compact modules” are that they should have tunable polarities affecting

the phys-chem and safety parameters as well as that they should be amenable to vectorization. The latter provides ready shape diversity, a feature critical to contemporary compound collections looking to access poorly or uncharted 3D-macromolecular biological targets. In this context, our recent pursuits have included thia-azaspiro[3.4]octanes (Figure 1). The concept driving our interest in these structures initially was their similarity to the parent diazaspiro[3.3]heptanes^{2b,c} and newly developed azaspiro[3.3]heptane scaffolds^{2a} in our group (Figure 1) and the fact that these position functional groups and attendant exit vectors in a cambered environment. Additionally, we reasoned that the inclusion of a heteroatom, such as sulfur, particularly in its oxidized state, at different positions of the spirocycle would serve to generate uncharted spiro[3.4]octanes that intuitively resembled “drug-like” fragments. Herein, we report convenient constructions of the novel thia-azaspirocyclic systems.

[†] ETH Zürich.

[‡] F. Hoffmann-La Roche AG.

(1) For oxetanes and related spirocycles, see: (a) Wuitschik, G.; Rogers-Evans, M.; Müller, K.; Fischer, H.; Wagner, B.; Schuler, F.; Polonchuk, L.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2006**, *45*, 7736–7739. (b) Wuitschik, G.; Rogers-Evans, M.; Buckl, A.; Bernasconi, M.; Märki, M.; Godel, T.; Fischer, H.; Wagner, B.; Parrilla, I.; Schuler, F.; Schneider, J.; Alker, A.; Schweizer, W. B.; Müller, K.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2008**, *47*, 4512–4515. (c) Wuitschik, G.; Carreira, E. M.; Rogers-Evans, M.; Müller, K. Oxetan-3-one: chemistry and synthesis. In *Process Chemistry in the Pharmaceutical Industry*; Gadamasetti, K., Braish, T., Eds.; CRC Press: Boca Raton, FL, 2008; pp 217–229. (d) Wuitschik, G.; Carreira, E. M.; Wagner, B.; Fischer, H.; Parrilla, I.; Schuler, F.; Rogers-Evans, M.; Müller, K. *J. Med. Chem.* **2010**, *53*, 3227–3246. (e) Burkhard, J. A.; Wuitschik, G.; Rogers-Evans, M.; Müller, K.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2010**, *49*, 9052–9067.

(2) For azaspiro[3.3]heptanes, see: (a) Guérot, C.; Tchitchanov, B. H.; Knust, H.; Carreira, E. M. *Org. Lett.* **2011**, *13*, 780–783. (b) Burkhard, J. A.; Wagner, B.; Fischer, H.; Schuler, F.; Müller, K.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2010**, *49*, 3524–3527. (c) Burkhard, J. A.; Guérot, C.; Knust, H.; Rogers-Evans, M.; Carreira, E. M. *Org. Lett.* **2010**, *12*, 1944–1947. (d) Burkhard, J.; Carreira, E. M. *Org. Lett.* **2008**, *10*, 3525–3526.

(3) For other recent related work, see: (a) Xu, R.; Czarniecki, M.; de Man, J.; Pan, J.; Qiang, L.; Root, Y.; Ying, S.; Su, J.; Sun, X.; Zhang, Y.; Yu, T.; Zhang, Y.; Hu, T.; Chen, S.-H. *Tetrahedron Lett.* **2011**, *52*, 3266–3270. (b) Meyers, M. J.; Muizebelt, I.; van Wiltenburg, J.; Brown, D. L.; Thorarensen, A. *Org. Lett.* **2009**, *11*, 3523–3525. (c) Duncton, M. A. J.; Estiarte, M. A.; Johnson, R. J.; Cox, M.; O'Mahony, D. J. R.; Edwards, W. T.; Kelly, M. G. *J. Org. Chem.* **2009**, *74*, 6354–6357. (d) Duncton, M. A. J.; Estiarte, M. A.; Tan, D.; Kaub, C.; O'Mahony, D. J. R.; Johnson, R. J.; Cox, M.; Edwards, W. T.; Wan, M.; Kincaid, J.; Kelly, M. G. *Org. Lett.* **2008**, *10*, 3259–3262.





Figure 1. Azaspiro[3.4]octanes incorporating sulfones shown with exit vectors.

The first targets for synthesis were 8-functionalized 5-thia-2-azaspiro[3.4]octanes **5** and **7** (Scheme 1) from commercially available 1-*N*-Boc-3-azetidinone⁴ as a starting point. Thus, aldol addition of 4-butyrothiolactone with 1-*N*-Boc-3-azetidinone produced adduct **1** on an 8-g scale in 85% yield. Subsequent dehydration provided conjugated thiolactone **2** (82%). The key spirocyclization was then realized upon exposure of **2** to alkaline methanol solution, which produced **3b**. We hypothesize the observed formation of **3b** from **2** proceeds by methanol-triggered opening of the thiolactone⁵ followed by intramolecular conjugate addition onto the unsaturated ester in **3a**. After removal of volatiles, the unpurified spirocyclic ester **3b** was directly subjected to oxidation with mCPBA to afford sulfone **4** in 70% yield (two steps). Saponification furnished the corresponding acid **5** (86%). When **5** was subjected to the Curtius rearrangement protocol, orthogonally protected bisamine **6** was isolated in 91% yield. Finally, after hydrogenolytic removal of the Cbz-group, amine **7** was obtained in 82% yield.

In order to render the new spirocycles more versatile, further manipulations were investigated that permitted installation of carbonyl and alcohol groups. However, attempts to implement Barton's decarboxylative oxygenation⁶ method on acid **5** were unsuccessful, despite the clean formation of the intermediate thiopyridone adduct. Examination of ketone derivatives under conditions leading to Bayer–Villiger rearrangement did not meet with success. Our attention was then shifted to developing a de novo route analogous to the approach to thiolanes reported by Warren.⁷

As shown in Table 1, the first step in the implementation of Warren's method was to access an efficient sulfenylation

Scheme 1. Synthesis of Spirocyclic Modules **5** and **7**



to reach key intermediates **8a** or **8b**. Although access to **8a** was possible by conversion of commercially available 1-*N*-Boc-3-azetidinone into the corresponding homologous enol ether, which was then subjected to the sulfenylation with BnSCl to provide the required aldehyde **8a**, this two-step protocol provided **8a** in only 16% poor yield (entry 1, Table 1). Formation of the corresponding enol ether under the Peterson conditions⁸ did not lead to improvements (entry 2, Table 1), and the attempted preparation of silyl enol ether of the parent aldehyde⁹ was unsatisfactory as well (entries 3 and 4,¹⁰ Table 1). Furthermore, the formation of silyl ketene acetal of the parent ester¹¹ was not detected (entry 5, Table 1); rather C-silylation was exclusively observed. The direct sulfenylation of the enolate derived from the parent ester with BnSSBn, by using LiHMDS as a base, yielded an unknown product (entry 6, Table 1). However, ester enolization with LDA allowed direct sulfenylation to give BnS-ester **8b** in 40% yield (entry 7, Table 1). The yield could be improved to 53% with the addition of HMPA as a cosolvent (entry 8, Table 1).¹² It should be emphasized that, unlike 1-*N*-Boc-3-azetidinone, the chemistry of 1-*N*-Boc-3-azetidinecarboxaldehyde and methyl 1-*N*-Boc-3-azetidinecarboxylate is not well-documented. Thus the enolization studies provide important opportunities for this class of building blocks.

Ester **8b** underwent Claisen condensation to provide ketoester **9** in 84% yield (Scheme 2). Although the reduction to 1,3-diol **10** proceeded cleanly with DIBAL, the yield was modest. Following the Warren conditions,⁷ treatment of BnS-1,3-diol **10** with TsCl in pyridine smoothly produced spirocyclic alcohol **13** at room temperature. We hypothesize that the reaction proceeds by [1,4]-SBn participation

(4) (a) From PharmaBlock, catalog no. PB00003. (b) For the chemistry of the ketone, see: Dejaegher, Y.; Kuz'menok, N. M.; Zvonok, A. M.; De Kimpe, N. *Chem. Rev.* **2002**, *102*, 29–60.

(5) The mild, ring opening method has been reported for the synthesis of methyl-4-thio-butanoate, see: De Gregori, A.; Jommi, G.; Sisti, M.; Gariboldi, P.; Merati, F. *Tetrahedron* **1988**, *44*, 2549–2568.

(6) For recent review and references therein, see: Saraiva, M. F.; Couri, M. R. C.; Hyaric, M. L.; de Almeida, M. V. *Tetrahedron* **2009**, *65*, 3563–3572.

(7) Eames, J.; Kuhnert, N.; Warren, S. *J. Chem. Soc., Perkin Trans. 1* **2001**, 1504–1510.

(8) Magnus, P.; Roy, G. *Organometallics* **1982**, *1*, 553–559.


(9) From Apollo Scientific Ltd., catalog no. OR15687.

(10) Tanabe, Y.; Misaki, T.; Kurihara, M.; Iida, A.; Nishii, Y. *Chem. Commun.* **2002**, 1628–1629.

(11) From Apollo Scientific Ltd., catalog no. OR42009.

(12) For HMPA as cosolvent, see: Grieco, P. A.; Flynn, D. L.; Zelle, R. E. *J. Am. Chem. Soc.* **1984**, *106*, 6414–6417.

Table 1. Screening for an Efficient Sulfonylation to Reach Key Intermediate

				
entry	starting material	conditions	product	total yield
1	R ₁ , R ₂ = -O-	1) Ph ₃ P ⁺ Cl ⁻ CH ₂ OMe 2) BnSCl	8a	16%
2	R ₁ , R ₂ = -O-	1) TMSCH ₂ OMe 2) KH 3) BnSCl	8a	15%
3	R ₁ = CHO R ₂ = H	1) TMSCl, Et ₃ N 2) BnSCl	8a	trace
4	R ₁ = CHO R ₂ = H		N.D. ^a	– ^b
5	R ₁ = CO ₂ Me R ₂ = H	TMSCl, LDA	N.D. ^a	– ^b
6	R ₁ = CO ₂ Me R ₂ = H	BnSSBn, LiHMDS	N.D. ^a	– ^b
7	R ₁ = CO ₂ Me R ₂ = H	BnSSBn, LDA	8b	40%
8	R ₁ = CO ₂ Me R ₂ = H	BnSSBn, LDA, HMPA	8b	53%

^a Not desired product. ^b The yield was not determined.

and debenzoylation with the chloride generated *in situ*.⁷ Finally, oxidation with *m*CPBA furnished sulfonyl alcohol **14** in 67% yield, and subsequent oxidation with DMP proceeded very well to provide the corresponding sulfonyl ketone **15**.

We next examined the construction of 6-thia-2-azaspiro-[3.4]octanes (**20** and **21**, Scheme 3). The aldol addition reaction involving BnSCH₂CHO¹³ and protected azetidine ester produced adduct **16** on gram scale in acceptable yield. Azetidine **16** underwent stepwise reduction (DIBAL-NaBH₄) to afford 1,3-diol **17**. Subsequent treatment of 1,3-diol **17** with TsCl in pyridine produced a mixture of tosylate **18** along with desired spirocyclic thiolane **19**. The complete conversion of the intermediate tosylate into the spirocyclic thiolane could be smoothly realized (50% for **19** overall from **16**) by heating in DMF. Finally, oxidation with *m*CPBA gave sulfone alcohol **20** in excellent yield. Parikh–Doering oxidation¹⁴ of **19** proceeded well to provide the corresponding sulfonyl ketone **21** in 63% yield.

In summary, we have described the synthesis of six new multifunctional modules through the use of highly step-economic sequences. The syntheses of 5-thia-2-azaspiro-

Scheme 2. Synthesis of Spirocyclic Modules **14** and **15**

