## Electrochemical Carboxylation of *N*-(2-Bromopropionyl)-4*R*phenyloxazolidin-2-one: An Efficient Route to Unsymmetrical Methylmalonic Ester Derivatives

2002 Vol. 4, No. 16

2617 - 2620

ORGANIC LETTERS

Marta Feroci,\*,† Achille Inesi,\*,† Monica Orsini,† and Laura Palombi‡

Dipartimento di Ingegneria Chimica, Materiali, Materie Prime e Metallurgia, Università "La Sapienza", via Castro Laurenziano, 7, I-00161 Roma, Italy, and Dipartimento di Chimica, Ingegneria Chimica e Materiali, Università degli Studi, I-67040, Monteluco di Roio, L'Aquila, Italy

marta.feroci@uniroma1.it

Received March 28, 2002

ABSTRACT



major diastereoisomer

Electrochemical carboxylation of *N*-(2-bromopropionyl)-4*R*-phenyloxazolidin-2-one aimed at the synthesis and chiral resolution of unsymmetrical methylmalonic ester derivatives is described. The presence of the Evans' chiral auxiliary permits an easy resolution of the mixture of the two diastereoisomers.

Optically active  $\alpha$ -monoalkylmalonic derivatives are valuable building blocks in asymmetric synthesis, providing access to unsymmetrical 2-substituted 1,3-propandiols in an enantiomerically enriched form.<sup>1,2</sup> The common chemical approach to these chiral malonates in enantiopure form consists of the resolution of the diastereoisomeric mixtures by means of a chiral resolving agent (typically an optically active alcohol<sup>1</sup> such as menthol or 8-phenylmenthol). Menthyl and 8-phenylmenthyl half-esters of methylmalonic acid have been obtained in 62 and 60% yields, respectively, and were composed of the two isomers in a ~3:2 ratio. They could be converted into the corresponding chiral diol derivative by reduction (after activation).<sup>1</sup>

The electrochemical methodology permits the carboxylation<sup>3</sup> in the  $\alpha$ -position to carbonyl groups by direct reduction of  $\alpha$ -haloketones in the presence of carbon dioxide<sup>4,5</sup> (50– 57% yields in  $\alpha$ -ketocarboxylic acids) or by direct reduction of vinyl triflates also in the presence of carbon dioxide<sup>6,7</sup> (28–77% yields in cyclic  $\alpha$ -ketocarboxylic acids and 33– 67% yields in linear ones). To our knowledge, no electro-

<sup>\*</sup> Corresponding authors. Phone: +39-06-49766780. Fax: +39-06-49766749. E-mail (A.I.): inesi@ing.univaq.it.

University of Rome.

<sup>&</sup>lt;sup>‡</sup> University of L'Aquila.

<sup>(1)</sup> Ihara, M.; Takahashi, M.; Taniguchi, N.; Yasui, K.; Fukumoto, K.; Kametani, T. J. Chem. Soc., Perkin Trans. 1 **1989**, 897–903.

<sup>(2)</sup> Harada, T.; Hayashiya, T.; Wada, I.; Iwa-ake, N.; Oku, A. J. Am. Chem. Soc. 1987, 109, 527-532.

<sup>(3)</sup> Silvestri, G. In *Carbon Dioxide as a Source of Carbon*; Aresta, M., Forti, G., Eds.; D. Reidel Publishing Company: Dordrecht, Holland, 1987; pp 339–369.

<sup>(4)</sup> Carelli, I.; Curulli, A.; Inesi, A.; Zeuli, E. J. Chem. Res., Synop. 1988, 154–155; J. Chem. Res., Miniprint 1988, 1276–1296.

<sup>(5)</sup> Carelli, I.; Curulli, A.; Inesi, A.; Zeuli, E. J. Chem. Res., Synop. 1990, 74–75; J. Chem. Res., Miniprint 1990, 620–644.

<sup>(6)</sup> Kamekawa, H.; Senboku, H.; Tokuda, M. Tetrahedron Lett. 1998, 39, 1591-1594.

<sup>(7)</sup> Senboku, H.; Fujimura, Y.; Kamekawa, H.; Tokuda, M. *Electrochim. Acta* **2000**, *45*, 2995–3003.

## Table 1. Electrolyses<sup>a</sup> of Solutions of 1 under Different Experimental Conditions

				products (yield, %) <sup>b</sup>			
entry	anode/cathode	solvent/electrolyte	Т	3	4	2a + 2b	$\mathrm{d}\mathbf{r}^{c}$
1	$Mg/Pb^d$	MeCN/Et <sub>4</sub> NClO <sub>4</sub>	rt	33	30	36	57:43
2	$Mg/Pb^d$	MeCN/Et <sub>4</sub> NClO <sub>4</sub>	−10 °C	65	20	15	65:3 <i>5</i>
3	$Mg/Pb^d$	MeCN/Et <sub>4</sub> NClO <sub>4</sub>	−20 °C	72	22	5	83:17
4	Mg/Pb	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C	37		62	$63:37^{e}$
5	Mg/Pb <sup>f</sup>	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C	32	5	40	69:31 <sup>e</sup>
6	Mg/Pb	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−40 °C	22		55	59:41 <sup>e</sup>
7	Mg/Pt	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C			70	60:40 <sup>e</sup>
8	Mg/Pt <sup>f</sup>	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C	53	10	36	68:32
9	Al/Pt	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C	7		78	59:41 <sup>e</sup>
10	$Al/Pt^{f}$	THF/Bu <sub>4</sub> NBF <sub>4</sub>	−20 °C	22		27	$52:48^{e}$

<sup>&</sup>lt;sup>*a*</sup> Compound 1 (2 F/mol), undivided cells, galvanostatic conditions ( $I = 4 \text{ mA/cm}^2$ ). <sup>*b*</sup> Yields of isolated products, calculated with respect to the starting oxazolidinone 1. <sup>*c*</sup> Diastereoisomeric ratio was determined by <sup>1</sup>H NMR. <sup>*d*</sup> Et<sub>3</sub>N was added to the solution prior to electrolysis. These electrolyses were carried out under potentiostatic conditions (E = -1.7 V vs SCE). <sup>*e*</sup> Starting material was recovered at the end of the electrolysis. <sup>*f*</sup> Et<sub>3</sub>N was added to the solution prior to electrolysis.

chemical carboxylation of  $\alpha$ -haloesters or amides has been reported.

Here we report our preliminary results for the electrochemical carboxylation of N-(2-bromopropionyl)-4R-phenyloxazolidin-2-one aimed at the synthesis and chiral resolution of unsymmetrical methylmalonic ester derivatives. The subsequent selective reduction of the amidic function in the presence of the methyl ester group, to yield the corresponding chiral alcohol, is well described in the literature.<sup>8</sup>

The advantage of this kind of approach is that the resolving agent is present in the molecule prior to the introduction of the second carbonyl group by carboxylation, so it allows the transformation of this second function in an acid or ester in an easy way without the side reaction of diester formation.

The starting material was N-(2-bromopropionyl)-4R-phenyloxazolidin-2-one **1** (Scheme 1), both as an epimeric



mixture and as pure diastereoisomers, and the outcome of the reaction was the same in all cases (no memory of chirality was evidenced).

The electrolyses<sup>9</sup> were carried out on solutions (solventsupporting electrolyte) containing 1 in which carbon dioxide was continuously bubbling. Various experimental conditions have been considered, and the results are reported in Table 1. All the electrolysis products are described in Figure 1.



The two diastereoisomers **2a** and **2b** could be easily separated by column chromatography.

The yields in carboxylated products are influenced by various factors: electrolysis temperature (Table 1, entries 1-4 and 6), solvent system (Table 1, entries 3 and 5), and electrode material (cathode, Table 1, entries 4 and 7; anode, Table 1, entries 7 and 9). The higher yields in 2a + 2b (78%, Table 1, entry 9) were obtained using aluminum as a sacrificial anode and platinum as a cathode, in THF/Bu<sub>4</sub>-NBF<sub>4</sub> as a solvent–electrolyte system, at -20 °C under galvanostatic conditions. In this case, the diastereoisomeric ratio between the two carboxylated products was  $\sim$ 3:2. If the reaction was carried out in MeCN–Et<sub>4</sub>NClO<sub>4</sub>,<sup>10</sup> the yields were lower (Table 1, entries 1–3), while the dr increased slightly with decreasing the temperature. MeCN probably

<sup>(8)</sup> Tietze, F.; Schünke, C. Angew. Chem., Int. Ed. Engl. 1995, 34 (16), 1731–1733.





acts as a proton donor; in fact, when this solvent was used, the yields in reduced product 3 were very high to the detriment of those of the carboxylated products.

The role of  $Et_3N$  in this reaction is not so clear:  $Evans^{11}$  reported that  $Et_3N$  forms a kind of aggregated complex with the enolate that derives from the reduction of the C–Br bond, so it should enhance the diastereoselectivity of the carboxylation reaction; on the other hand, the yields in 2a + 2b are considerably lowered by the presence of the amine. The

(9) General Procedure. A solution of 3-(2-bromopropionyl)-4R-phenyl-2-oxazolidinone<sup>16</sup> 1 in 30 mL of THF-0.2 mol/L Bu<sub>4</sub>NBF<sub>4</sub> was electrolyzed (undivided cells, Pt or Pb cathode, Al or Mg anode, at -20 °C) under galvanostatic conditions ( $I = 4 \text{ mA/cm}^2$ ) in the presence of carbon dioxide (p = 1 atm). After the consumption of 2 Faradays per mol of 1, the current flow was stopped, the solvent evaporated under reduced pressure, and the residue poured into water. This aqueous phase was extracted with diethyl ether (3  $\times$  30 mL), and this organic solution was worked up as usual, giving the starting material 1, (R)-(-)-4-phenyl-3-propionyl-2-oxazolidinone 3,<sup>1</sup> and (R)-(-)-4-phenyl-2-oxazolidinone 4, if any. The aqueous solution was then acidified (pH  $\approx$  3) with dilute HCl and extracted again with ether. This second ethereal phase was cooled at 0 °C and treated with ethereal CH<sub>2</sub>N<sub>2</sub>.<sup>18</sup> (CAUTION! Diazomethane is toxic and prone to cause development of specific sensitivity; in addition, it is potentially explosive). The usual workup gave the mixture of 2a and 2b, whose ratio was calculated by <sup>1</sup>H NMR. The two pure isomers were obtained after column chromatography (8:2 n-hexanes-ethyl acetate as the eluent). 3-(2-Methoxycarbonylpropionyl)-4S-phenyl-2-oxazolidinone (Less Polar Isomer) 2a. 1H NMR  $\delta$  (CDCl<sub>3</sub>): 7.40–7.25 (m, 5H, ar), 5.46 (dd, 1H, J = 8.9 Hz, J =3.7 Hz, OCHH), 4.70 (t, 1H, J = 8.9 Hz, N.–CH), 4.53 (q, 1H, J = 7.2Hz,  $CHCO_2CH_3$ ), 4.29 (dd, 1H, J = 8.9 Hz, J = 3.7 Hz, OCHH), 3.71 (s, 3H, CO<sub>2</sub>CH<sub>3</sub>), 1.40 (d, 3H, J = 7.2 Hz, CH<sub>3</sub>CHCO<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR  $\delta$ (CDCl<sub>3</sub>): 170.81, 169.08, 153.69, 138.80, 129.25, 128.85, 125.94, 70.19, 57.79, 52.53, 45.66, 13.19. GC-MS m/z: 277 (M<sup>+,</sup>, 3%), 218 (M<sup>+</sup> - CO<sub>2</sub>-CH<sub>3</sub>, 3%), 162 (73%), 104 (85%), 77 (79%), 59 (100%).  $[\alpha]_{\rm D}^{20}$  -96.9 (c 0.96, AcOEt). 3-(2-Methoxycarbonylpropionyl)-4R-phenyl-2-oxazolidinone (More Polar Isomer) 2b. <sup>1</sup>H NMR  $\delta$  (CDCl<sub>3</sub>): 7.40–7.31 (m, 5H, ar), 5.44 (dd, 1H, J = 9.0 Hz, J = 3.9 Hz, OCHH), 4.71 (t, 1H, J = 9.0Hz, N.-CH), 4.49 (q, 1H, J = 7.2 Hz, CHCO<sub>2</sub>CH<sub>3</sub>), 4.25 (dd, 1H, J = 9.0 Hz, J = 3.9 Hz, OCHH), 3.67 (s, 3H, CO<sub>2</sub>CH<sub>3</sub>), 1.39 (d, 3H, J = 7.2 Hz, CH<sub>3</sub>CHCO<sub>2</sub>CH<sub>3</sub>). <sup>13</sup>C NMR δ (CDCl<sub>3</sub>): 170.67, 168.81, 153.68, 138.39, 129.08, 128.70, 125.82, 70.30, 57.88, 52.36, 45.55, 12.95. GC-MS m/z: 277 (M<sup>+</sup>, 3%), 218 (M<sup>+</sup> - CO<sub>2</sub>CH<sub>3</sub>, 3%), 162 (73%), 104 (85%), 77 (79%), 59 (100%).  $[\alpha]_{\rm D}^{20}$  -44.6 (*c* 0.87, AcOEt).

(10) CAUTION! Although in more than 100 experiments no particular safety problem has been encountered, the use of perchlorates in organic solvent must be considered as potentially dangerous (explosive). Take adequate precautions.

(11) Evans, D. A.; Rieger, D. L.; Bilodeau, M. T.; Urpì, F. J. Am. Chem. Soc. **1991**, *113*, 1047–1049.



presence of **4** among the electrolysis products is probably due to the decomposition of the enolate ion via a ketene pathway.<sup>12</sup>

When the ammonium salt was replaced by LiClO<sub>4</sub>, no carboxylated product could be isolated from the electrolyzed solution. Only 6% in 2a + 2b was obtained working with divided cells in the same experimental conditions as those of entry 9 (Table 1).

In all cases, the more abundant diastereoisomer<sup>13</sup> is **2a**, as demonstrated by the X-ray analysis (Figure 2). This is probably due to the preferential "non-Evans" conformation<sup>14</sup> (that is, the nonchelated one) of the enolate ion during the carboxylation (Figure 3). In fact, in our experimental conditions, the concentration of the metallic ion, which derives from the consumption of the sacrificial anode and could chelate both oxygen atoms, is probably too low with respect to the concentration of the supporting electrolyte cation  $R_4N^+$ , which can coordinate but not chelate both carbonyl oxygen atoms. It is also possible that in a solvent such as THF, the metallic cation is not free but ion-paired with the bromide released during the electrolysis.

When the two diastereoisomeric esters (2a and 2b) were allowed to epimerize separately (by reaction with Et<sub>3</sub>N in CD<sub>3</sub>CN), the equilibrium ratio of 2a:2b was 2:3 in both cases, so we can deduce that in our experimental conditions the reaction is partially controlled by kinetic factors.

Any attempt to carboxylate by electrochemical means (using electrogenerated bases) the starting material not containing the bromine atom (i.e., *N*-propionyl-4*R*-phenyl-oxazolidin-2-one) failed, so the use of a halogenated derivative seems necessary.<sup>15</sup>

(17) Ager, D. J.; Allen, D. R.; Schaad, D. R. Synthesis 1996, 1283–1285.

(18) de Boer, T. J.; Backer, H. J. Organic Syntheses; Wiley: New York, 1963; Collect. Vol. IV, pp 250-253.

<sup>(12)</sup> Evans, D. A.; Ennis, M. D.; Mathre, D. J. J. Am. Chem. Soc. 1982, 104, 1737-1739.

<sup>(13)</sup> This kind of molecule exhibits a quite low kinetic acidity; they are, in fact, quite stable to silica gel chromatography and can be isolated in pure enantiomeric form. See: Evans, D. A.; Ennis, M. D.; Le, T. J. Am. Chem. Soc. **1984**, 106, 1154–1156.

<sup>(14)</sup> Yan, T.-H.; Tan, C.-W.; Lee, H.-C.; Lo, H.-C.; Huang, T.-Y. J. Am. Chem. Soc. **1993**, 115, 2613–2621.

<sup>(15)</sup> Any attempt to obtain 2a + 2b by metalation and carboxylation (using Mg or Zn under classical reaction conditions, followed by the addition of CO<sub>2</sub>) failed.

<sup>(16)</sup> Ito, Y.; Sasaki, A.; Tamoto, K.; Sunagawa, M.; Terashima, S. *Tetrahedron* **1991**, *47*, 2801–2820.

Table 2. Electrochemical Carboxylation of N-(α-Bromoacyl)-4R-phenyloxazolidin-2-ones<sup>a</sup>



<sup>*a*</sup> Starting bromide (2 F/mol). Electrolysis conditions: entry 9, Table 1. <sup>*b*</sup> Yields of isolated products, calculated with respect to the starting bromide. <sup>*c*</sup> Diastereoisomeric ratio was determined by <sup>1</sup>H NMR. <sup>*d*</sup> Chemical shifts of the hydrogen atoms of the methoxycarbonyl group (<sup>1</sup>H NMR, ppm with respect to tetramethylsilane) of the major and minor diastereoisomers, respectively. <sup>*e*</sup> This electrolysis is reported in Table 1 (entry 9), and cited here for comparison. <sup>*f*</sup> Starting material was a chloride (instead of a bromide), and the electrolysis was stopped after 3 F/mol.

To test the applicability of this electrochemical carboxylation, the investigation was extended to *N*-( $\alpha$ -bromoacyl)-4*R*-phenyloxazolidin-2-ones other than the title one (see Table 2). As expected, when the alkyl group is replaced by a phenyl one, the electrochemical carboxylation does not take place and only the reduction product can be detected. Further investigations are necessary to improve these results.

In conclusion, the electrochemical methodology permits the synthesis in good yields and the easy resolution of methylmalonic ester derivatives by reduction and carboxylation of (4R)-3-(2-bromopropionyl)-4-phenyl-2-oxazolidinone.

Acknowledgment. This work was supported by research grants from M.U.R.S.T. (Cofin 2000) and C.N.R., Rome, Italy. The authors thank Mr. M. Di Pilato for his contribution to the experimental part of this work and Prof. Fabio Marchetti for the X-ray analysis.

**Supporting Information Available:** Characterization data for all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

OL025939Z