

Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: S. Batsyts, E. G. Hübner, J. C. Namyslo, M. Gjika and A. Schmidt, *Org. Biomol. Chem.*, 2019, DOI: 10.1039/C9OB00357F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Synthesis and characterization of propeller-shaped mono- to hexacationic quinolinium-substituted benzenes

Sviatoslav Batsyts,^a Eike G. Hübner,^a Jan C. Namyslo,^a Mimoza Gjika,^b and Andreas Schmidt^{a*}

Received 00th January 2019,
Accepted 00th January 2019

DOI: 10.1039/x0xx00000x


www.rsc.org/

Diels-Alder reaction of 2-, 3- and 4-(phenylethynyl)quinolines and tetraphenylcyclopentadienone gave three regioisomeric 2,3,4,5,6-pentaphenyl-1-(quinolin-2-yl, -3-yl, and -4-yl)benzenes. Restricted rotation of the 3-yl and 4-yl substituted derivatives is observed between the central core and the substituents, resulting in propeller-shaped molecules. Likewise, 1,2-diquinolyl-3,4,5,6-tetraphenylbenzenes with 3-yl,3-yl and 3-yl,4-yl connectivity were prepared. As evidenced by NMR spectroscopy, they form two diastereomers due to their restricted rotation. A cobalt-catalyzed [2+2]-cyclootrimerization of 2-(phenylethynyl)quinoline resulted in the formation of triphenyl-2,4,6- and -3,5,6-tri(quinolin-2-yl)benzenes. The same reaction was applied to 3,3'-ethyne-1,2-diylquinoline which formed hexa(quinolin-3-yl)benzene. *N*-Methylation gave the title compounds. Among those, the hexacationic hexa(*N*-methylquinolinio-3-yl)benzene is described. Stereochemical aspects are predominantly discussed by means of results of NMR experiments. DFT-calculations on the most stable conformations and the frontier orbital profiles of the hexacation as well as of its neutral precursor have been carried out.

Introduction

Considerable attention has been directed toward heterocycle polycations as they are interesting in natural product chemistry, heterocyclic chemistry, and materials chemistry. Thus, polypyridinium alkaloids (halitoxin,¹ cyclostelletamin C,² amphitoxin,³ viscosamine,⁴ viscosaline,⁵ pachychalines A-C⁶) are polycationic molecules from nature. Streitwieser et al. reported on series of polypyridinium salts and betaines which have adjacent heteroarene rings in conjugation.⁷ Thus, the five-fold pyridinium substituted cyclopentadiene anion **1**⁸ possesses structure elements of conjugated mesomeric betaines. The SASAPOS protocol (self-activated silyl-assisted polyon substitution) by Weiss et al. allowed for the synthesis of a variety of heteroarene substituted substrates, for which **2** is given as an example here.⁹ This widely applicable protocol¹⁰ takes advantage of the fact that a substrate which bonds neutral ligands such as chloride undergoes a substitution with heteroaromatic nucleophiles equivalent to the quantity of the bonding ligands in the presence of the same number of equivalents of trimethylsilyltriflate (TMSOTf). We reported on heterocycle polycations with heteroaromatic central cores such as pyridine,¹¹ pyrimidine,¹² pyrazine,¹³ pyridazine,¹³ 1,3,5-triazine,¹³ and purine.¹³ An example is pentacation **3**.¹⁴ Pyridine polycations proved to be versatile starting materials for the synthesis of highly substituted pyridines with various substitution patterns.¹¹ Apart from their synthetic applicabilities, heterocycle polycations are of interest as

potential semiconductors,¹⁵ photosensitive materials,¹⁶ oxidants,¹⁷ and biologically active compounds like herbicides,¹⁸ acetylcholinesterase reactivators,¹⁹ and cholinesterase inhibitors.²⁰ A recent review article summarizes results achieved so far.²¹



Scheme 1. Examples of heterocycle polycations and of propeller-shaped molecules

^a Clausthal University of Technology, Institute of Organic Chemistry, Leibnizstraße 6, D-38678 Clausthal-Zellerfeld, Germany. Email: schmidt@ioc.tu-clausthal.de

^b Clausthal University of Technology, Institute of Inorganic and Analytical Chemistry, Paul-Ernst-Straße 4, D-38678 Clausthal-Zellerfeld, Germany.


Electronic Supplementary Information (ESI) available: Calculations, NMR spectra. See DOI: 10.1039/x0xx00000x



Steric hindrance exerted by the peripheral ligands cause propeller-shaped molecules. Recently, the sterically congested hexa(heteroaryl)benzene (HHAB) **4** was formed as two isolable isomers which differ only in the rotation of one methyl-pyridine group.²² In general, propeller-shaped hexaarylbenzenes (HABs) such as hexa(β -naphthyl)benzene **5**²³ have numerous applications in materials sciences as they play roles as liquid crystals,²⁴ microporous organic solids,²⁵ molecular capsules,²⁶ supramolecular electronic materials,²⁷ molecular rotors,²⁸ nonlinear optical materials,²⁹ metal sensors,³⁰ redox materials,³¹ and molecular wires.³² We report here on quinolinium-substituted benzenes which combine the features of HHABs and polycations as they have to adopt propeller-shaped configurations due to the additional benzo-annulation in comparison to their pyridinium derivatives.

Results and discussion

First, Sonogashira-Hagihara coupling reactions were used to prepare three isomeric ethynyl-substituted quinolines as starting materials for the synthesis of monoquinolinyl-substituted pentaphenylbenzenes, which are of potential interest as ligands of organic electroluminescent device materials.³³ Thus, the 2-, 3-, and 4-halosubstituted quinolines **6a-c** and phenylacetylene **7** were reacted to give the quinolines **8a-c** in good yields (Scheme 2). With these compounds in hand, we examined the [4+2]-cycloaddition with tetraphenylcyclopentadiene **9**. The reaction did not proceed under a variety of different conditions that were tested, among those reflux temperature in benzene, toluene, and xylene, respectively. On increasing the reaction temperature to 305 °C by using benzophenone as a solvent, however, the quinolines **10a-c** were finally prepared within 30 min. in moderate yields.



Scheme 2. Synthesis of monoquinoline-substituted HHABs.

Similar to hexaphenylbenzene, the six peripheral rings of **10a-c** cannot lie in the plane of the central benzene ring. Whereas in solution, on the NMR time scale, the peripheral rings are perpendicular to the plane of the central ring in the absence of appropriate substitutions, the X-ray structure of hexaphenylbenzene itself showed a propeller conformation with angles around the Ph-Ph bonds of approximately 65°. ³⁴ It is

known that methyl and methoxy groups in C-Ar systems in *ortho* position cause a barrier of rotation of approximately 33 kcal/mol, whereas this value is considerably decreased to approximately 17 kcal/mol for the case of *meta*-substitutions.²³ Hexaphenylbenzene C₆H₆ consequently displays one set of signals in the ¹³C NMR spectra and one overlapped signal with a center of gravity at 6.83 ppm (30H) in the proton resonance spectra.³⁵ The chemical shifts of **10a-c** are different as a consequence of their isomerism and, in addition, intermolecular interactions (Fig. 1). The broadening of the signals at approximately 7.00 ppm in the spectrum of **10a** (blue) can be attributed to N...H-C interactions to the nitrogen atom, which is not possible in **10b,c**. These interactions are thought to be within the limits of hydrogen bonds and classical van der Waals contacts.³⁶ The upfield shift of phenyl ring signals due to intramolecular π -interactions caused by the quinolin-4-yl substitution of **10c** can clearly be seen (black), which are less in **10a,b**.





Figure 1. Comparison of ¹H NMR spectra of HHABs **10a** (1), **10b** (2), and **10c** (3) (DMSO-d₆, 25 °C).

The steric hindrance of the quinoline rings can be compared with 1-(3,4-dimethylphenyl)-2,3,4,5,6-pentaphenylbenzene³⁷ for **10a,b** and 1-(2,3-dimethylphenyl)-2,3,4,5,6-pentaphenylbenzene, which seems to be unknown, for **10c**. However, the quinolin-4-yl-substitution pattern of **10c** imitates the steric hindrance of the methyl group in the model compound 1-(2-methylphenyl)-2,3,4,5,6-pentaphenylbenzene which has a propeller-type topology in the solid³⁸ and a slow rotation of the *o*-tolyl group 25 °C in solution.³⁹ For the case of free rotation under the measurement conditions, 25 distinct ¹³C NMR resonance frequencies can be expected for **10a-c**, among those 15 signals of CH groups. For the case of restricted rotation, this number is increased to 29 ¹³C NMR signals in total, and 19 signals of CH groups, because the *ortho*- and *meta*-positions of the 2- and 3-phenyl substituents become non-isochronous. The least sterically hindered compound **10a** displays 22 distinct signals due to overlapping in DMSO-d₆ at 25 °C. Compound **10b**




shows the expected 29 signals for a restricted rotation, and **10c** displays in total 27 distinct resonance frequencies under the same measurement conditions, presumably due to overlapping of two signals.

The same protocol was applied to the reaction of quinoline **14** with two alkynyl residues which was prepared in 21% yield in three steps from 1,4-dibromobenzene **11** via **12** and **13** as shown (Scheme 3). Quinoline **14** reacted with an excess of cyclopentadienone **9** to give compound **15** which precipitated as exclusive reaction product during the work-up procedure in good yield.




Scheme 3. Synthesis of a bulky monoquinoline HHAB.

The method is also applicable to prepare isomeric diquinoliny substituted tetraphenylbenzenes (Scheme 4). Thus, under analogous conditions, the compound **19a** with quinolin-3-yl/3-yl connectivity and the isomeric quinolin-3-yl/4-yl derivative **19b** were prepared in 90% and 78% yield, respectively, starting from **18a,b** which are available starting from **6b,c** and **17** by standard procedures. The quinolin-3-yl,3-yl compound **19a** can exist in two diastereomeric forms due to restricted rotation, a C_s conformation and a racemic pair with C_2 symmetry, similar to hexaarylbenzenes with two meta-substituents at 0 °C on the NMR time scale.^{34,40} The ^1H NMR spectrum of **19a** showed 37 of expectable 48 ^{13}C NMR signals. Two sets of partially overlapped signals can be identified in a ratio of 1:0.8 under the measuring conditions (DMSO- d_6 , 25 °C). The 4-H hydrogen atom of the quinoline residues of one isomer appears at 7.97 ppm, whereas the other is detectable at 7.93 ppm. The ^{13}C NMR spectrum of isomer **19b**, which possesses a quinolin-3-yl,4-yl connectivity, shows 72 distinct signals of expectable 96. This number is due to the fact, that the 3-yl,4-yl connectivity causes an additional non-symmetry of the molecule and consequently non-isochronous substituents of the two magnetically inequivalent isomers. Under the same measuring conditions, two isochronous rotameric forms are present in a ratio of 1:1 in the NMR spectrum of **19b**.



Scheme 4. Synthesis of diquinoline-substituted HHABs.


For the synthesis of HHABs with three quinoline residues another approach was applied. Thus, the asymmetric acetylene **8a** was subjected to a cobalt-catalyzed [2+2+2]-trimerization reaction which has already been applied for the synthesis of HABs before (Scheme 5).⁴¹ The trimerization of **8a** gave a mixture of two separable regioisomers **20a** and **20b** (1:3) with a total yield of 40%. Compound **20a** is a heteroaromatic analogue of 1,3,5-tri(α -naphthyl)-benzene, a propeller-like, non-planar molecule that is known to interlock in the melt.⁴² It displays 15 ^{13}C NMR signals, similar to the corresponding number of signals of D_{3h} symmetric oligophenylenes which were expected within the fast exchange limit for all single bond rotations.⁴³ Similar to the spectra of the mono(quinolin-2-yl)derivative **10a**, the signals of the *ortho*- and *meta*-protons appear as broad singlets due to N...H-C interactions. Assignment of all atoms using 2D spectra was possible; one set of proton signals for all three quinoline residues of **20a** was found. It is known that *ortho*-substitutions, which cannot be taken as a model for the quinolin-2-yl substitution, can prevent rapid interconversion of the C_{3v} and C_s isomeric forms which are the result of the formation of *syn*- and *anti*-conformations.⁴⁴ The 1,2,4-substituted compound **20b** displays 36 of expectable 45 distinct ^{13}C NMR resonance frequencies in DMSO- d_6 at 25 °C, if free rotation under the measurement conditions is presumed. Similar to compound **10a**, a considerable broadening of the ^1H NMR signals of the quinolines in DMSO- d_6 at 25 °C as well as the phenyl rings is observable.



Scheme 5. Cobalt-catalyzed [2+2+2]-cyclotrimerization.




Trimerization of the acetylene **18a** was tested to prepare a hexaquinolin-3-yl substituted benzene. Indeed, HHAB **21** was formed in 85% yield (Scheme 6). Unfortunately, due to the rather limited solubility of **21** NMR analyses were not possible. Nonetheless, the corresponding HRMS ($m/z = 863.2896$) is in accord with the structure of **21** which can exist in eight rotameric forms. Literature-known²³ calculated minimized energies of hexa(β -naphthyl)benzene, the non-heteroaromatic analogue of **21**, revealed that the most stable rotamer is the one in which all β -naphthyl residues are twisted onto the same side (6,0). The X-ray crystal analysis of hexa(2-pyridyl)benzene, however, revealed the $\alpha,\beta,\alpha,\beta,\alpha,\beta$ arrangement of pyridine rings and dihedral angles between the pyridyl substituent and the benzene ring of approximately 90° .²²



Scheme 6. Cobalt-catalyzed [2+2+2]-cyclotrimerization of **18a**.


Synthesis of polycationic HABs

Based on former experiences of our group,⁴⁵ dimethyl sulfate was used as a methylation agent to convert the quinoline derivatives into cationic species. First, the monoquinoline substituted HABs **10a-c** were successfully methylated in anhydrous toluene under reflux conditions (Scheme 7). The N-methyl group of **22a** interlocks the molecule and 19 distinct ^{13}C NMR signals of CH groups are detectable plus the signals of the anion and the methyl group at 52.8 and 42.5 ppm, respectively. Thus, in contrast to **10a-c**, **22a-c** all display the same number of signals. The N-methylquinolinium salts of **15** with methylsulfate (**23**) and hexafluorophosphate anions (**23PF₆**) were prepared as well, and the diquinoline HABs **19a,b** were also successfully methylated. An immediate anion exchange reaction during work-up yielded the dihexafluorophosphate salts **24a,b**. Similar to the non-methylated precursors **19a** and **19b**, the number of ^{13}C NMR resonance frequencies of **24a** and **24b** is 38 and 72, respectively. The 3-yl,3-yl salt **24a** has an almost identical ratio of rotamers (1:0.9) as its non-methylated precursor, and the corresponding ratio of the salt **24b** in $\text{DMSO-}d_6$ is 1:1.27. In summary, except for **10a**, no spectroscopically detectable changes of the symmetry of the molecules is caused by the methylations. On methylation, the UV/Vis absorption maxima display bathochromic shifts. The spectra are shown in the Supporting Information.



Scheme 7. Methylation of quinoline-based HABs.

Finally, N-methylation of **21** with an excess of dimethyl sulfate followed by precipitation with NH_4PF_6 successfully gave the fully methylated hexacationic HHAB **25** (Scheme 8) which is soluble in $\text{DMSO-}d_6$ and which could be characterized completely.



Scheme 8. Methylation of perquinoline-substituted HAB **21**.

Theoretically, the salt **25** can exist in eight rotameric forms A-H. Similar to hexa(β -naphthyl)benzene,²³ solving the simple combinatoric problem created by propeller-like compounds such as **21** or **25** results in the following ratio: A:B:C:D:E:F:G:H = 1:6:3:6:6:3:6:1. Figure 2 shows possible interconversions between the forms A-H, which we calculated in form of the hexacationic species (DFT, 6-31G*/PBE0). Schematically, the rotamers can be classified as those in which N-methylquinolinium residues are 6:0 (all-syn), 5-up/1-down, 4 up/2-down (1,2; 1,3; 1,4), and 3-up/3-down (1,2,3; 1,2,4; 1,3,5) (Figure 2). The calculated energy differences between the eight rotamers of **25** are extremely small, and so are the differences



between the rotamers of **21**. The lowest energy rotamer of **25** is the 3-up-3-down (1,3,5) isomer, whereas the all-syn isomer (6,0) has an energy of 7.65 kJ mol^{-1} and is thus the less stable. Similarly, the (6,0) rotamer of the neutral precursor **21** is by only 3.8 kJ mol^{-1} less stable than the (3,3)-1,3,5 rotamer. As a matter of fact, the rotamers of **25** cause signal overlaps in the ^1H NMR spectrum at ambient temperature (Figure 1, spectrum 1, Supplementary Information). The spectra change reversibly on heating of the NMR sample successively from rt to 100°C (Figure 1, spectra 2-9); on cooling, the original spectrum is reconstituted (Figure 1, spectrum 10). Obviously, no other ratio of sets of isochronous rotamers is formed during this temperature experiment. Calculations of all true minimum structures of **25** show that in all rotamers the dihedral angles between the quinoline and the phenyl rings are in the range from 71° to 88° .




Figure 2. Eight energy-minimized rotamers of **25**, showing which can be directly interconverted by rotation of a single substituent. Relative energies (in kJ mol^{-1}) are shown in red boxes.

Figure 3 shows the calculated most stable conformer of **25** and the HOMO/LUMO profiles of the **25** (3,3)-1,3,5 rotamer. In contrast to **22b**,⁴⁶ in which the HOMO is separated from the LUMO, the HOMO (-19.41 eV) as well as LUMO (-15.48 eV) of **25** are located individually on the six N-methylquinolinium-3-yl residues. Considerable parts of the LUMO are located on the nitrogen atoms as well as on the C2/C4 carbon atoms of the quinolinium residues, whereas significant parts of the HOMO are located on their fused benzene rings. The fraction of the HOMO located at C3 of the (3,3)-1,3,5- as well as of the all-syn (6,0) rotamer (c.f. Supplementary Information) is lying in one plane and is orientated to the opposite phases of neighboring C3, similar to the in-plane-aromatic stabilizations due to interactions between the in-plane ethynyl π -orbitals in hexaethynylbenzene.⁴⁷ This may contribute to the stability of **25**. The frontier orbitals of the neutral precursor **21** are shown in the Supporting Information.




Figure 3. Most stable conformation of **25** (calculated) (above). HOMO (left) and LUMO (right) of **25** (3,3)-1,3,5.

Conclusions

Series of propeller-shaped quinolinium-substituted benzenes as well as their electrostatically neutral precursors have been prepared under variation of the substitution site of the quinoline ring (2-yl, 3-yl, 4-yl), covering the range from the monocationic quinolinium-2,3,4,5,6-pentaphenylbenzene to the hexacationic hexakis(1-methylquinolinium-3-yl)benzene. The latter can exist in eight different rotamers, the (3,3)-1,3,5-isomer of which was calculated to be the most stable one. Due to the propeller-shape geometry which suppresses conjugation throughout the entire π -electron system, the frontier orbitals are located in the individual quinolinium rings.

Experimental

All reactions were carried out under an atmosphere of nitrogen in flame or oven-dried glassware. All chemicals were purchased and used without further purification unless otherwise mentioned. Anhydrous solvents were dried according to standard procedures before usage. Melting points are uncorrected and were determined in an apparatus according to Dr. Tottoli (Büchi). The ATR-IR spectra were obtained on a Bruker Alpha in the range of 400 to 4000 cm^{-1} . ^1H NMR spectra were recorded at 600 MHz . ^{13}C NMR spectra were recorded at 150 MHz , with the solvent peak used as the internal reference. Multiplicities are described by using the following abbreviations: s = singlet, d = doublet, t = triplet, q = quartet, and m = multiplet. Spectroscopic atom numberings are shown in the Supporting Information. Signal orientations in DEPT



experiments were described as follows: o = no signal; + = up (CH₃); - = down (CH₂). The mass spectra (ESIMS) were measured with a Varian 320 MS Triple Quad GC/MS/MS (EIMS) or with an Agilent LCMSD series HP 1100 with APIES at fragmentor voltages as indicated. Samples were sprayed from MeOH at 4000 V capillary voltage and fragmentor voltages of 30 V unless otherwise noted. The HRMS spectra were obtained with a Bruker Impact II, a Bruker Daltonik Tesla-Fourier transform-ion cyclotron resonance mass spectrometer, or with a Waters Micromass LCT with the direct inlet. Chromatography: The reactions were traced by thin layer chromatography with silica gel 60 (F254, company MERCK KGAA). For the detection of substances, quenching was used at either 254 nm or 366 nm with a mercury lamp. The preparative column chromatography was conducted through silica gel 60 (230 400 mesh) of the company MERCK KGAA. Yields are not optimized.

Calculations: All density-functional theory (DFT)-calculations were carried out by using the Firefly 8.2.0 QC package,⁴⁸ which is partially based on the GAMESS (US) source code,⁴⁹ running on Linux 2.6.18-238.el5 SMP (x86_64) on five AMD Phenom II X6 1090T processor workstations (Beowulf-cluster) with Infiniband interconnect and parallelized with MPICH 1.2.7p1. MM2 optimized structures were used as starting geometries. Complete geometry optimizations were carried out on the implemented 6-31G* basis set and with the PBE0 density functional. All calculated structures were proven to be true minima by the absence of imaginary frequencies. Partial charges were obtained with NBO 5.9⁵⁰ from the results of the DFT calculations. Orbital plots were obtained using Jmol 14.27.2.

Synthesis.

General procedure of Sonogashira-Hagihara coupling (Procedure 1)

The reactions were carried out under a nitrogen atmosphere. A mixture of 5 mmol of corresponding haloquinoline 1, 1 mol % of Pd(PPh₃)₂Cl₂, and 2 mol % of CuI was suspended in 7 mL of dry NEt₃ with stirring. A sample of the corresponding ethynylbenzene 2 (1.05 equiv) in dry NEt₃ was added dropwise at ambient temperature. The resulting solutions were then stirred at reflux temperature until complete conversion was monitored by TLC. The mixtures were then allowed to cool to rt. The solvents were removed in vacuo. The resulting residues were finally purified by column chromatography (petroleum ether: ethyl acetate) to afford the products.

2-(Phenylethynyl)quinoline (8a). According to Procedure 1, a solution of 0.491 g (3.00 mmol) of 2-chloroquinoline **6a**, 0.042 g (0.06 mmol) of Pd(PPh₃)₂Cl₂, 0.012 g (0.06 mmol) CuI and 0.367 g (3.6 mmol) of ethynylbenzene **7** in 20 mL of anhydrous NEt₃ was heated over the period of 2 h under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 5:1) gave 2-(phenylethynyl)quinoline **8a**. Yield 0.653 g, 95%, a yellow solid, mp 75 °C (72-75 °C⁵¹). ¹H NMR (600 MHz, CDCl₃): δ = 8.44 (d, *J* = 8.3 Hz, 1H, 4-H), 8.03-8.01 (m, 2H, 5-H, 8-H), 7.82 (ddd, *J* = 1.5, 6.7, 8.5 Hz, 1H, 7-H), 7.74 (d, *J*

= 8.3 Hz, 1H, 3-H), 7.70-7.68 (m, 2H, 2'-H, 6'-H), 7.66 (ddd, *J* = 1.2, 6.9, 8.1 Hz, 1H, 6-H), 7.53-7.48 (m, 3H, 3'-H, 4'-H, 5'-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 147.7 (o, C8a), 142.6 (o, C2), 136.7 (+, C4), 131.9 (+, C2', C6'), 129.7 (+, C4'), 128.9 (+, C3', C5'), 128.6 (+, C5), 128.0 (+, C8), 127.4 (+, C6), 126.9 (o, C4a), 124.3 (+, C3), 121.2 (o, C1'), 89.5 (o, Cα), 89.1 (o, Cβ) ppm. IR (ATR): 3053, 2208, 2161, 1616, 1591, 1552, 1498, 1441, 1425, 1310, 1295, 1211, 1142, 1115, 1070, 989, 829, 788, 769, 760, 693, 620, 548, 526, 478 cm⁻¹. HRMS (ESI): *m/z* calcd for C₁₇H₁₂N [M + H]⁺ 230.0965, found 230.0969; *m/z* calcd for C₁₇H₁₁NNa [M + Na]⁺ 252.0789, found 252.0789.

3-(Phenylethynyl)quinoline (8b): According to Procedure 1, a solution of 2.080 g (10.00 mmol) of 3-bromoquinoline **6b**, 0.070 g (0.10 mmol) of Pd(PPh₃)₂Cl₂, 0.038 g (0.20 mmol) CuI and 1.071 g (10.50 mmol) of ethynylbenzene **7** in 30 mL of anhydrous NEt₃ was heated over the period of 1.5 h under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 3:1) gave 3-(phenylethynyl)quinoline **8b**. Yield 2.290 g, 100%, a white solid, mp 83 °C. ¹H NMR (600 MHz, CDCl₃): δ = 9.00 (d, *J* = 2.0 Hz, 1H, 2-H), 8.30 (d, *J* = 2.0 Hz, 1H, 4-H), 8.10 (d, *J* = 8.3 Hz, 1H, 8-H), 7.79 (d, *J* = 8.3 Hz, 1H, 5-H), 7.72 (ddd, *J* = 1.5, 6.9, 8.3 Hz, 1H, 7-H), 7.63-7.53 (m, 3H, 6-H, 2'-H, 6'-H), 7.42-7.35 (m, 3H, 3'-H, 4'-H, 5'-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 152.1 (+, C2), 146.8 (o, C8a), 138.3 (+, C4), 131.8 (+, C2', C6'), 130.1 (+, C7), 129.4 (+, C8), 128.9 (+, C4'), 128.5 (+, C3', C5'), 127.6 (+, C6), 127.3 (+, C5), 127.3 (o, C4a), 122.6 (o, C1'), 117.5 (o, C3), 92.7 (o, Cβ), 86.7 (o, Cα) ppm. IR (ATR): 3008, 2162, 1619, 1599, 1564, 1484, 1443, 1409, 1368, 1294, 1196, 1124, 1113, 1072, 980, 905, 860, 784, 756, 691, 656, 640, 539, 515, 498, 474, 429 cm⁻¹. HRMS (ESI): *m/z* calcd for C₁₇H₁₁NNa [M + Na]⁺ 252.0789, found 252.0796.

4-(Phenylethynyl)quinoline (8c): According to Procedure 1, a solution of 0.624 g (3.00 mmol) of 4-bromoquinoline **6c**, 0.042 g (0.06 mmol) of Pd(PPh₃)₂Cl₂, 0.012 g (0.06 mmol) CuI and 0.367 g (3.6 mmol) of ethynylbenzene **7** in 20 mL of anhydrous NEt₃ was heated over the period of 2 h under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 5:1) gave 4-(phenylethynyl)quinoline **8c**. Yield 0.674 g, 98%, of a yellow solid, mp 45 °C. ¹H NMR (600 MHz, CDCl₃): δ = 8.90 (d, *J* = 4.4 Hz, 1H, 2-H), 8.38 (ddd, *J* = 0.6, 1.4, 8.3 Hz, 1H, 5-H), 8.14 (d, *J* = 8.1 Hz, 1H, 8-H), 7.76 (ddd, *J* = 1.4, 6.9, 8.4 Hz, 1H, 7-H), 7.68-7.66 (m, 2H, 2'-H, 6'-H), 7.64 (ddd, *J* = 1.4, 6.9, 8.2 Hz, 1H, 6-H), 7.57 (d, *J* = 4.4 Hz, 1H, 3-H), 7.44-7.42 (m, 3H, 3'-H, 4'-H, 5'-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 149.9 (+, C2), 148.3 (o, C8a), 132.1 (+, C2', C6'), 130.0 (+, C7, C8), 129.9 (o, C4), 129.5 (+, C4'), 128.7 (+, C3', C5'), 127.9 (o, C4a), 127.3 (+, C6), 126.1 (+, C5), 123.7 (+, C3), 122.4 (o, C1'), 98.8 (o, Cβ), 85.2 (o, Cα) ppm. IR (ATR): 3085, 3051, 3029, 2998, 2212, 1961, 1919, 1595, 1576, 1505, 1487, 1441, 1418, 1390, 1363, 1310, 1277, 1217, 1192, 1175, 1134, 1068, 1029, 919, 871, 846, 815, 756, 687, 641, 579, 550, 529, 506, 485, 442 cm⁻¹. HRMS (ESI): *m/z* calcd for C₁₇H₁₂N [M + H]⁺ 230.0965, found 230.0965.

General procedure of preparation of propeller-like compounds (Procedure 2): Benzophenone (10 g) was melted in a 50 mL round-bottomed flask fitted with an air condenser. Corresponding phenylethynylquinoline **8** (2.00 mmol) and



tetraphenylcyclopentadienone **9** (2.50 mmol) were added to the flask, which was heated for 0.5 h using a heat gun. The solution was cooled to rt and toluene (10 mL) was added to prevent the solidification of the benzophenone. After cooling, *n*-hexane (50 mL) was added, resulting in the precipitation of a product, which was collected by vacuum filtration.

2,3,4,5,6-Pentaphenyl-1-(quinolin-2-yl)benzene (10a):

2,3,4,5,6-pentaphenyl-1-(quinoline-2-yl)benzene **10a** was prepared by Procedure 2 using 2-(phenylethynyl)quinoline **8a** (0.158 g, 0.690 mmol) and tetraphenylcyclopentadienone **9** (0.394 g, 1.035 mmol) in benzophenone (5 g) in 10 mL round-bottomed flask in 0.75 h. Yield 0.260 g, 64%, a white solid, mp >360 °C. ¹H NMR (600 MHz, CDCl₃): δ = 7.75 (d, *J* = 8.6 Hz, 1H, 8ⁱ-H), 7.64 (d, *J* = 8.5 Hz, 1H, 4ⁱ-H), 7.55 (dd, *J* = 1.2, 8.2 Hz, 1H, 5ⁱ-H), 7.51 (ddd, *J* = 1.5, 6.9, 8.5 Hz, 1H, 7ⁱ-H), 7.36 (ddd, *J* = 1.1, 7.0, 8.1 Hz, 1H, 6ⁱ-H), 7.01 (d, *J* = 8.5 Hz, 1H, 3ⁱ-H), 7.00-6.96 (m, 2H, Ph), 6.93-6.83 (m, 16 H, 2ⁱⁱ-H, 6ⁱⁱ-H, 2ⁱⁱⁱ-H, 6ⁱⁱⁱ-H, 2^v-H, 6^v-H, 2^{vi}-H, 6^{vi}-H, 3ⁱⁱ-H, 5ⁱⁱ-H, 3ⁱⁱⁱ-H, 5ⁱⁱⁱ-H, 3^v-H, 5^v-H, 3^{vi}-H, 5^{vi}-H), 6.79-6.72 (m, 7H, Ph) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 160.2 (o, C2ⁱ), 147.1 (o, C8aⁱ), 141.6 (o, C1^{iv}), 140.8/140.4/140.3/140.1 (o, C2, C3, C5, C6, C1ⁱⁱ, C1ⁱⁱⁱ, C1^v, C1^{vi}), 140.6 (o, C4), 140.0 (o, C1), 134.2 (+, C4ⁱ), 131.7/131.5/131.4/130.7 (+, Ph), 129.05 (+, C7ⁱ), 129.00 (+, C8ⁱ), 127.3 (+, C5ⁱ), 126.8 (+, Ph), 126.0 (+, C6ⁱ), 125.7 (o, C4aⁱ), 125.51/125.45 (+, Ph) ppm. IR (ATR): 3055, 3025, 1597, 1558, 1501, 1441, 1403, 1323, 1300, 1221, 1154, 1073, 1029, 948, 839, 812, 753, 718, 695, 617, 583, 555, 531, 478 cm⁻¹. HRMS (ESI): *m/z* calcd for C₄₅H₃₂N [M + H]⁺ 586.2530, found 586.2531.

2,3,4,5,6-Pentaphenyl-1-(quinolin-3-yl)benzene (10b):

2,3,4,5,6-Pentaphenyl-1-(quinoline-3-yl)benzene **10b** was prepared by Procedure 2 using 3-(phenylethynyl)quinoline **8b** (0.458 g, 2.00 mmol) and tetraphenylcyclopentadienone **9** (0.961 g, 2.50 mmol) in benzophenone (10 g) in 50 mL round-bottomed flask in 0.5 h. Yield 0.590 g, 51%, a white solid, mp 334 °C. ¹H NMR (600 MHz, CDCl₃): δ = 8.45 (d, *J* = 2 Hz, 1H, 2ⁱ-H), 7.84 (d, *J* = 8.4 Hz, 1H, 8ⁱ-H), 7.59 (d, *J* = 2 Hz, 1H, 4ⁱ-H), 7.53-7.51 (m, 1H, 7ⁱ-H), 7.43 (d, *J* = 8.0 Hz, 5ⁱ-H), 7.35-7.33 (m, 1H, 6ⁱ-H), 6.88-6.85 (m, 20H, 2ⁱⁱ-H, 6ⁱⁱ-H, 2ⁱⁱⁱ-H, 6ⁱⁱⁱ-H, 2^{iv}-H, 6^{iv}-H, 2^v-H, 6^v-H, 2^{vi}-H, 6^{vi}-H, 3ⁱⁱ-H, 5ⁱⁱ-H, 3ⁱⁱⁱ-H, 5ⁱⁱⁱ-H, 3^{iv}-H, 5^{iv}-H, 3^v-H, 5^v-H, 3^{vi}-H, 5^{vi}-H), 6.83-6.78 (m, 4H, 4ⁱⁱ-H, 4ⁱⁱⁱ-H, 4^v-H, 4^{vi}-H), 6.78-6.77 (m, 1H, 4^{iv}-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 152.9 (+, C2ⁱ), 145.6 (o, C8aⁱ), 141.4 (o, C4), 141.07 (o, C2, C6), 140.88 (o, C3, C5), 140.44 (o, C1^{iv}), 140.34 (o, C1ⁱⁱ, C1^{vi}), 139.96 (o, C1ⁱⁱⁱ, C1^v), 137.8 (+, C4ⁱ), 136.51 (o, C1), 134.2 (o, C3ⁱ), 131.51/131.48/131.45/131.41/131.37 (+, C2ⁱⁱ, C6ⁱⁱ, C2ⁱⁱⁱ, C6ⁱⁱⁱ, C2^{iv}, C6^{iv}, C2^v, C6^v, C2^{vi}, C6^{vi}), 129.0 (+, C8ⁱ), 128.9 (+, C7ⁱ), 127.6 (+, C5ⁱ), 127.4/127.0/126.85/126.83/126.78 (+, C3ⁱⁱ, C5ⁱⁱ, C3ⁱⁱⁱ, C5ⁱⁱⁱ, C3^{iv}, C5^{iv}, C3^v, C5^v, C3^{vi}, C5^{vi}), 127.1 (o, C4aⁱ), 126.2 (+, C6ⁱ), 125.87 (+, C4ⁱⁱ, C4^{vi}), 125.54 (+, C4ⁱⁱⁱ, C4^v), 125.52 (+, C4^{iv}) ppm. IR (ATR): 3055, 3024, 1662, 1600, 1495, 1440, 1401, 1354, 1311, 1275, 1261, 1127, 1070, 1029, 966, 906, 838, 817, 779, 747, 738, 719, 695, 647, 638, 555, 544, 535, 480, 443, 413 cm⁻¹. HRMS (APCI): *m/z* calcd for C₄₅H₃₂N [M + H]⁺ 586.2529, found 586.2530.

2,3,4,5,6-Pentaphenyl-1-(quinolin-4-yl)benzene (10c):

2,3,4,5,6-Pentaphenyl-1-(quinoline-4-yl)benzene **10c** was prepared by Procedure 2 using 4-(phenylethynyl)quinoline **8c**

(0.302 g, 1.319 mmol) and tetraphenylcyclopentadienone **9** (0.430 g, 1.121 mmol, 0.85 equiv) in benzophenone (10 g) in 50 mL round-bottomed flask in 0.5 h (total conversion of dienone). Yield 0.529 g, 81%, a white solid, mp 323 °C. ¹H NMR (600 MHz, CDCl₃): δ = 8.49 (d, *J* = 4.4 Hz, 1H, 2-H), 7.86 (dd, *J* = 0.5, 7.9 Hz, 1H, 5-H), 7.80 (d, *J* = 8.3, 1H, 8-H), 7.50 (ddd, *J* = 1.5, 6.9, 8.4 Hz, 1H, 7-H), 7.39 (ddd, *J* = 1.3, 6.9, 8.3 Hz, 1H, 6-H), 7.05 (d, *J* = 4.4 Hz, 1H, 3-H), 6.93-6.82 (m, 17H, Ph), 6.77-6.75 (m, 2H, Ph), 6.68-6.64 (m, 4H, Ph), 6.55-6.52 (m, 2H, Ph) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 148.8 (+, C2), 147.9 (o, C4), 147.7 (o, C8a), 141.5 (o), 140.74 (o), 140.70 (o), 140.4 (o), 140.2 (o), 139.7 (o), 136.0 (o, C2), 131.53 (+), 131.46 (+), 131.44 (+), 130.9 (+), 130.2 (+), 129.3 (+, C8), 128.8 (+, C7), 128.0 (o, C4a), 127.2 (+, C5), 126.84 (+), 126.81 (+), 126.7 (+), 126.6 (+), 125.8 (+), 125.7 (+, C6), 125.5 (+), 124.4 (+, C3) ppm. IR (ATR): 3052, 3025, 1598, 1496, 1440, 1385, 1291, 1071, 1028, 905, 852, 817, 759, 736, 718, 695, 616, 586, 562, 551, 542, 463, 449, 425 cm⁻¹. HRMS (ESI): *m/z* calcd for C₄₅H₃₂N [M + H]⁺ 586.2530, found 586.2522.

3-((4-Ethynylphenyl)ethynyl)quinoline (14): According to Procedure 1, a solution of 2.080 g (10.00 mmol) of 3-bromoquinoline **6b**, 0.070 g (0.10 mmol) of Pd(PPh₃)₂Cl₂, 0.038 g (0.20 mmol) of CuI and 1.197 g (9.50 mmol) of 1,4-diethynylbenzene **13** in 50 mL of anhydrous NEt₃ was heated (3.5 h) under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 3:1) gave 3-((4-ethynylphenyl)ethynyl)quinoline **14**. Yield 0.889 g, 37%, a brown solid, mp 113 °C. ¹H NMR (600 MHz, CDCl₃): δ = 8.99 (d, *J* = 2.1 Hz, 1H, 2-H), 8.30 (d, *J* = 2.1 Hz, 1H, 4-H), 8.10 (d, *J* = 8.5 Hz, 1H, 8-H), 7.70 (d, *J* = 8.1 Hz, 1H, 5-H), 7.73 (ddd, *J* = 1.5, 7.0, 8.5 Hz, 1H, 7-H), 7.57 (ddd, *J* = 1.5, 7.0, 8.1 Hz, 1H, 6-H), 7.55-7.53 (m, 2H, 2'-H, 6'-H), 7.51-7.49 (m, 2H, 3'-H, 5'-H), 3.20 (s, 1H, CCH) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 152.1 (+, C2), 147.1 (o, C8a), 138.5 (+, C4), 132.3 (+, C3', C5'), 131.7 (+, C2', C6'), 130.4 (+, C7), 129.6 (+, C8), 127.8 (+, C5), 127.5 (+, C6), 127.4 (o, C4a), 123.2 (o, C1'), 122.6 (o, C4'), 117.2 (o, C3), 92.1 (o, Cβ), 88.7 (o, Cα), 83.2 (o, Cγ), 79.4 (o, Cδ) ppm. IR (ATR): 3265, 3060, 3034, 2101, 1969, 1710, 1699, 1602, 1566, 1487, 1404, 1351, 1266, 1145, 1105, 1010, 981, 958, 906, 861, 838, 782, 752, 691, 653, 622, 548, 471, 419 cm⁻¹. HRMS (ESI): *m/z* calcd for C₁₉H₁₂N [M + H]⁺ 254.0964, found 254.0972; *m/z* calcd for C₁₉H₁₁NNa [M + Na]⁺ 276.0789, found 276.0785.

1-(Quinolin-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)-benzene (15):

1-(Quinolin-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene **15** was prepared by Procedure 2 using 3-((4-ethynylphenyl)ethynyl)quinoline **14** (0.290 g, 1.146 mmol) and tetraphenylcyclopentadienone **9** (1.320 g, 3.438 mmol, 3.0 equiv) in benzophenone (10 g) in 50 mL round-bottomed flask in 0.5 h. Yield 0.885 g, 77%, a brown solid, mp 368 °C (decomp.). ¹H NMR (600 MHz, CDCl₃): δ = 8.42 (d, *J* = 2.0 Hz, 1H), 7.90 (d, *J* = 8.3 Hz, 1H), 7.57 (ddd, *J* = 1.4, 6.8, 8.3 Hz, 1H), 7.55 (d, *J* = 2.0 Hz, 1H), 7.46-7.45 (m, 1H), 7.41-7.38 (m, 1H), 7.23 (s, 1H), 7.15-1.13 (m, 2H), 7.07-7.06 (m, 2H), 6.94-6.55 (m, 40H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 152.9 (+), 145.7 (o), 141.8 (o), 141.7 (o), 141.3 (o), 141.1 (o), 140.86 (o), 140.82 (o), 140.80 (o), 140.73 (o), 140.48 (o), 140.43 (o), 140.34 (o), 140.27 (o), 140.21 (o), 140.07 (o), 139.95 (o), 139.7 (o), 139.3 (o), 139.1 (o), 138.8



(o), 137.9 (o), 137.8 (+), 136.5 (o), 134.2 (o), 132.5 (+), 131.6 (+), 131.5 (+), 131.39 (+), 131.35 (+), 131.0 (+), 130.9 (+), 130.2 (+), 130.0 (+), 129.1 (+), 128.89 (+), 128.87 (+), 128.77 (+), 128.4 (+), 127.69 (+), 127.62 (+), 127.3 (+), 127.1 (+), 127.0 (+), 126.96 (o), 126.82 (+), 126.76 (+), 126.66 (+), 126.31 (+), 126.2 (+), 125.8 (+), 125.64 (+), 125.55 (+), 125.50 (+), 125.46 (+), 125.36 (+) ppm. IR (ATR): 3024, 1599, 1490, 1441, 1398, 1275, 1071, 1021, 907, 851, 763, 696, 638, 615, 566, 554, 480, 425 cm⁻¹. HRMS (ESI): *m/z* calcd for C₇₅H₅₁NNa [M + Na]⁺ 988.3914, found 988.3901.

3,3'-Ethyne-1,2-diylquinoline (18a): According to Procedure 1, a solution of 0.208 g (1.00 mmol) of 3-bromoquinoline **6b**, 0.007 g (0.01 mmol) of Pd(PPh₃)₂Cl₂, 0.0038 g (0.02 mmol) CuI and 0.2184 g (1.05 mmol) of 3-ethynylquinoline **17** in 10 mL of anhydrous NEt₃ was heated (2 h) under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 3:1) gave 3,3'-ethyne-1,2-diylquinoline **18a**. Yield 0.196 g, 70%, a brownish solid, mp 173 °C. ¹H NMR (600 MHz, CDCl₃): δ = 9.05 (d, *J* = 2 Hz, 2H, 2-H, 2'-H), 8.36 (d, *J* = 2 Hz, 2H, 4-H, 4'-H), 8.12 (d, *J* = 8.1 Hz, 2H, 8-H, 8'-H), 7.82 (d, *J* = 8.3 Hz, 2H, 5-H, 5'-H), 7.74 (ddd, *J* = 1.4, 6.9, 8.4 Hz, 2H, 7-H, 7'-H) 7.58 (ddd, *J* = 1.1, 6.9, 8.0 Hz, 2H, 6-H, 6'-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 152.0 (+, C2, C2'), 147.2 (o, C8a, C8a'), 138.8 (+, C4, C4'), 130.5 (+, C7, C7'), 129.6 (+, C8, C8'), 127.8 (+, C5, C5'), 127.6 (+, C6, C6'), 127.3 (o, C4a, C4a'), 116.9 (o, C3, C3'), 90.0 (o, Cα) ppm. IR (ATR): 3061, 2169, 1839, 1616, 1565, 1487, 1468, 1413, 1357, 1291, 1226, 1192, 1122, 991, 955, 906, 878, 869, 786, 770, 761, 748, 737, 640, 614, 594, 553, 527, 491, 479, 472, 461, 445, 415 cm⁻¹. MS (ESI): *m/z* = 281.1 [M + H]⁺. HRMS (ESI): *m/z* calcd for C₂₀H₁₃N₂ [M + H]⁺ 281.1074, found 281.1072.

3-(Quinolin-4-ylethynyl)quinoline (18b): According to Procedure 1, a solution of 0.208 g (1.00 mmol) of 4-bromoquinoline **6c**, 0.070 g (0.10 mmol) of Pd(PPh₃)₂Cl₂, 0.038 g (0.20 mmol) CuI and 0.218 g (1.05 mmol) of 3-ethynylquinoline **17** in 10 mL of anhydrous NEt₃ was heated (3.5 h) under reflux temperature. Finally, a purification by column chromatography (petroleum ether: ethyl acetate = 3:1) gave 3-(quinolin-4-ylethynyl)quinoline **18b**. Yield 0.160 g, 57%, a yellow solid, mp 134 °C. ¹H NMR (600 MHz, CDCl₃): δ = 9.11 (d, *J* = 2.1 Hz, 1H, 2-H), 8.94 (d, *J* = 4.5 Hz, 1H, 2'-H), 8.45 (d, *J* = 2.0 Hz, 1H, 4-H), 8.40 (dd, *J* = 0.9, 8.3 Hz, 1H, 5'-H) 8.16 (d, *J* = 7.8, 1H, 8'-H), 8.14 (d, *J* = 7.8 Hz, 1H, 8-H) 7.85 (d, *J* = 7.9 Hz, 1H, 4-H), 7.80–7.76 (m, 2H, 7-H, 7'-H), 7.68 (ddd, *J* = 1.2, 6.9, 8.2 Hz, 1H, 6'-H), 7.63 (d, *J* = 4.3 Hz, 1H, 3'-H), 7.61 (ddd, *J* = 1.2, 7.1, 8.2 Hz, 1H, 6-H) ppm. ¹³C NMR (150 MHz, CDCl₃): δ = 151.9 (+, C2), 149.9 (+, C2'), 148.3 (o, C8a'), 147.5 (o, C8a), 139.2 (+, C4), 130.9 (+, C7), 130.2 (+, C7'), 130.2 (+, C8'), 129.7 (+, C8), 129.1 (o, C4'), 127.9 (+, C5), 127.7 (+, C6), 127.6 (o, C4a'), 127.6 (+, C6'), 127.3 (o, C4a), 126.0 (+, C5'), 123.9 (+, C3'), 116.5 (o, C3), 95.7 (o, Cα), 88.3 (o, Cβ) ppm. IR (ATR): 3033, 2209, 2203, 1615, 1572, 1484, 1419, 1282, 1220, 1192, 1123, 1011, 976, 905, 845, 787, 758, 651, 609, 542, 514, 491, 476, 445 cm⁻¹. MS (ESI): *m/z* = 281.1 [M + H]⁺. HRMS (ESI): *m/z* calcd for C₂₀H₁₃N₂ [M + H]⁺ 281.1074, found 281.1062.

1,2-Di(quinolin-3-yl)-3,4,5,6-tetraphenylbenzene (19a): According to Procedure 2, benzophenone (5 g) was melted in a 50 mL round-bottomed flask fitted with an air condenser. 3,3'-

Ethyne-1,2-diylquinoline **18a** (0.280 g, 1.00 mmol) and tetraphenyl-cyclopentadienone **9** (0.576 g, 1.50 mmol) were added to the flask, which was heated for 1 h using an open flame. The solution was cooled to rt and toluene (3 mL) was added to prevent the solidification of the benzophenone. After cooling, *n*-hexane (200 mL) was added, resulting in the precipitation of **19a** as a white powder, which was washed with *n*-hexane and toluene, and collected by vacuum filtration. Yield 0.570 g, 90%, a white solid, mp 344 °C. ¹H NMR (600 MHz, DMSO-*d*₆), two sets of isochronous rotameric forms (1:0.8): δ = 8.52 (d, *J* = 2.1 Hz, 3.6H, 2-H, 2'-H), 7.97 (d, *J* = 1.7 Hz, 2H, 4-H), 7.93 (d, *J* = 1.7 Hz, 1.6H, 4'-H), 7.64–7.61 (m, 3.6H, 8-H, 8'-H), 7.59 (d, *J* = 7.9 Hz, 2H, 5-H), 7.52 (d, *J* = 8.3 Hz, 1.6H, 5'-H), 7.50–7.47 (m, 3.6H, 7-H, 7'-H), 7.38–7.36 (m, 2H, 6-H), 7.35–7.32 (m, 1.6H, 6'-H), 7.01–6.74 (m, 36H, Ph) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 152.0 (+, C2), 151.8 (+, C2'), 144.99 (o, C8a'), 144.98 (o, C8), 141.06 (o), 141.04 (o), 140.9 (o), 139.5 (o), 139.29 (o), 139.25 (o), 137.4 (+, C4'), 137.0 (+, C4), 136.95 (o, Cα), 136.93 (o, Cβ), 132.97 (o), 132.94 (o), 130.93 (+), 130.89(+), 130.77 (+), 130.72 (+), 130.70 (+), 129.66 (+), 129.51 (+), 129.18 (+, C7 or C7'), 129.16 (+, C7 or C7'), 128.34 (+, C8 or C8'), 128.31 (+, C8 or C8'), 127.8 (+, C5), 127.6 (+, C5'), 127.0 (+), 126.8 (+), 126.7 (+), 126.5 (+, C6, C6'), 126.3 (o, C4a or C4a'), 126.20 (o, C4a or C4a'), 125.9 (+), 125.7 (+) ppm. IR (ATR): 3055, 3025, 1600, 1552, 1490, 1401, 1314, 1241, 1127, 1071, 966, 907, 812, 784, 747, 696, 632, 536, 476 cm⁻¹. HRMS (ESI): *m/z* calcd for C₄₈H₃₃N₂ [M + H]⁺ 637.2639, found 637.2637.

1-(Quinolin-3-yl)-2-(quinolin-4-yl)-3,4,5,6-tetraphenylbenzene (19b): 1-(Quinoline-3-yl)-2-(quinoline-4-yl)-3,4,5,6-tetraphenylbenzene **19b** was prepared by Procedure 2 using 4-(quinolin-3-ylethynyl)quinoline **18b** (0.264 g, 0.943 mmol) and tetraphenylcyclopentadienone **9** (0.543 g, 1.414 mmol) in benzophenone (5 g) in 10 mL round-bottomed flask in 0.5 h. The solution was cooled to rt and toluene (4 mL) was added to prevent the solidification of the benzophenone. Then solvents were evaporated with silica gel and obtained solid was purified by column chromatography with EE:PE (1:1) as eluents to give **19b**. Yield 0.469 g, 78%, a white solid, mp 300 °C (decomp.). ¹H NMR (600 MHz, DMSO-*d*₆): δ = 8.58 (d, *J* = 2.2 Hz, 1H), 8.46–8.44 (m, 2H), 8.25 (d, *J* = 2.2 Hz, 1H), 7.97–7.96 (m, 1H), 7.91–7.87 (m, 2H), 7.61–7.49 (m, 10H), 7.47 (ddd, *J* = 1.6, 6.8, 8.4 Hz, 2H), 7.45 (ddd, *J* = 1.5, 6.6, 8.2 Hz, 1H), 7.41–7.39 (m, 2H), 7.36 (ddd, *J* = 1.2, 6.8, 8.1 Hz, 1H), 7.26–7.24 (m, 1H), 7.21–7.19 (m, 1H), 7.13–7.00 (m, 8H), 6.96–6.81 (m, 22H), 6.79–6.72 (m, 3H), 6.71–6.67 (m, 4H), 6.55 (t, *J* = 7.6 Hz, 2H) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 151.50 (+), 150.81 (+), 148.76 (+), 148.58 (+), 146.85 (o), 146.78 (o), 145.98 (o), 145.89 (o), 145.00 (o), 144.90 (o), 141.21 (o), 141.18 (o), 141.07 (o), 141.01 (o), 140.86 (o), 140.70 (o), 140.29 (o), 140.20 (o), 139.52 (o), 139.45 (o), 139.17 (o), 139.15 (o), 139.05 (o), 138.94 (o), 136.72 (+), 136.35 (o), 136.29 (o), 135.89 (o), 135.82 (o), 135.61 (+), 132.66 (o), 132.55 (o), 131.11 (+), 130.98 (+), 130.94 (+), 130.90 (+), 130.81 (+), 130.76 (+), 130.71 (+), 130.63 (+), 130.60 (+), 130.58 (+), 130.53 (+), 130.42 (+), 129.36 (+), 129.07 (+), 129.01 (+), 128.95 (+), 128.75 (+), 128.68 (+), 128.21 (+), 127.63 (+), 127.31 (+), 126.93 (+), 126.89 (+), 126.81 (+), 126.76 (+), 126.72 (+), 126.67 (+), 126.63 (+), 126.49 (+), 126.45 (+), 126.40 (+), 126.33 (+), 126.24 (+), 125.89



(+), 125.83 (+), 125.73 (+), 125.65 (+), 124.37 (+), 123.82 (+) ppm. IR (ATR): 3055, 3027, 1699, 1600, 1585, 1567, 1505, 1490, 1463, 1442, 1384, 1359, 1126, 1072, 1027, 965, 908, 854, 814, 761, 696, 617, 533, 480 cm⁻¹. HRMS (ESI): *m/z* calcd for C₄₈H₃₃N₂ [M + H]⁺ 637.2639, found 637.2629.

General procedure for the synthesis of substituted benzenes (Procedure 3) 2-(Phenylethynyl)quinoline **8a** (0.21 g, 0.92 mmol) was dissolved in anhydrous dioxane (4 mL) in an oven dried Schlenk flask under a nitrogen atmosphere and the flask was evacuated and filled with nitrogen repeatedly (3×). Then, Co₂(CO)₈ (0.015 g) was added to the flask under a nitrogen atmosphere and the flask was evacuated and filled with nitrogen again (3×). The resulting mixture was refluxed for 14 h and dioxane was evaporated. The resulting residue was dissolved in dichloromethane and filtered through a short pad of silica gel. Evaporation of the solvent afforded a dark colored solid which was purified by column chromatography with PE - EE as an eluent.

1,3,5-Triphenyl-2,4,6-tri(quinolin-2-yl)benzene (20a): Yield 0.021 g, 10%, a white solid, mp 337 °C. ¹H NMR (600 MHz, DMSO-*d*₆): δ = 7.83 (d, *J* = 8.5 Hz, 3H, 4'-H), 7.72 (d, *J* = 8.5 Hz, 3H, 8'-H), 7.67 (d, *J* = 7.9 Hz, 3H, 5'-H), 7.57 (t, *J* = 7.3 Hz, 3H, 7'-H), 7.41 (t, *J* = 7.3 Hz, 3H, 6'-H), 7.22 (d, *J* = 8.5 Hz, 3H, 3'-H), 7.05 (br s, 6H, 2''-H, 6''-H), 6.70 (br s, 6H, 3''-H, 5''-H), 6.63 (t, *J* = 7.3 Hz, 3H, 4''-H) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 158.9 (o, C2'), 146.4 (o, 8a'), 139.9 (o, C2, C4; C6), 139.8 (o, C1, C3, C5), 138.7 (o, C1''), 134.1 (+, C4'), 130.6 (+, C2'', C6''), 129.0 (+, C7'), 128.6 (+, C8'), 127.4 (+, C5'), 126.3 (+, C3'', C5''), 126.1 (+, C6'), 125.8 (+, C4''), 125.4 (o, C4a'), 124.2 (+, C3') ppm. IR (ATR): 3058, 3038, 3007, 1617, 1595, 1557, 1501, 1443, 1424, 1332, 1305, 1253, 1223, 1156, 1139, 1113, 1073, 1031, 959, 906, 838, 753, 698, 619, 587, 515, 476, 431, 423, 412 cm⁻¹. MS (ESI): *m/z* = 688.5 [M + H]⁺. HRMS (ESI): *m/z* calcd for C₅₁H₃₄N₃ [M + H]⁺ 688.2745, found 688.2743.

1,2,4-Triphenyl-3,5,6-tri(quinolin-2-yl)benzene (20b): Yield 0.063 g, 30%, a white solid, mp 318 °C. ¹H NMR (600 MHz, DMSO-*d*₆): δ = 7.85 (d, *J* = 7.5 Hz, 1H), 7.76-7.68 (m, 4H), 7.57-7.53 (m, 5H), 7.45-7.42 (m, 3H), 7.32-7.29 (m, 2H), 7.22-7.20 (m, 3H), 7.04-6.99 (m, 6H), 6.83-6.67 (m, 9H) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 159.1 (o), 159.0 (o), 158.8 (o), 146.4 (o), 146.3 (o), 146.2 (o), 140.5 (o), 140.4 (o), 140.2 (o), 139.8 (o), 139.6 (o), 139.3 (o), 139.0 (o), 138.9 (o), 138.6 (o), 134.1 (+), 133.6 (+), 130.8 (+), 130.3 (br, +), 129.0 (+), 128.8 (+), 128.6 (+), 128.3 (+), 127.4 (+), 127.3 (+), 126.6 (+), 126.5 (+), 126.1 (+), 125.92 (+), 125.85 (+), 125.81 (+), 125.7 (+), 125.3 (o), 125.2 (o), 124.3 (+), 124.2 (+) ppm. IR (ATR): 3056, 3023, 1737, 1597, 1559, 1501, 1441, 1424, 1294, 1160, 1142, 1294, 1160, 1142, 1111, 1072, 1029, 942, 849, 835, 818, 751, 722, 699, 606, 531, 514, 479, 423, 417 cm⁻¹. MS (ESI): *m/z* = 688.3 [M + H]⁺. HRMS (ESI): *m/z* calcd for C₅₁H₃₄N₃ [M + H]⁺ 688.2745, found 688.2739.

Hexakis(quinolin-3-yl)benzene (21): Hexakis(quinoline-3-yl)benzene **11** was prepared by an identical procedure (Procedure 3) using *bis*(quinoline-3-yl)acetylene **18a** (0.295 g, 0.35 mmol) as above. A dark colored solid was purified by column chromatography with EE, then MeOH - CHCl₃ (1 : 3) as eluents. Yield 0.295 g, 85%, a white solid, mp >370 °C. The compound is insoluble in all NMR solvents which we tested. IR

(ATR): 3025, 1619, 1602, 1567, 1489, 1355, 1317, 1260, 1195, 1126, 1110, 1043, 1017, 974, 953, 905, 858, 785, 744, 696, 605, 527, 474, 439 cm⁻¹. HRMS (ESI): *m/z* calcd for C₆₀H₃₆N₆Na [M + Na]⁺ 863.2899, found 863.2896.

General procedure for the preparation of the salts (Procedure 4): Samples of 0.50 mmol of the corresponding quinolines were dissolved in toluene containing 1 drop of nitrobenzene. Then 0.75 mmol of dimethyl sulfate was added with stirring. Thereafter the resulting mixture was stirred under reflux temperature. After completion of the reaction (controlled by TLC), the solution was cooled, the crude product was filtered off, washed with ethyl acetate (3×10 mL), and dried to afford the product.

2,3,4,5,6-Pentaphenyl-1-(1-methylquinolinium-2-yl)benzene methylsulfate (22a): According to Procedure 4, a solution of 0.100 g (0.171 mmol) of 2,3,4,5,6-pentaphenyl-1-(quinoline-2-yl)benzene **10a**, 1 drop of nitrobenzene and 0.06 mL (0.63 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 2 h under reflux temperature to give 2,3,4,5,6-pentaphenyl-1-(1-methylquinolinium-2-yl)benzene methylsulfate **22a**. Yield 0.115 g, 95%, a white solid, mp 190 °C. ¹H NMR (600 MHz, DMSO-*d*₆): δ = 8.88 (d, *J* = 8.5 Hz, 1H, 4ⁱ-H), 8.31 (d, *J* = 9.0 Hz, 1H, 8ⁱ-H), 8.24-8.23 (m, 2H, 3ⁱ-H, 5ⁱ-H), 8.14 (t, *J* = 8.0 Hz, 7ⁱ-H), 7.93 (t, *J* = 7.4 Hz, 6ⁱ-H), 7.10-6.84 (m, 25H, Ph), 4.41 (s, 3H, N⁺CH₃), 3.38 (s, 3H, CH₃SO₄) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 159.1 (o, C2ⁱ), 144.2 (+, C4ⁱ), 143.9 (o, C1), 141.2 (o), 139.1 (o), 138.9 (o), 138.3 (o), 137.5 (o, C8aⁱ), 137.4 (o), 136.0 (+, C7ⁱ), 131.7 (+), 130.9 (+), 130.6 (+), 130.4 (+), 130.4 (+, C6ⁱ), 130.3 (+, C5ⁱ), 130.26 (+), 130.0 (+), 129.3 (+), 126.7 (+), 127.43 (o, C4aⁱ), 127.39 (+, C3ⁱ), 127.2 (+), 126.95 (+), 126.91 (+), 126.86 (+), 126.2 (+), 126.0 (+), 119.1 (+, C8ⁱ), 52.8 (+, CH₃SO₄), 42.5 (+, N⁺CH₃) ppm. IR (ATR): 3056, 3023, 1619, 1600, 1577, 1519, 1497, 1442, 1409, 1347, 1252, 1224, 1176, 1152, 1073, 1059, 1013, 918, 849, 817, 770, 756, 723, 698, 644, 608, 576, 554, 532, 428 cm⁻¹. HRMS (ESI): *m/z* calcd for C₄₆H₃₄N [M]⁺ 600.2686, found 600.2692.

2,3,4,5,6-Pentaphenyl-1-(1-methylquinolinium-3-yl)benzene methylsulfate (22b): According to Procedure 4, a solution of 0.100 g (0.171 mmol) of 2,3,4,5,6-pentaphenyl-1-(quinoline-3-yl)benzene **10b**, 1 drop of nitrobenzene and 0.06 mL (0.63 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 2 h under reflux temperature to give 2,3,4,5,6-pentaphenyl-1-(1-methylquinolinium-3-yl)benzene methylsulfate **22b**. Yield 0.122 g, 99%, a light green solid, mp 321 °C (decomp.). ¹H NMR (600 MHz, DMSO-*d*₆): δ = 9.35 (d, *J* = 1.4 Hz, 1H, 2-H), 8.84 (s, 1H, 4-H), 8.24 (d, *J* = 8.1 Hz, 1H, 8-H), 8.14 (ddd, *J* = 1.5, 7.1, 8.8 Hz, 1H, 7-H), 8.06 (dd, *J* = 1.1, 8.4 Hz, 1H, 5-H), 7.92-7.90 (m, 1H, 6-H), 7.05 (d, *J* = 7.7 Hz, 2H, Ph), 7.02 (d, *J* = 7.7 Hz, 2H, Ph), 6.96-6.82 (m, 21H, Ph), 4.34 (s, 3H, NCH₃), 3.37 (s, 3H, CH₃SO₄) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 150.6 (+, C2), 147.5 (+, C4), 142.3 (o), 140.7 (o), 140.5 (o), 139.3 (o), 139.0 (o), 138.4 (o), 135.53 (o, C8a), 135.48 (+, C7), 134.4 (o, C3), 132.8 (o, C1'), 131.0 (+), 130.7 (+), 130.64 (+), 130.60 (+), 130.54 (+), 130.5 (+, C6), 129.9 (+, C5), 127.7 (o, C4a), 127.4 (o), 126.89 (+), 126.87 (+), 126.80 (+), 126.77 (+), 126.4 (+), 125.92 (+), 125.85 (+), 118.9 (+, C8), 52.8 (+, CH₃SO₄), 44.8 (+, NCH₃) ppm. IR (ATR): 3051, 3023, 2961, 1600, 1524, 1496, 1442, 1378,



1254, 1215, 1063, 1012, 760, 697, 618, 577, 558, 441 cm^{-1} . HRMS (ESI): m/z calcd for $\text{C}_{46}\text{H}_{34}\text{N} [\text{M}]^+$ 600.2686, found 600.2687.

2,3,4,5,6-Pentaphenyl-1-(1-methylquinolinium-4-yl)benzene methylsulfate (22c): According to Procedure 4, a solution of 0.154 g (0.263 mmol) of 2,3,4,5,6 pentaphenyl-1-(quinoline-4-yl)benzene **10c**, 1 drop of nitrobenzene and 0.06 mL (0.63 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 2 h under reflux temperature to give 2,3,4,5,6-pentaphenyl-1-(1-methylquinolinium-4-yl)benzene methylsulfate **22c**. Yield 0.185 g, 99%, a brownish solid, mp 355 °C (decomp.). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ = 9.14 (d, J = 6.0 Hz, 1H, 2-H), 8.27 (dd, J = 1.0, 8.4 Hz, 1H, 5-H), 8.20 (d, J = 8.9 Hz, 1H, 8-H), 8.15 (d, J = 6.0 Hz, 1H, 3-H), 8.11 (ddd, J = 1.4, 7.1, 8.7 Hz, 1H, 7-H), 7.96-7.94 (m, 1H, 6-H), 7.06 (d, J = 7.7 Hz, 2H, Ph), 7.01 (d, J = 7.7 Hz, 2H, Ph), 6.95-6.83 (m, 14H, Ph), 6.74-6.71 (m, 4H, Ph), 6.60-6.57 (m, 2H, Ph), 4.42 (s, 3H, NCH_3), 3.37 (s, 3H, CH_3SO_4) ppm. ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$): δ = 158.5 (o, C4), 147.5 (+, C2), 142.3 (o), 140.6 (o), 139.36 (o), 139.34 (o), 138.9 (o), 138.3 (o), 137.1 (o, C8a), 135.2 (+, C7), 133.1 (o, C α), 131.0 (+), 130.8 (+), 130.52 (+), 130.49 (+), 129.83 (+, C6), 128.80 (+), 129.3 (+, C5), 128.3 (o, C4a), 126.83 (+), 126.78 (+), 126.74 (+), 126.71 (+), 126.67 (+), 126.4 (+), 125.81 (+), 125.77 (+), 125.3 (+, C3), 118.8 (+, C8), 52.8 (+, CH_3SO_4), 45.0 (+, NCH_3) ppm. IR (ATR): 3051, 3025, 1622, 1615, 1599, 1586, 1577, 1531, 1495, 1441, 1401, 1389, 1375, 1335, 1265, 1216, 1118, 1059, 1015, 869, 820, 766, 726, 695, 609, 576, 553, 542, 464, 430 cm^{-1} . HRMS (ESI): m/z calcd for $\text{C}_{46}\text{H}_{34}\text{N} [\text{M}]^+$ 600.2686, found 600.2673.

1-(1-Methylquinolinium-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene methylsulfate (23): According to Procedure 4, a solution of 0.481 g (0.5 mmol) of 1-(quinolin-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene **15**, 1 drop of nitrobenzene and 0.06 mL (0.63 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 2 h under reflux temperature to give 1-(1-methylquinolinium-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene methylsulfate **23**. Yield 0.521 g, 96%, a khaki solid, mp 222 °C (decomp.). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ = 9.28 (d, J = 1.3 Hz, 1H, Q), 8.73 (s, 1H, Q), 8.35 (d, J = 6.7 Hz, 1H, Q), 8.20 (ddd, J = 1.6, 7.0, 8.7 Hz, 1H, Q), 8.05-8.04 (m, 1H, Q), 7.98-7.95 (m, 1H, Q), 7.26-7.23 (m, 1H, Ph), 7.19-7.13 (m, 4H, Ph), 7.02-6.99 (m, 3H, Ph), 6.95-6.74 (m, 28H, Ph), 6.71-6.64 (m, 5H, Ph), 6.61-6.53 (m, 3H, Ph), 6.49-6.47 (m, 1H, Ph), 4.33 (s, 3H, NCH_3), 3.36 (s, 3H, CH_3SO_4) ppm. ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$): δ = 150.6 (+), 147.5 (+), 142.2 (o), 141.5 (o), 140.9 (o), 140.7 (o), 140.6 (o), 140.4 (o), 140.3 (o), 140.0 (o), 139.6 (o), 139.3 (o), 139.01 (o), 139.00 (o), 138.96 (o), 138.8 (o), 138.6 (o), 138.4 (o), 137.3 (o), 136.5 (o), 135.6 (o), 135.5 (+), 134.4 (o), 132.8 (o), 131.0 (+), 130.9 (+), 130.8 (+), 130.7 (+), 130.64 (+), 130.60 (+), 130.55 (+), 130.53 (+), 130.44 (+), 130.36 (+), 130.28 (+), 130.0 (+), 129.4 (+), 128.9 (+), 128.6 (+), 128.4 (+), 128.2 (+), 127.73 (+), 127.66 (+), 127.4 (+), 126.85 (+), 126.81 (+), 126.51 (+), 126.46 (+), 126.4 (+), 125.91 (+), 125.85 (+), 125.8 (+), 125.44 (+), 125.37 (+), 125.31 (+), 118.9 (+), 52.7 (+), 44.8 (+) ppm. IR (ATR): 3055, 3023, 1600, 1524, 1442, 1378, 1249, 1222, 1178, 1157, 1138, 1058, 1008, 911, 852, 797, 766,

697, 565, 496, 432 cm^{-1} . HRMS (ESI): m/z calcd for $\text{C}_{76}\text{H}_{54}\text{N} [\text{M}]^+$ 980.4251, found 980.4257. DOI: 10.1039/C9OB00357F

1-(1-Methylquinolinium-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene hexafluorophosphate (23PF₆): A suspension of 0.050 g (0.046 mmol) of 1-(1-methylquinolinium-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene methylsulfate **23** and 0.008 g (0.049 mmol) of NH_4PF_6 in 4 mL of water was stirred for 1 day at rt to give 1-(1-methylquinolinium-3-yl)-2,3,4,5-tetraphenyl-6-(4-(1,2,3,4-tetraphenyl)phenyl)benzene hexafluorophosphate **23PF₆**. Yield 0.049 g, 95%, a yellow solid, mp 220 °C (decomp.). ^1H NMR (600 MHz, $\text{DMSO}-d_6$): δ = 9.28 (s, 1H, Q), 8.73 (s, 1H, Q), 8.35 (d, J = 8.9 Hz, 1H, Q), 8.20 (t, J = 7.7 Hz, 1H, Q), 8.05 (d, J = 7.9 Hz, 1H, Q), 7.96 (t, J = 7.5 Hz, 1H, Q), 7.26 (m, 1H, Ph), 7.17-7.14 (m, 4H, Ph), 7.02-6.47 (m, 40H, Ph), 4.33 (s, 3H, NCH_3) ppm. ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$): δ = 150.6 (+), 147.5 (+), 142.2 (o), 141.5 (o), 140.9 (o), 140.7 (o), 140.6 (o), 140.4 (o), 140.3 (o), 140.0 (o), 139.6 (o), 139.3 (o), 139.01 (o), 139.00 (o), 138.96 (o), 138.8 (o), 138.6 (o), 138.4 (o), 137.3 (o), 136.5 (o), 135.6 (o), 135.5 (+), 134.4 (o), 132.8 (o), 131.0 (+), 130.9 (+), 130.8 (+), 130.7 (+), 130.64 (+), 130.60 (+), 130.55 (+), 130.53 (+), 130.44 (+), 130.36 (+), 130.28 (+), 130.0 (+), 129.4 (+), 128.9 (+), 128.6 (+), 128.4 (+), 128.2 (+), 127.73 (+), 127.66 (+), 127.4 (+), 126.85 (+), 126.81 (+), 126.51 (+), 126.46 (+), 126.4 (+), 125.91 (+), 125.85 (+), 125.8 (+), 125.44 (+), 125.37 (+), 125.31 (+), 118.9 (+), 44.8 (+) ppm. IR (ATR): 3055, 3027, 1600, 1521, 1495, 1442, 1379, 1072, 1028, 835, 766, 697, 557, 427 cm^{-1} . HRMS (ESI): m/z calcd for $\text{C}_{76}\text{H}_{54}\text{N} [\text{M}]^+$ 980.4251, found 980.4238.

1,2-Di(1-methylquinolinium-3-yl)-3,4,5,6-tetraphenylbenzene dihexafluorophosphate (24a): A solution of 0.050 g (0.079 mmol) of 1,2-di(quinoline-3-yl)-3,4,5,6-tetraphenylbenzene **19a**, 1 drop of nitrobenzene and 0.02 mL (0.21 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 3 h under reflux temperature, cooled to rt and extracted with water (3 \times 5 mL) and precipitated with excess of NH_4PF_6 (1.3 equiv) to give 1,2-di(1-methylquinolinium-3-yl)-3,4,5,6-tetraphenylbenzene dihexafluorophosphate **24a**. Yield 0.068 g, 90%, a white solid, mp 240 °C. ^1H NMR (600 MHz, $\text{DMSO}-d_6$): two sets of isochronous rotameric forms (0.9:1): δ = 9.43 (d, J = 1.3 Hz, 0.9H, 2-H), 9.40 (d, J = 1.3 Hz, 1H, 2'-H), 9.05 (s, 1H, 4'-H), 8.88 (s, 0.9H, 4-H), 8.30 (d, J = 8.8 Hz, 1H, 8'-H), 8.27 (d, J = 8.9 Hz, 0.9H, 8'-H), 8.16-8.09 (m, 3.8H, 5-H, 5'-H, 7-H, 7'-H), 7.93-7.90 (m, 1.9H, 6-H, 6'-H), 7.05-6.84 (m, 48H, Ph), 4.38 (s, 6H, NCH_3), 4.32 (s, 5.4H, NCH_3) ppm. ^{13}C NMR (150 MHz, $\text{DMSO}-d_6$): δ = 150.6 (+, C2), 149.8 (+, C2'), 147.9 (+, C4'), 147.6 (+, C4), 142.8 (o), 142.7 (o), 141.9 (o), 141.8 (o), 138.5 (o), 137.6 (o), 136.5 (o, C8a'), 136.4 (o, C8a), 135.94 (+, C7 or C7'), 135.85 (+, C7 or C7'), 133.10 (o, C β), 132.99 (o, C α), 132.4 (o), 132.2 (o), 130.89 (+), 130.75 (+), 130.69 (+), 130.60 (+), 130.48 (+), 130.44 (+), 130.42 (+), 130.38 (+), 130.33 (+), 130.31 (+), 130.22 (+, C5), 130.07 (+, C5'), 127.92 (o, C4a'), 127.89 (o, C4a), 127.6 (+), 127.0 (+), 126.7 (+), 126.3 (+), 119.00 (+, C8'), 118.90 (+, C8), 45.4 (+, NCH_3), 45.2 (+, NCH_3) ppm. IR (ATR): 3057, 3027, 1631, 1602, 1583, 1523, 1497, 1443, 1380, 1356, 1334, 1230, 1174, 1141, 1115, 1073, 1024, 936, 830, 769, 744, 702, 619, 556, 524, 504,



419 cm⁻¹. HRMS (ESI): *m/z* calcd for C₅₀H₃₈N₂ [M]²⁺ 333.1512, found 333.1518.

1-(1-methylquinolinium-3-yl)-2-(1-methylquinolinium-4-yl)-3,4,5,6-tetraphenylbenzene dihexafluorophosphate (24b): A solution of 0.050 g (0.079 mmol) of 1-(quinoline-3-yl)-2-(quinoline-4-yl)-3,4,5,6-tetraphenylbenzene **19b**, 1 drop of nitrobenzene and 0.02 mL (0.21 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 3 h under reflux temperature, cooled to rt and extracted with water (3×5 mL) and then precipitated with excess of NH₄PF₆ (1.3 equiv) to give 1-(1-methylquinolinium-3-yl)-2-(1-methylquinolinium-4-yl)-3,4,5,6-tetraphenylbenzene dihexafluorophosphate **24b**. Yield 0.067 g, 89%, a yellow solid, mp 245 °C (decomp.). ¹H NMR (600 MHz, DMSO-*d*₆), two sets of isochronous rotameric forms (24b*/24b# = 1 : 1.27) with close chemical shifts, exact description of the coupling constants is not possible: δ = 9.54[#] (d, *J* = 1.3 Hz, 1H, 2-H), 9.23* (d, *J* = 1.5 Hz, 1H, 2-H), 9.20[#] (d, *J* = 6.3 Hz, 1H, 2'-H), 9.18* (d, *J* = 6.4 Hz, 1H, 2'-H), 9.02* (d, *J* = 1.5 Hz, 1H, 4-H), 8.81[#] (s, 1H, 4-H), 8.34[#] (dd, *J* = 1.1, 8.6 Hz, 1H), 8.26-8.20 (m, 6H), 8.16[#] (d, *J* = 6.3 Hz, 1H, 3'-H), 8.15-8.07 (m, 7H), 8.04-8.01[#] (m, 2H), 7.92* (ddd, *J* = 1.2, 6.8, 8.1 Hz, 1H), 7.85-7.79[#] (m, 2H), 7.21-6.62*[#] (m, 40H, Ph), 4.37[#] (s, 3H, N'CH₃), 4.346[#] (s, 3H, NCH₃), 4.343* (s, 3H, N'CH₃), 4.10* (s, 3H, NCH₃) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 155.64*[#] (o, C4'), 155.59*[#] (o, C4'), 149.53[#] (+, C2), 149.31* (+, C2), 148.47[#] (+, C2'), 148.42* (+, C2'), 147.14* (+, C4), 146.81[#] (+, C4), 142.92 (o), 142.90 (o), 142.89 (o), 142.84 (o), 141.63 (o), 141.49 (o), 140.37 (o), 140.21 (o), 138.56 (o), 138.48 (o), 138.35 (o), 138.31 (o), 137.54 (o), 137.52 (o), 137.48 (o), 137.43 (o), 136.38[#] (o, C8a), 136.10* (o, C8a), 135.96 (+), 135.88 (+), 135.46 (+), 135.38 (+), 133.36* (o, Cβ), 133.23[#] (o, Cβ), 132.26 (o), 132.21 (o), 132.05 (o), 131.97 (o), 131.16 (+), 130.88 (+), 130.73 (+), 130.63 (+), 130.61 (+), 130.58 (+), 130.55 (+), 130.51 (+), 130.42 (+), 130.30 (+), 130.26 (+), 130.23 (+), 130.22 (+), 129.71 (+), 129.63 (+), 129.50 (+), 128.60 (+), 128.32 (+), 127.88 (o), 127.71 (o), 127.61 (+), 127.59 (+), 127.44 (+), 127.37 (+), 127.15 (+), 127.07 (+), 127.03 (+), 126.98 (+), 126.93 (+), 126.85 (+), 126.79 (+), 126.64 (+), 126.30 (+), 126.26 (+), 126.24 (+), 125.24 (+), 124.78 (+), 119.34 (+), 119.02 (+), 118.86 (+), 45.37 (+), 45.33 (+), 45.30 (+), 44.67* (+, NCH₃) ppm. IR (ATR): 3057, 1619, 1605, 1588, 1529, 1498, 1443, 1234, 1175, 1115, 1073, 1025, 1001, 918, 827, 765, 702, 619, 556, 492, 434 cm⁻¹. HRMS (ESI): *m/z* calcd for C₅₀H₃₈N₂ [M]²⁺ 333.1512, found 333.1512.

Hexakis(1-methylquinolinium-3-yl)benzene hexakis hexafluorophosphate (25): According to Procedure 4, a solution of 0.084 g (0.10 mmol) of hexakis(quinoline-3-yl)benzene **21**, 1 drop of nitrobenzene and 0.10 mL (1.00 mmol) of dimethyl sulfate in 5 mL of anhydrous toluene was heated for 3 h under reflux temperature. Then the obtained salt was dissolved in water and precipitated with 1.05 equiv of NH₄PF₆ to give hexakis(1-methylquinolinium-3-yl)benzene hexakis-hexafluorophosphate **25**. Yield 0.171 g, 95%, a white solid, mp 195 °C (decomp.). ¹H NMR (600 MHz, DMSO-*d*₆): δ = 9.41-8.89 (m, 12H), 8.34-8.16 (m, 18H), 8.00-7.92 (m, 6H), 4.44-4.28 (m, 18H) ppm. ¹³C NMR (150 MHz, DMSO-*d*₆): δ = 148.7, 148.2, 137.9, 137.8, 137.7, 137.5, 137.0, 131.0, 130.4, 128.6, 128.0, 119.4, 45.97, 45.89, 45.85, 45.79 ppm. IR (ATR): 3084, 1631,

1582, 1524, 1451, 1382, 1228, 1173, 1040, 952, 920, 827, 771, 753, 740, 614, 556, 494, 440, 413 cm⁻¹. DOI: 10.1039/C9OB00357F

- 1 F. J. Schmitz, K. H. Hollenbeak, D. C. Campbell, *J. Org. Chem.*, 1978, **43**, 3916.
- 2 (a) N. Fusetani, N. Asai, S. Matsunaga, *Tetrahedron Lett.*, 1994, **23**, 3967. (b) H. Anan, N. Seki, O. Noshiro, K. Hona, K. Yasumuro, T. Ozasa, N. Fusetani, *Tetrahedron*, 1996, **52**, 10849.
- 3 S. Albrizio, P. Ciminiello, E. Fattorusso, S. Magno, J. R. Pawlik, *J. Nat. Prod.*, 1995, **58**, 647.
- 4 C. A. Volk, M. Köck, *Org. Lett.*, 2003, **5**, 3567.
- 5 C. A. Volk, M. Köck, *Org. Biomol. Chem.*, 2004, **2**, 1827.
- 6 R. Laville, O. P. Thomas, F. Berrue, F. Reyes, P. Amade, *Eur. J. Org. Chem.*, 2008, 121.
- 7 (a) K. A. Smith, A. Streitwieser, Jr., *J. Org. Chem.*, 1983, **48**, 2629. (b) K. C. Waterman, A. Streitwieser, Jr., *J. Am. Chem. Soc.*, 1984, **106**, 3874. (c) K. A. Smith, K. C. Waterman, A. Streitwieser, Jr., *J. Org. Chem.*, 1985, **50**, 3360. (d) K. C. Waterman, D. V. Speer, A. Streitwieser, Jr., G. C. Look, K. O. Nguyen, J. G. Stack, *J. Org. Chem.*, 1988, **53**, 583. (e) A. S. Koch, K. C. Waterman, K. Banks, A. Streitwieser, *J. Org. Chem.*, 1990, **55**, 6166. (f) A. S. Feng, D. V. Speer, S. G. DiMaggio, M. S. Konings, A. Streitwieser, *J. Org. Chem.*, 1992, **57**, 2902.
- 8 A. S. Koch, A. S. Feng, T. A. Hopkins, A. Streitwieser, *J. Org. Chem.*, 1993, **58**, 1409.
- 9 F. Pühlhofer and R. Weiss, *Eur. J. Org. Chem.*, 2004, 1002.
- 10 (a) R. Weiss, R. Roth, R. H. Lowack, H. Bremer, *Angew. Chem.*, 1990, **102**, 1164; *Angew. Chem., Int. Ed. Engl.*, 1990, **29**, 1132. (b) R. Weiss, B. Pomrehn, F. Hampel, W. Bauer, *Angew. Chem.*, 1995, **107**, 1446; *Angew. Chem., Int. Ed. Engl.*, 1995, **34**, 1319. (c) R. Weiss, R. May, B. Pomrehn, *Angew. Chem.*, 1996, **108**, 1319; *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1232. (d) R. Weiss, S. M. Huber, F. G. Pühlhofer, *Eur. J. Org. Chem.*, 2005, 3530. (e) R. Weiss, S. M. Huber, F. W. Heinemann, P. Audebert, F. G. Pühlhofer, *Angew. Chem.*, 2006, **118**, 8228; *Angew. Chem. Int. Ed.*, 2006, **45**, 8059. (f) R. Weiss, F. G. Pühlhofer, *J. Am. Chem. Soc.*, 2007, **129**, 547.
- 11 (a) A. Schmidt, T. Mordhorst, M. Nieger, *Synthesis*, 2006, 3987. (b) A. Schmidt, J. C. Namyslo, T. Mordhorst, *Tetrahedron*, 2006, **62**, 6893. (c) A. Schmidt, T. Mordhorst, M. Nieger, *Tetrahedron*, 2006, **62**, 1667. (d) A. Schmidt, T. Mordhorst, *Synthesis*, 2005, 781. (e) A. Schmidt, *J. Heterocycl. Chem.*, 2002, **39**, 949.
- 12 (a) A. Schmidt, A. Hetzheim, *Tetrahedron*, 1997, **53**, 1295. (b) A. Schmidt, M. K. Kindermann, *J. Org. Chem.*, 1998, **63**, 4636. (c) A. Schmidt, M. Nieger, *J. Chem. Soc., Perkin Trans. 1*, 1999, 1325.
- 13 A. Schmidt, T. Mordhorst, *Heterocycles*, 2006, **68**, 1393.
- 14 (a) A. Schmidt, T. Mordhorst, M. Nieger, *Org. Biomol. Chem.*, 2005, **3**, 3788. (b) A. Schmidt, T. Mordhorst, T. Habeck, *Org. Lett.*, 2002, **4**, 1375.
- 15 (a) S. G. DiMaggio, K. C. Waterman, D. V. Speer, A. Streitwieser, *J. Am. Chem. Soc.*, 1991, **113**, 4679. (b) A. S. Koch, A. S. Feng, T. A. Hopkins, A. Streitwieser, *J. Org. Chem.*, 1993, **58**, 1409.
- 16 (a) K. Akiyama, S. Tero-Kubota, Y. Ikegami, *J. Am. Chem. Soc.*, 1983, **105**, 3601. (b) Y. Ikegami, T. Muramatsu, K. Hanaya, *J. Am. Chem. Soc.*, 1989, **111**, 5782. (c) K. Okada, K. Matsumoto, M. Oda, H. Murai, K. Akiyama, Y. Ikegami, *Tetrahedron Lett.*, 1995, **36**, 6689.



- 17 R. Weiss, R. May, B. Pohmrehn, *Angew. Chem.*, 1996, **108**, 1319; *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1232.
- 18 (a) W. R. Boon, *Chem. Ind. (London)*, 1965, 782. (b) A. L. Black, L. A. Summers, *J. Chem. Soc. C*, 1969, 610. (c) E. C. Campbell, E. E. Glover, G. Trenholm, *J. Chem. Soc.*, 1969, 1987. (d) A. D. Dodge, *Endeavour*, 1970, **111**, 130.
- 19 (a) C. N. Corder, J. L. Way, *J. Med. Chem.*, 1966, **9**, 638. (b) Y. Ashani, S. Cohen, *J. Med. Chem.*, 1971, **14**, 621. (c) C. F. Barfknecht, F. W. Benz, J. P. Long, *J. Med. Chem.*, 1971, **14**, 1003.
- 20 M. Komloova, K. Musilek, A. Horova, O. Holas, V. Dohnal, F. Gunn-Moore, K. Kuca, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 2505.
- 21 J. Hiller, M. Liu, A. Schmidt, *Heterocycles*, 2017, **94**, 821.
- 22 M. Drev, U. Grošelj, B. Ledinek, F. Perdih, J. Svete, B. Štefane, F. Požgan, *Org. Lett.*, 2018, **20**, 5268.
- 23 L. E. Harrington, J. F. Britten, K. Nikitin, M. J. McGlinchey, *ChemPlusChem*, 2017, **82**, 433.
- 24 Y. Geng, A. Fechtenkötter, K. Müllen, *J. Mater. Chem.*, 2001, **11**, 1634.
- 25 K. Kobayashi, A. Sato, S. Sakamoto, K. Yamaguchi, *J. Am. Chem. Soc.*, 2003, **125**, 3035.
- 26 S. Hiraoka, T. Nakamura, M. Shiro, M. Shionoya, *J. Am. Chem. Soc.*, 2010, **132**, 13223.
- 27 Z. Tomović, J. van Dongen, S. J. George, H. Xu, W. Pisula, P. Leclère, M. M. J. Smulders, S. De Feyter, E. W. Meijer, A. P. H. J. Schenning, *J. Am. Chem. Soc.*, 2007, **129**, 16190.
- 28 S. Hiraoka, Y. Hisanaga, M. Shiro, M. Shionoya, *Angew. Chem.*, 2010, **122**, 1713; *Angew. Chem., Int. Ed.*, 2010, **49**, 1669.
- 29 B. Traber, J. J. Wolff, F. Rominger, T. Oeser, R. Gleiter, M. Goebel, R. Wortmann, *Chem. Eur. J.*, 2004, **10**, 1227.
- 30 R. Shukla, S. V. Lindeman, R. Rathore, *J. Am. Chem. Soc.*, 2006, **128**, 5328.
- 31 M. Steeger, C. Lambert, *Chem. Eur. J.* 2012, **18**, 11937.
- 32 Y. Tanaka, T. Koike, M. Akita, *Chem. Commun.*, 2010, **46**, 4529.
- 33 S. H. Shin, N. Y. Sim, Korean Kongkae Taeho Kongbo (2016), KR 2016072868 A 20160624.
- 34 (a) D. Gust, *J. Am. Chem. Soc.*, 1977, **99**, 6980. (b) J. C. J. Bart, *Acta Crystallogr., Sect. B*, 1968, **24**, 1277. (c) A. Almenningsen, O. Bastiansen, P. N. Skancke, *Acta Chem. Scand.*, 1958, **12**, 1215.
- 35 G. Hilt, T. Vogler, W. Hess, F. Galbiati, *Chem. Commun.*, 2005, **11**, 1474.
- 36 (a) M. Mazik, D. Bläser, R. Boese, *Tetrahedron Lett.*, 2000, **41**, 5827. (b) R. Taylor, O. Kennard, *J. Am. Chem. Soc.*, 1982, **104**, 5063. (c) F. A. Cotton, L. M. Daniels, G. T. Jordan IV, C. A. Murillo, *Chem. Commun.*, 1997, 1673. (d) G. R. Desiraju, *Crystal Engineering. The Design of Organic Solids*; Elsevier: Amsterdam, 1989; pp. 166.
- 37 (a) K. Ghosh, A. J. Bhattacharya, *Indian J. Chem. B*, 1977, **15**, 678. (b) P. K. Banerjee, S. R. Samanta, A. J. Bhattacharya, *Curr. Sci.*, 1976, **45**, 254.
- 38 E. Gagnon, T. Maris, P.-M. Arseneault, K. E. Maly, J. D. Wuest, *Cryst. Growth Des.*, 2010, **10**, 648.
- 39 E. Gagnon, S. D. Halperin, V. Métivaud, K. E. Maly, J. D. Wuest, *J. Org. Chem.*, 2010, **75**, 399.
- 40 D. Gust, A. Patton, *J. Am. Chem. Soc.*, 1978, **100**, 8175.
- 41 (a) M. A. Ogliaruso, M. G. Romanelli, E. I. Becker, *Chem. Rev.*, 1965, **65**, 261. (b) S. Saito, Y. Yamamoto, *Chem. Rev.*, 2000, **10**, 2901.
- 42 (a) J. H. Magill, A. R. Ubbelohde, *Trans. Faraday Soc.*, 1958, **54**, 1811. (b) D. J. Plazek, J. H. Magill, *J. Chem. Phys.*, 1966, **45**, 3038.
- 43 D. Wasserfallen, G. Mattersteig, V. Enkelmann, K. Müllen, *Tetrahedron*, 2006, **62**, 5417.
- 44 H. D. Katz, *J. Org. Chem.*, 1987, **52**, 3932.
- 45 (a) A. Schmidt, S. Batsyts, A. Smevanov, T. Freese, E. G. Hübner, M. Nieger, *J. Org. Chem.*, 2016, **81**, 4202. (b) S. Batsyts, F. J. Ramirez, J. Casado, J. C. Namyslo, A. Schmidt, *Z. Naturforsch.*, 2018, **73**, 481. (c) A. Dreger, R. Cisneros Camuña, N. Münster, T. A. Rokob, I. Pápai, A. Schmidt, *Eur. J. Org. Chem.*, 2010, 4296. (d) M. Liu, M. Nieger, E. Hübner, A. Schmidt, *Chem. Eur. J.*, 2016, **22**, 5416.
- 46 S. Batsyts, R. Vedmid, J. C. Namyslo, M. Nieger, A. Schmidt, *Eur. J. Org. Chem.*, 2019, 1301.
- 47 M. S. El-Shall, K. P. C. Vollhardt, *J. Mol. Struct. (Theochem)*, 1989, 183, 175.
- 48 A. A. Granovsky, Firefly version 8, www http://classic.chem.msu.su/gran/firefly/index.html.
- 49 M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. Su, T. L. Windus, M. Dupuis, J. A. Montgomery, *J. Comput. Chem.*, 1993, **14**, 1347.
- 50 NBO 5.9, E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales, and F. Weinhold (Theoretical Chemistry Institute, University of Wisconsin, Madison, WI) 2012.
- 51 M. -H. Son, J. Y. Kim, E. J. Lim, D. -J. Beak, K. Choi, J. K. Lee, A. N. Pee, S. -J. Min, Y. S. Cho, *Bioorg. Med. Chem. Lett.*, 2013, **23**, 1472.

View Article Online
DOI: 10.1039/C9OB00055F