

This article was downloaded by: [University of Tennessee, Knoxville]

On: 22 March 2013, At: 09:15

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

### IMPROVED SYNTHESIS OF PHENYLSELENOGLYCOLIC ACIDS

Hamdi Özkan <sup>a</sup>, Ali Disli <sup>a</sup> & Yilmaz Yildirim <sup>a</sup>

<sup>a</sup> Department of Chemistry, Faculty of Arts and Sciences, Gazi University, 06500, Teknikokullar, Ankara, TURKEY

Version of record first published: 09 Feb 2009.

To cite this article: Hamdi Özkan, Ali Disli & Yilmaz Yildirim (2004): IMPROVED SYNTHESIS OF PHENYLSELENOGLYCOLIC ACIDS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 36:2, 161-163

To link to this article: <http://dx.doi.org/10.1080/00304940409355388>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

14. "Chemical Reagent. Fine Chemicals Handbook (Chinese)", Chemical Industry Publishing House, Huixin li 3, Chaoyang District, Beijing, 2002
15. M. Darmon, *Bull. Soc. Chem.Fr.*, **8**, 414 (1941).
16. N. Allentoff and G. F Wright, *J. Org. Chem.*, **22**, 1 (1957).


\*\*\*\*\*

## IMPROVED SYNTHESIS OF PHENYLSELENOGLYCOLIC ACIDS

Submitted by Hamdi Özkan, Ali Disli, and Yılmaz Yıldırım\*  
(10/28/03)

*Department of Chemistry, Faculty of Arts and Sciences  
Gazi University, 06500, Teknikokullar, Ankara, TURKEY*


Selenium is a nonmetallic trace element recognized as a nutrient essential to human health.<sup>1,2</sup> Selenium is also an essential constituent of extracellular and cellular glutathione peroxidases, thyroidal and extrathyroidal iodothyronine 5'-deiodinases, thioredoxin reductase, and other selenoproteins.<sup>2</sup> Various experimental models showed that selenium inhibits tumorigenesis.<sup>3</sup> Low serum selenium levels are associated with an increased risk of developing cancer at several sites, especially cancers of the stomach and lung for men<sup>4</sup>. Thus, many organoselenium compounds have been synthesized.<sup>5,6</sup> Some substituted phenylselenoglycolic acids have been synthesized by using Grignard reagent<sup>7</sup> (*Scheme 1*) (yields 20-25%) and from



**Scheme 1**

diazonium salts.<sup>8,9</sup> In the studies with diazonium salts, substituted anilines were diazotized in aqueous medium, followed by addition of potassium selenocyanide. Although yields were not reported,<sup>8,9</sup> we found them to be lower (25-30%) when the last traces of acids were not removed and reaction performed in aqueous medium. In the presence of acid, potassium selenocyanide decomposes to release poisonous hydrogen cyanide and concurrent decrease in the yields.

In our method (*Scheme 2*), substituted anilinium chlorides are prepared, dried and washed with ether to remove excess acid from the salts. The anilinium chlorides are diazotized with ethyl nitrite in non-aqueous medium, thus avoiding the decomposition of potassium selenocyanide.



Scheme 2

Table 1. Yields, mps <sup>1</sup>H NMR and IR Spectral Data of 4

Cmpd	mp (°C)	<i>lit</i> mp (°C)	Yield (%)	<sup>1</sup> H NMR (δ) (CDCl <sub>3</sub> ; CF <sub>3</sub> COOH)	IR (cm <sup>-1</sup> )			
					O-H,	C=O,	=CH,	CH
4a	89-90	87-89 <sup>a</sup>	34	3.70 (s, 2H); 7.24 (m, 4H); 10 (s, 1H)	3480,	1703,	3075,	2993
4b	110-113	114 <sup>b</sup>	46	3.40 (s, 2H); 7.0-7.78 (m, 4H); 9.8 (s, 1H)	3428,	1705,	3074,	2921
4c	90-91	92 <sup>a</sup>	36	3.60 (s, 2H); 7.1-7.4 (m, 4H); 9.1 (s, 1H)	3461,	1708,	3082,	2921
4d	126-127	127 <sup>c</sup>	50	3.40 (s, 2H); 7.0 (m, 4H); 9.0 (s, 1H)	3454,	1709,	3082,	2903
4e	94-95	97-98 <sup>b</sup>	48	2.19 (s, 3H); 3.38 (s, 2H); 6.95-7.24 (m, 4H); 9.78 (s, 1H)	3435,	1702,	3070,	2928
4f	160-163	165 <sup>b</sup>	35	3.40 (s, 2H); 7-8 (m, 4H) 9.2 (s, 1H)	3461,	1715,	3024,	2921
4g	88-90	90-91 <sup>b</sup>	40	3.90 (s, 2H); 7.40-8.11 (m, 4H)	3505,	1708,	3089,	2921
4h	117-120	119-120 <sup>b</sup>	48	3.38 (s, 2H); 6.95-7.24 (m, 4H)	3416,	1702,	3262,	2900
4i	245-250	250 <sup>a</sup>	45	3.20 (s, 2H); 7.0-8.0 (m, 4H); 9.1 (s, 1H)	3448,	1683,	3121	2928

a) Ref 9; b) Ref 8; c) Ref 7

## EXPERIMENTAL SECTION

The anilines and other chemicals were supplied by Merck Co. All mps were determined in sealed capillaries and are uncorrected. FT-IR spectra were recorded on a Matson 1000 spectrometer as KBr pellets. <sup>1</sup>H NMR spectra were obtained on a Varian EM-360 L (60 MHz) NMR spectrometer and a Varian Gemini 200 (200 MHz) NMR spectrometer in CDCl<sub>3</sub>.

**2-Chlorophenylselenoglycolic acid (4a). Typical Procedure.** 2-Chloroaniline (1.08 mL, 0.01 mole) and conc. HCl (2.7 mL, 0.04 mole) were mixed slowly. The anilinium chloride was collected, air-dried and dissolved in 30 mL of absolute alcohol. The stirred solution was cooled -5-0°C and diazotized by the careful dropwise addition (over a period of 30 minutes) of ethyl nitrite (1 mL, 0.006 mole). Both the solution and ethyl nitrite (its bp 17°C) must be cold (-5-0°C) during the addition. Then a solution of KSeCN (1.44 g, 0.01 mole) in 30 mL of absolute alcohol was added into the solution of the diazonium salt. The precipitate which formed was collected, air dried and dissolved in about 30 mL of boiling ethanol. Then, a solution of NaOH (0.8 g, 0.02 mole) in ethanol was added to the boiling solution and this mixture was refluxed 2 hrs. The mixture was cooled at room temperature and then 5 mL water was added. Then a solution of

sodium chloroacetate (0.01 mole, 1.165 g) in cold water was poured into the final solution and the mixture was boiled under reflux for 15 minutes. The alcohol was evaporated *in vacuo*. The cooled mixture was filtered and HCl was added the filtrate until precipitation was complete. The precipitate obtained was recrystallized from benzene (water for **4c-4i**) to afford 0.85 g (34%) of colorless needles, mp. 89-90°C, *lit.*<sup>a</sup> 87-89°C.

## REFERENCES

1. X. Chen, G. Yang, J. Chen, X. Chen, Z. Wen and K. Ge, *Biol. Trace. Elem. Res.*, **2**, 91 (1979).
2. E. A. Klein, I. M. Thompson, S. M. Lippmann, P. J. Goodman, D. Albanes, P. R. Taylor and C. Coltman, *Urologic Oncology: Seminars and Original Investigations*, **21**, 59 (2003).
3. Jr. G. F. Combs and L. C. Clark, Selenium and Cancer in: H. Garewall, editor, *Antioxidants and Disease Prevention*, New York, CRC Press (1997).
4. P. Knekt, A. Aromaa, J. Maatela, G. Alfthan, R. K. Aaran, M. Hakama, T. Hakulinen, R. Peto and L. Teppo, *J. Natl. Cancer Inst.*, **82**, 864 (1990).
5. Y. Xie, M. D. Short, P. B. Cassidy and J.C. Roberts, *Bioorg. Med. Chem. Lett.*, **11-22**, 2911 (2001).
6. K. Overvad, *Bibl. Nutr. Diet.*, **54**, 141 (1998).
7. G. T. Morgan and W. H. Porritt., *J. Chem. Soc.*, 1755 (1925).
8. O. Behagel and M. Rollmann, *J. prakt. Chem.*, **123**, 336 (1929).
9. L. D. Pettit, A. Royston, C. Sherrington and R. J. Whewell, *J. Chem. Soc. (B)*, 588 (1968).

\*\*\*\*\*