

Synthesis and Antifungal Activity of β -Trifluoroalkyl Aminovinyl Ketone Derivatives

Gary Gellerman,^{*,†} Natali Pariente,[†] Zahi Paz,[‡] Anat Shnaiderman,[‡] and Oded Yarden^{*,‡}

[†]Department of Biological Chemistry, Ariel University Center of Samaria, Ariel 40700, Israel, and [‡]Department of Plant Pathology and Microbiology, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot 76100, Israel

Ten β -trifluoroalkyl aminovinyl ketone derivatives were synthesized, and their inhibitory effects on several phytopathogenic fungi, an oomycete and plants were assessed. The various compounds were fungitoxic at the 10–100 μ M range, with (*Z*)-3-amino-4,4,4-trifluoro-1-(4-chlorophenyl)but-2-en-1-one exhibiting the highest inhibitory effect on most of the test pathogens. *Alternaria alternata* and *Neurospora crassa* were the most tolerant and sensitive fungi to the compounds, respectively. We propose that (*Z*)-3-amino-4,4,4-trifluoro-1-phenylbut-2-en-1-one is the minimal structural requirement for a β -trifluoroalkyl aminovinyl ketone fungitoxic derivative.

KEYWORDS: Antifungal; β -trifluoroalkyl aminovinyl ketone derivatives; structure–activity relationship; Alternaria alternata; Penicillium digitatum; Rhizoctonia soliani; Botrytis cinerea; Sclerotinia sclerotiorum; Neurospora crassa; Pythium aphanidermatum

INTRODUCTION

In many agricultural practices, fungicides are extensively used to prevent and curb the progression of plant diseases that have severe adverse effects on crop yields and quality. In some cases, the prolonged and repeated use of many fungicides has resulted in the proliferation of plant pathogen isolates that are resistant to the fungicides employed. Therefore, identifying new classes of antifungal compounds based on structures not previously used in combating plant diseases may assist in the control of plant pathogens that are tolerant or resistant to known compounds.

Sclerotinia sclerotiorum and Botrytis cinerea are necrotrophic phytopathogenic filamentous ascomycetes known to attack more than 400 and 200 plant species, respectively (1-7). Diseases caused in economically important plants by these fungi occur worldwide, cause considerable damage, and have proven difficult to control (culturally or chemically), and host resistance to these fungi is inadequate. Annual losses of crops from diseases caused by S. sclerotiorum and B. cinerea are in the multimillion dollar range (e.g., http://www.whitemoldresearch.com). In addition, their ubiquitous prevalence along with the intensive use of fungicides for their control has resulted in repeated appearance of fungicide-resistant strains (8-11). In order to expand the range of chemicals that could potentially be harnessed to control these fungi, we have synthesized a group of (Z)- β -trifluoroalkyl aminovinyl ketones and assessed their fungitoxic activity. We have determined that some of these compounds can curb fungal growth at the 10–100 μ M range. The potential utilization of these molecules is supported by the fact that synthesis of these compounds consists of two simple stages using inexpensive and commercially available compounds.

MATERIALS AND METHODS

Analytical HPLC was performed on a 250×4.2 mm Lichroprep RP-18 column from Merck, with a 1 mL/min flow and detection at 214 nm. The eluents were triply distilled water and HPLC-grade CH₃CN (containing 0.1% TFA) or MeOH. Mass spectra were measured in the positive and negative modes using a quadrupole mass spectrometer equipped with an electrospray ionization source and cross-flow inlet. ¹H and ¹³C NMR spectra were recorded at 300 and 75 MHz, respectively in CDCl₃, unless otherwise indicated. Assignments in the final products were supported by 2D COSY, HMBC and HMQC spectroscopy. All chemical shifts are reported with respect to TMS. Chromatography was carried out by standard flash chromatography and TLC on silica gel (Merck 7735).

Compounds and Syntheses. Preparation of Trifluoroacetonitrile. Sulfuric acid (2 mL) was slowly added to 100 mL of methanol while stirring in an ice bath. Trifluoroacetic acid (118 gr, 78 mL) was added to the cooling methanol solution. The mixture was stirred while heating (40–45 °C) for 6 h. After cooling to rt, water was added and the resulting trifluoromethyl ester organic phase was separated and dried over anhydrous sodium sulfate. After filtration, 1 L of a 2.0 M solution of ammonia in methanol was added to the filtrate, and the mixture was stirred overnight. Solvent and unreacted ammonia were evaporated, and the white solid was dried under vacuum (98 g) and used for the next step without any further purification. The obtained trifluoroacetamide was rapidly mixed with phosphorus pentaoxide (P₂O₅) (in 3-fold excess) in a 500 mL flask (the condensation reaction starts immediately). The reaction mixture was heated to 150 °C, and the obtained gaseous trifluoroacetonitrile was directly bubbled into the following reaction.

General Procedure for Preparation of β -Trifluoroalkyl Aminovinyl Ketones **5**a-**j**. 4 mmol of acetophenone was added in a dropwise manner to a mixture of 10 mmol of sodium *tert*-butoxide in

^{*}Authors to whom correspondence should be addressed. G.G.: tel, (972) 03 9371442; fax, 972-3-9066634; e-mail, Garyg@ariel.ac.il. O.Y.: tel, (972) 08 9489298; fax: (972) 08 9468785; e-mail, Oded.Yarden @huji.ac.il.

100 mL of dry THF. Then, trifluoroacetonitrile (produced from 15 g of amide and 45 g of P_2O_5) was bubbled in for 2 h to the reaction vessel equipped with a dry ice-acetone condenser. After bubbling, the mixture was stirred for 5 h at rt. A solution (10 mmol) of commercial HCl-dioxane 4 N was added dropwise, and after 10 min the mixture was filtrated and dioxane was evaporated under vacuum. The crude product was recrystallized from hexane.

Preparation of (*Z*)-4-(3-*Amino*-4,4,4-*trifluorobut*-2-*enoyl*)*benzonitrile* (5*a*). Compound 5*a* was prepared from 3 and 4-cyanoacetophenone (1 g, 6.9 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent, yielded 1.44 g of 5*a* (87% yield) as a colorless oil: v_{max} (KBr) 2880, 1670, 1635, 1270 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₁H₇F₃N₂O (MH⁺) 241.05, found 241.04; ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 7.68 (d, 2, *J* = 10.7 Hz, H-2), 8.14 (d, 2, *J* = 10.7 Hz, H-3); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 160.7, 142.2, 132.7, 131.5, 130.6, 118.4, 115.7, 95.5.

Preparation of (Z)-3-Amino-4,4,4-trifluoro-1-(4-fluorophe-nyl)but-2-en-1-one (5b). Compound **5b** was prepared from **3** and 4-fluoroacetophenone (1 g, 7.3 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.28 g of **5b** (85% yield) as a colorless oil: ν_{max} (KBr) 1670, 1650, 1190 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₀H₇F₄NO (MH⁺) 234.046, found 234.052; ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 7.06 (t, 2, *J* = 5.35 Hz, H-3), 7.66 (m, 2, H-2); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 168.7, 160.7, 133.5, 131.5, 116, 95.5.

Preparation of (Z)-3-Amino-4,4,4-trifluoro-1-(4-chlorophenyl)but-2-en-1-one (5c). Compound **5c** was prepared from **3** and 4-chloroacetophenone (1 g, 6.5 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.38 of **5c** (81% yield) as a colorless oil: ν_{max} (KBr) 1670, 1590, 1380, 1060 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₀H₇ClF₃NO (MH⁺) 250.017, found 250.020: 252.018 (3:1); ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 7.41 (d, 2, *J*=10.7 Hz, H-3), 7.82 (d, 2, *J*=10.7 Hz, H2). ¹³C NMR (δ , ppm, CDCl₃) 189.7, 160.7, 140.1, 136, 131.5, 131.3, 129.3, 95.0.

Preparation of (Z)-3-Amino-4,4,4-trifluoro-1-(4-hydroxy-phenyl)but-2-en-1-one (5d). Compound **5d** was prepared from **3** and 4-hydroxyacetophenone (1 g, 7.4 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (6:4) as eluent yielded 1.19 g of **5d** (76% yield) as a white powder; ν_{max} (KBr) 3500–3180 (bs), 1660, 1520, 1450, 1230 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₀H₈F₃NO₂ (MH⁺) 232.05, found 232.04; ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 6.89 (d, 2, *J*=10.7 Hz, H-3), 7.82 (d, 2, *J*=10.7 Hz, H-2); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 164.3, 160.7, 131.3, 130.5, 125.5, 116.4, 94.5.

Preparation of (Z)-tert-Butyl-4-(3-amino-4,4,4-trifluorobut-2-enoyl)phenyl carbamate (5e). Compound 5e was prepared from 3 and 4-BOC-amino acetophenone (1 g, 4.3 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (6:4) as eluent yielded 1.01 g of 5e (72% yield) as a colorless oil: ν_{max} (KBr) 1660, 1500, 1350, 1120 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₅H₁₇F₃N₂O₃ (MH⁺) 331.119, found 331.126; ¹H NMR (CDCl₃) δ 1.5 (s, 9, ¹Bu), 6.19 (s, 1, CH=C), 7.45 (d, 2, *J*=10.7 Hz, H-3), 7.85 (d, 2, *J*=10.7 Hz, H-2); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 164.3, 160.7, 131.5, 131.3, 130.5, 116.4, 90.0.

Preparation of (Z)-3-Amino-4,4,4-trifluoro-1-(naphthalen-2-yl)but-2-en-1-one (5f). Compound **5f** was prepared from **3** and acetonaphthone (1 g, 5.9 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (7:3) as eluent yielded 1.11 g of **5f** (74% yield) as a colorless oil: ν_{max} (KBr) 1655, 1570, 1360, 1240 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₄H₁₀F₃NO (MH⁺) 266.071, found 266.077; ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 7.58 (m, 3, H-3,7,8), 7.99 (m, 3, H-4,5,6), 8.40 (s, 1, H-2); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 160.7, 135.6, 134.7, 132.5, 131.5, 129.5, 128.6, 128.5, 128.3, 127.7, 126.9, 124.2, 93.0.

Preparation of (Z)-3-Amino-4,4,4-trifluoro-1-(3-(trifluoromethyl)phenyl)but-2-en-1-one (5g). Compound 5g was prepared from 3 and 3-trifluoromethyl acetonaphthone (1 g, 4.29 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.02 of **5g** (84% yield) as a colorless oil: ν_{max} (KBr) 1665, 1570, 1360, 1240 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₁H₇F₆NO (MH⁺) 284.048, found 284.051; ¹H NMR (CDCl₃) δ 6.21 (s, 1, CH=C), 7.6 (t, 1, *J*=8.82 Hz, H-3), 7.8 (d, 2, *J*=8.82 Hz, H-2), 8.09 (d, 2, *J*=8.82 Hz, H-4), 8.19 (s, 1, H-6); ¹³C NMR (δ , ppm, CDCl₃) 189.7, 160.7, 138.2, 133.2, 131.5, 130.9, 129.6, 126.2, 124.2, 94.0.

Preparation of (*Z*)-3-*Amino-1-(3-bromophenyl*)-4,4,4-tri*fluorobut-2-en-1-one* (*5h*). Compound **5h** was prepared from **3** and 3-bromoacetophenone (1 g, 5.0 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.32 g of **5h** (89% yield) as a colorless oil: ν_{max} (KBr) 1675, 1640, 1410. 1305, 1060 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₀H₇BrF₃NO (MH⁺) 294.295, found 293.972, 295.974 (1:1); ¹H NMR (CDCl₃) δ 6.19 (s, 1, CH=C), 7.38 (t, 1, *J*=6.25, 1H, H-4), 7.69 (d, 1 *J*=12.5 Hz, H-3), 7.81 (d, 1, *J*=12.5 Hz, H-2), 8.09 (s, 1, H-6). ¹³C NMR (δ, ppm, CDCl₃) 189.7, 160.7, 140.1, 137.4, 133.4, 131.5, 128.9, 123.6, 95.5.

*Preparation of (Z)-3-Amino-1-(4-chlorophenyl)-2-ethyl-*4,4,4-trifluorobut-2-en-1-one (5i). Compound 5i was prepared from 3 and 4-chlorobutyrophenone (1 g, 5.5 mmol) following the general procedure described above. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.17 g of 5i (68% yield) as a colorless oil: v_{max} (KBr) 1665, 1630, 1390 cm⁻¹; HRMS (DI, *m/ z*) calcd for C₁₂H₁₁ClF₃NO (MH⁺) 277.048, found 277.043:277.044 (3:1); ¹H NMR (CDCl₃) δ 0.98 (t, 3, *J*=9.31 Hz, CH₂CH₃), 2.00 (q, 2, *J*=9.31 Hz, C=CCH₂CH₃), 7.51 (d, 2, *J*=10.34, H-3), 7.86 (d, 2, *J*=10.34, H-4); ¹³C NMR (δ, ppm, CDCl₃) 190.5, 152.5, 140.1, 136, 131.3, 129.3, 125.3, 112.9, 12.6, 11.6.

Preparation of (*Z*)-3-*Amino-1-(4-chlorophenyl)-4,4,5,5,5pentafluoropent-2-en-1-one* (*5j*). Compound *5j* was prepared from gaseous 2,2,3,3,3-pentafluoropropanenitrile (prepared from commercial CF₃CF₂CONH₂ in the same manner as for **3** and 4-chlorobutyrophenone (1 g, 6.9 mmol) following the general procedure. Purification by column chromatography on silica gel using *n*-hexane–EtOAc (8:2) as eluent yielded 1.18 g of *5j* (57% yield) as a colorless oil: v_{max} (KBr) 1660, 1600, 1470, 1290 cm⁻¹; HRMS (DI, *m/z*) calcd for C₁₁H₁₇CIF₅NO (MH⁺) 300.014, found 300.016:302.017 (3:1); ¹H NMR (CDCl₃) δ 6.23 (s, 1, CH=C), 7.40 (d, 2, *J* = 10.7 Hz, H-3), 7.84 (d, 2, *J* = 10.7 Hz, H-4; ¹³C NMR (δ, ppm, CDCl₃) 191.2, 161.6, 141.7, 1364, 136.9, 133.8, 130.6, 127.1, 97.3.

Fungal Strains and Culturing Conditions. The fungi Sclerotinia sclerotiorum (isolate 1980 (12)), Botrytis cinerea (isolate BcI16 (13)), Penicillium digitatum (14), Alternaria alternata and Rhizoctonia solani (both from the fungal collection of the Dept. of Plant Pathology and Microbiology) and Neurospora crassa (strain 74-OR23-1A, The Fungal Genetics Stock Center (15)) were used during this study. In addition, the oomycete Pythium aphanidermatum (from the Dept. of Plant Pathology and Microbiology) was also used as a test organism. Strains were cultured on potato dextrose agar (PDA, Difco Laboratories, Detroit, MI), unless otherwise stated.

Scheme 1. General Synthesis of β -Trifluoroalkyl Aminovinyl Ketone Derivatives 5a-j



 $\begin{array}{l} \textbf{5a} \ \mathbf{R_1} = p \ - \mathbf{CN}, \ \mathbf{R_2} = \ \mathbf{H}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \\ \textbf{5b} \ \mathbf{R_1} = p \ - \mathbf{F}, \ \mathbf{R_2} = \ \mathbf{H}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \\ \textbf{5c} \ \mathbf{R_1} = p \ - \mathbf{CR}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \\ \textbf{5c} \ \mathbf{R_1} = p \ - \mathbf{OH}, \ \mathbf{R_2} = \ \mathbf{H}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \\ \textbf{5c} \ \mathbf{R_1} = p \ - \mathbf{OHBoc}, \ \mathbf{R_2} = \ \mathbf{H}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \\ \textbf{5c} \ \mathbf{R_1} = p \ - \mathbf{OHBoc}, \ \mathbf{R_2} = \ \mathbf{H}, \ \mathbf{R_3} = \ \mathbf{H}, \ \mathbf{n} = 0 \end{array}$

 $\begin{array}{l} \textbf{5f} \ R_1, R_2= naphthalen-2-yl, R_3=H, n=0\\ \textbf{5g} \ R_1=m\text{-}CF_3 \ , R_2=H, R_3=H, n=0\\ \textbf{5h} \ R_1=m\text{-}Br, R_2=H, R_3=H, n=0\\ \textbf{5i} \ R_1=p\text{-}Cl, R_2=H, R_3=Et \ , n=0\\ \textbf{5j} \ R_1=p\text{-}Cl, R_2=H, R_3=H \ , n=1 \end{array}$

Article

Assessment of Fungitoxicity. Initial toxicity against the fungi (excluding *S. sclerotiorum*) and the oomycete was assessed by culturing the test organisms on Petri dishes containing 10 mL of PDA amended with either 10 or 100 μ M of the ten original test compounds. Stock solutions of the different compounds were prepared in dimethyl sulfoxide (DMSO), and aliquots (not exceeding 0.001% v/v, a concentration that had no observable effect on the test organisms) were added to warm (50 °C) sterile medium prior to dispensing to the Petri dishes. *S. sclerotiorum* and *B. cinerea* were used for the advanced phase of tests, where the effects of compound concentrations (0–50 μ M) were tested. The two fungi were cultured at 17 and 22 °C, respectively, for 48 h, after which radial growth was measured and compared to that of the fungal strains cultured in medium not amended with the antifungal compounds.

RESULTS AND DISCUSSION

Phenyl α , β -unsaturated carbonyl derivatives of cinnamic and coumaric are known backbones found in potent compounds in antifungal discovery (16). In recent years research based on these core structures has become an important tool for the generation





 $(Z)\beta$ -trifluoroalkyl aminovinyl ketones

 $(CF_2)_n CF_3$

Figure 1. Structure of the compounds prepared in this study.

of new leads and for providing access to diverse chemical entities with novel structures and properties (17). We decided to explore additional compounds that share some similarity with the cinnamic acid backbone in a scaffold hopping strategy toward the generation of novel antifungal compounds. In order to identify potential antifungal compounds, we designed and synthesized β -trifluoroalkyl aminovinyl ketone molecules that differed in the R₁, R₂, and R₃ substituents as well as in the length of the fluorocarbon chain in the scaffold (structure **5** in **Scheme 1**).

Several synthetic routes to the synthesis of fluoroalkyl-containing β -aminovinyl ketones starting from unsymmetrical fluoroalkyl-containing β -diketones and amines have been previously described (18-21). Unfortunately, these reactions result in poor to moderate yields. We adopted a different approach for trifluoroalkyl β -aminovinyl ketone synthesis: namely, condensation of trifluoroacetonitrile (22) with acetophenones in the presence of a strong base in dry aprotic solvents. After examining several alternatives, sodium tert-butoxide (NaO^tBu) and dry THF were chosen as a base and solvent of choice respectively. This general method was applied very successfully in the preparation of 5a-j(Scheme 1). Thus, we initially prepared gaseous CNCF₃ (3) from the corresponding trifluoroacetamide by dehydration with P_2O_5 at high temperature. Compound 3 was then directly bubbled into the reaction mixture of the appropriate acetophenone in the presence of NaO^tBu under nitrogen atmosphere in dry THF. The reaction vessel was equipped with an acetone/dry ice coldfinger to avoid the undesired escape of gaseous 3. After completion (monitored by TLC and HPLC), the reactions were quenched with dilute HCl, resulting in crude products that after



Figure 2. Effect of nine β -trifluoroalkyl aminovinyl ketone derivatives on radial growth of six test organisms. Test organisms were cultured on potato dextrose agar in the presence of either 10 or 100 μ M concentrations (empty and full bars, respectively) of the various compounds. Y-error bars indicate standard deviation.



Figure 3. Effect of five β -trifluoroalkyl aminovinyl ketone derivatives on radial growth of *Sclerotina sclerotiorum* and *Botrytis cinerea*. Y-error bars indicate standard deviation.

subsequent purification by flash chromatography on silica gel (EtOAc/ *n*-hexane), pure **5a**-**i** were obtained in good yields (68–89%). Even a compound with the longer flourocarbone chain (**5j**) was prepared in the same manner as **5a**-**i**, using gaseous 2,2,3,3,3-pentafluoropropanenitrile (initially prepared from commercial CF₃CF₂CONH₂ and P₂O₅) and 4-chlorobutyr-ophenone in 57% yield.

Coumaric and cinnamic acids (**Figure 1**), but not caffeic acid, have been shown to be inhibitory to fungal (*Aspergillus* 'spp') growth (23). Apparently, the location of the substituent on the cinnamic acid backbone plays crucial role in determining the antifungal activity of the compound. This data prompted us to examine other phenyl α , β -unsaturated carbonyl scaffolds substituted with different functional groups at various locations and study their potential antifungal activity.

Almost all of the synthesized compounds exhibited some degree of toxicity toward the filamentous fungi tested (Figure 2). Nonetheless, significant differences in toxicity were observed among these compounds. For example, the highest toxicity levels were evident when with (Z)-3-amino-4,4,4-trifluoro-1-(4-chlorophenyl)but-2-en-1-one (5c) was present in the growth medium, while 5j, 5i and 5e, respectively, were significantly less toxic. The fungi least susceptible to most of the compounds tested were A. alternata and P. digitatum. None of the compounds tested inhibited radial growth of A. alternata more than 50% (even at a 100 μ M concentration) and only 5c inhibited P. digitatum by more than 50%. In contrast, N. crassa appeared to be the most sensitive to the compounds at the higher concentrations tested, 5c, 5h and 5b inhibited radial growth almost entirely. The oomycete P. aphanidermatum was sensitive to higher concentrations of 5d, 4d, 5c and 5h. The other compounds tested inhibited its radial growth only to a limited extent. Based on the outcome of the initial assessment of toxicity of the compounds to the test organisms, we decided to focus our analysis on five of the original compounds tested 5d, 4b, 5c, 5h and 5j and their toxicity to the omnivorus phytopathogen B. cinerea and an additional broad-host pathogen related to B. cinerea: S. sclerotiorum. This analysis allowed us to determine if these phylogenetically related pathogens differ in their sensitivity to the structurally related tested compounds. In most instances, the fungal response to the tested compounds was dose dependent (Figure 3). Both fungi

Gellerman et al.

exhibited very similar sensitivity to 5d and 5c. Commercially available 4d, that lacks the trifluoroprop-1-en-2-amine part in the backbone, was the least active compound and, interestingly, at the lower concentrations may have even stimulated the radial growth of S. sclerotiorum. This was also evident at the lower concentrations of 5j. Overall, the most effective compound (in terms of curbing radial growth) was 5h, even though the sensitivity of *B. cinerea* to this compound was significantly higher, at all concentrations, than that of S. sclerotiorum. Based on the results obtained it appears that (Z)-3-amino-4,4,4-trifluoro-1-phenylbut-2-en-1-one is the minimal scaffold for presenting β -trifluoroalkyl aminovinyl ketone fungitoxic activity, preferably substituted with a weak electron withdrawal halogen atom at *para* or *meta* positions. These findings emphasize the potential that some of these molecules have for fungicide development. Some of these compounds are potential leads to be considered for practical use, therefore, future full analysis of their biological activities need to be determined. One of these is their potential phytotoxicity. To that end, we performed preliminary tests on effect of the presence of the compounds on seed germination of alfalfa (Medicago sativa L.) and on induced phytotoxic response on leaves of tomato (Solanum lycopersicum L.), which are both hosts of B. cinerea and S. sclerotiorum. In all cases, no significant phytotoxic effects were observed (Gellerman et al., unpublished results). Work is in progress to assess the phytotoxic effects of this family of compounds.

Taken together, we have determined that a novel group of compounds, sharing the β -trifluoroalkyl aminovinyl ketone scaffold, exhibit antifungal activities *in vitro*. On the one hand, their activity is evident at relatively high concentrations (similar to those determined for most cinnamic acid derivatives described by Bisogno et al. (16)), even though at the highest tested concentrations fungal inhibition was not complete. On the other hand, the ease of synthesis of these stable compounds along with their apparent lack of phytotoxicity makes these compounds attractive for future consideration as potential components of antifungal intervention.

ACKNOWLEDGMENT

O.Y. is the incumbent of the Buck Family Chair in Plant Pathology.

Supporting Information Available: NMR spectra of compounds **5a–c,h** and HRMS of compounds **5c,f–h**. This material is available free of charge via the Internet at http://pubs.acs.org.

LITERATURE CITED

- Boland, G. J.; Hall, R. Index of plant hosts of *Sclerotinia sclerotior-um. Can. J. Plant Pathol.* **1994**, *16*, 93–108.
- (2) Bolton, M. V.; Bart, P. H.; Thomma, J.; Nelson, B. D. Sclerotinia sclerotiorum (Lib.) de Bary: biology and molecular traits of a cosmopolitan pathogen. *Mol. Plant Pathol.* **2006**, 7, 1–16.
- (3) Choquer, M.; Fournier, E.; Kunz, C.; Caroline Levis, C.; Pradier, J.-M.; Simon, A.; Viaud, M. *Botrytis cinerea* virulence factors:new insights into a necrotrophicand polyphageous pathogen. *FEMS Microbiol. Lett.* 2007, 277, 1–10.
- (4) Erental, A.; Dickman, M. B.; Yarden, O. Sclerotial development in *Sclerotinia sclerotiorum*: awakening molecular analysis of a "dormant" structure. *Fung. Biol. Rev.* 2008, 22, 6–16.
- (5) Van Kan, J. A. L. Licensed to kill: the lifestyle of a necrotrophic plant pathogen. *Trends Plant Sci.* 2006, 11, 247–253.
- (6) Purdy, L. H. Sclerotinia sclerotiorum: history, diseases and symptomatology, host range, geographic distribution and impact. *Phyto*pathology **1979**, 69, 875–880.

- (7) Tu, J. C. An integrated control of white mold (*Sclerotinia sclerotiorum*) of beans, with emphasis on recent advances in biological control. *Bot. Bull. Acad. Sinica* 1997, *38*, 73–76.
- (8) Gossen, B. D.; Rimmer, S. R.; Holley, J. D. First report of resistance to benomyl fungicide in *Sclerotinia sclerotiorum*. *Plant Dis.* 2001, 85, 1206.
- (9) Ma, Z. H.; Michailides, T. J. Advances in understanding molecular mechanisms of fungicide resistance and molecular detection of resistant genotypes in phytopathogenic fungi. *Crop Prot.* 2005, 24, 853–863.
- (10) Mendoza, L.; Araya-Maturana, R.; Cardona, W.; Delgado-Castro, T.; Garcia, C.; Lagos, C.; Cotoras, M. In vitro sensitivity of *Botrytis cinerea* to anthraquinone and anthrahydroquinone derivatives. J. Agric. Food Chem. 2005, 53, 10080–10084.
- (11) Yarden, O.; Katan, T. Mutations leading to substitutions at amino acids 198 and 200 of beta tubulin that correlate with benomylresistance phenotypes of field strains of *Botrytis cinerea*. *Phytopathology* **1993**, *83*, 1478–1483.
- (12) Godoy, G.; Steadman, J. R.; Dickman, M. B.; Dam, R. Use of mutants to demonstrate the role of oxalic acid in pathogenicity of *Sclerotinia sclerotiorum* on *Phaseolus vulgaris*. *Physiol. Mol. Plant Pathol.* **1990**, *37*, 179–191.
- (13) Swartzberg, D., Kirshner, B., Elad, Y., Granot, D. 2008. *Botrytis cinerea* induces senescence and is inhibited by autoregulated expression of the IPT gene. *Eur. J. Plant Pathol.* 2008, *120*, 289-297.
- (14) Yarden, O.; Katan, J.; Aharonson, N. A rapid bioassay for the determination of carbendazim residues in soil. *Plant Pathol.* 1985, 34, 69–74.
- (15) McCluskey, K., The fungal genetics stock center: From molds to molecules. Adv. Appl. Microbiol. 2003, 52, 245–262.

- (16) Bisogno, F.; Mascoti, L.; Sanchez, C.; Enriz, R. Structure-antifungal activity relationship of cinnamic acid derivatives. J. Agric. Food Chem. 2007, 55, 10635–10640.
- (17) Narasimhan, B.; Belsare, D.; Pharande, D.; Dhake, D. Esters, amides and substituted derivatives of cinnamic acid: synthesis, antimicrobial activity and QSAR investigations. *Eur. J. Med. Chem.* 2004, *39*, 827–834.
- (18) Pashkevich, K. I.; Filyakova, V. I.; Postovskii, I. Ya. Competitive amination of carbonyl groups in unsymmetrical polyfluoroalkylarylcontaining β-diketones. *Izv. Akad. Nauk SSSR, Ser. Khim.* **1981**, *10*, 2346–2349.
- (19) Filyakova, V. I.; Ratner, V. G.; Karpenko, N. S.; Pashkevich, K. I. Reaction of fluoroalkyl-containing β-diketones with amines. *Izv. Akad. Nauk, Ser. Khim.* **1996**, *9*, 2278–2284.
- (20) Boltacheva, N. S.; Filyakova, V. I.; Charushin, V. N. Fluoroalkylcontaining lithium 1,3-diketonates in reactions with amines and ammonium salts. *Russ. J. Org. Chem.* 2005, *41*, 1452–1457.
- (21) Sosnovskikh, V. Ya.; Gor'kii, A. M. Condensation of nitriles of polyhalogenated carboxylic acids and benzonitrile with 2-hydroxy-4,6-dimethylacetophenone. *Russ. Chem. Bull.* **1998**, *47*, 354–356.
- (22) Sosnovskikh, V. Ya.; Ovsyannikov, I. S. Condensation of trichloroand trifluoroacetonitrile with 4-hydroxy-4-methyl-2-pentanone and 2'-hydroxyacetophenone. *Zh. Org. Khim.* **1993**, *29*, 89–93.
- (23) Kim, J. H.; Campbell, B. C.; Mahomey, N. E.; Chan, K. L.; Molyneux, R. J. Identification of phenolics for control of *Aspergillus flavus* using *Saccharomyces cerevisiae* as a target-gene bioassay. *J. Agric. Food Chem.* **2004**, *52*, 7814–7821.

Received March 19, 2009. Revised manuscript received August 17, 2009. Accepted August 17, 2009. This research was supported by BARD, the United States—Israel Binational Research and Development Fund (to O.Y.).