## Linking Ion and Neutral Chemistry in C–H Bond Electrophilic Activation: Generation and Detection of HO<sub>2</sub><sup>•</sup> Reactive Radicals in the Gas Phase\*\*

Giulia de Petris,\* Giancarlo Angelini, Ornella Ursini, Marzio Rosi, and Anna Troiani

The activation of inert C–H bonds by charged electrophiles is the subject of extensive investigation mainly motivated by one of the most widely debated questions, that is, the conversion of alkanes into more valuable compounds. Studies of ion-molecule reactions in the idealized gaseous state have provided insightful information on this elementary step, thus lending models and concepts which span from the molecular level to the borders of the nanoscale size.<sup>[1]</sup> These studies have shown the enhanced ability of oxygen-centered radical cations to activate C–H bonds;<sup>[2]</sup> the inventory of XO<sup>++</sup> reactants now available includes metal oxides and oligomeric cluster ions, mixed metal/nonmetal cluster ions,<sup>[2]</sup> and also effective metal-free oxide ions.<sup>[3]</sup>

As a result of the H abstraction, the charge and spin of  $XO^{++}$  are separated in the products [Eqs. (1) and (2)], that is, depending on whether one or two electrons are formally transferred in the process, the  $XOH^+/R^+$  or  $XOH^+/R^+$  products are formed, respectively (RH = alkane).<sup>[4]</sup>

$$XO^{+} + RH \to XOH^{+} + R^{-}$$
(1)

$$XO^{+} + RH \to XOH^{+} + R^{+}$$
<sup>(2)</sup>

A very important implication of these thermal ionic reactions is therefore the formation of reactive radicals. The most striking example is the production of methyl radical  $CH_3$ ,  $^{[2a-c,e-j,3a,b]}$  which can dimerize to give ethane. Likewise, processes of the type shown in Eq. (2) produce oxygencentered radicals, which can undergo oxyfunctionalization in condensed and gas phases. <sup>[11,5,6]</sup> It is worth noting that surface-generated  $CH_3$  and HO radicals have been detected in the catalyzed oxidation of methane. <sup>[7]</sup> Accordingly, the nature and

[*]	Prof. Dr. G. de Petris, Dr. A. Troiani Dipartimento di Chimica e Tecnologie del Farmaco "Sapienza" Università di Roma P. le Aldo Moro 5, 00185 Roma (Italy) E-mail: giulia.depetris@uniroma1.it
	Dr. G. Angelini, Dr. O. Ursini Istituto di Metodologie Chimiche Area della Ricerca di Roma del CNR CP 10, 00016 Monterotondo Stazione (RM) (Italy)
	Prof. Dr. M. Rosi Dipartimento di Ingegneria Civile e Ambientale Università di Perugia and ISTM-CNR Via Elce di Sotto 8, 06123 Perugia (Italy)
[**]	This work is supported by the Italian Covernment and the

[\*\*] This work is supported by the Italian Government and the "Sapienza" University of Rome. The authors thank Stefania Recaldin for editorial assistance. fate of the radicals formed are central to the evolution of the environment or microenvironment where they are generated, a concept that can easily be extended to biological and atmospheric systems.

In this light, the reactivity of XO<sup>++</sup> radical cations, so far investigated with alkanes and also alkenes, indicates the formation of many radicals and neutral species in addition to CH<sub>3</sub><sup>+</sup>: C<sub>2</sub>H<sub>5</sub><sup>,[2i,3c]</sup> C<sub>2</sub>H<sub>3</sub><sup>,[2i,k,3d]</sup> P<sub>4</sub>O<sub>9</sub>(OH)<sup>,[3c]</sup> C<sub>2</sub>H<sub>4</sub><sup>,[2i,3e]</sup> C<sub>2</sub>H<sub>4</sub>O,<sup>[1c,2d,k]</sup> CH<sub>3</sub>OH.<sup>[2b,j]</sup> While a number of mechanistic details have been elucidated by mass spectrometric experiments, the nature of the uncharged products has never been experimentally probed.<sup>[8]</sup> This lack becomes a significant limitation when the C–H bond is activated by the twoelectron channel [Eq. (2)], as the signature of the H transfer rests on the radical product of the reaction.

In this work, we provide direct experimental evidence for the radical produced by an ion-molecule C-H activation reaction. The reaction between  $O_2^{\star}$  and  $CH_2F_2$  was chosen as a model of a two-electron process leading to an oxygencentered radical.  $O_2^{\star+}$  is the prototype of metal-free oxygencentered radical cations,  $CH_2F_2$  (HFC-32; HFC = hydrofluorocarbon) is a methane derivative having a C-H bond strength (102.7 kcalmol<sup>-1</sup>) quite close to that of CH<sub>4</sub>,<sup>[9]</sup> and a global warming potential approximately 30 times higher for which it is covered by the Kyoto protocol.<sup>[10,11]</sup> The radical detected is the hydroperoxy radical HO<sub>2</sub> that can dimerize to  $H_2O_2$ .  $HO_2$  is a known important player in atmospheric ozone cycles,<sup>[10]</sup> biochemical systems,<sup>[12]</sup> solution chemistry,<sup>[13]</sup> and also suggested to be involved in catalyzed oxidation of alkanes.<sup>[7d-e]</sup> The reactions of O2.+ with some HFCs have previously been studied,<sup>[14]</sup> whereas to the best of our knowledge no data is available for CH<sub>2</sub>F<sub>2</sub>. Among the oxide ions, only the vanadium oxide cluster cations have been assayed with CH<sub>2</sub>F<sub>2</sub>; in these experiments O transfer and C-F activation products were observed.<sup>[15]</sup>

In our experiments performed under the pressure of approximately  $10^{-8}$  Torr, the reaction between isolated  $O_2^{+}$  ions and  $CH_2F_2$  gives  $CHF_2^+$  as the only charged product of the C–H bond activation.

$$O_2^{\cdot +} + CH_2F_2 \rightarrow CHF_2^{+} + HO_2^{\cdot}$$
(3)

The other product observed, the  $CH_2F^+$  ion, cannot be traced to C–F activation by  $O_2^{\bullet+}$ . The kinetics exhibits the time profile typical of a consecutive reaction (Figure 1), thus showing that the primary product  $CHF_2^+$  subsequently reacts activating the C–F bond of  $CH_2F_2$ .



Figure 1. Time profiles and best-fit lines of the  $O_2^{\star+}$  ( $\blacksquare$ ,  $R^2 = 0.999$ ), CHF<sub>2</sub><sup>+</sup> ( $\blacklozenge$ ,  $R^2 = 0.994$ ) and CH<sub>2</sub>F<sup>+</sup> ( $\blacklozenge$ ,  $R^2 = 0.997$ ) ions from the reaction of  $O_2^{\star+}$  with CH<sub>2</sub>F<sub>2</sub>. *P* CH<sub>2</sub>F<sub>2</sub> = 2.4×10<sup>-8</sup> Torr.

 $\mathrm{CHF_2}^+ + \mathrm{CH_2F_2} \to \mathrm{CH_2F^+} + \mathrm{CHF_3}$ 

Accordingly,  $O_2^{\cdot+}$  selectively activates the C–H bond of CH<sub>2</sub>F<sub>2</sub> with a rate constant  $k_3 = 3.3 \times 10^{-10}$  (± 30%) cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>, which corresponds to the efficiency  $k/k_{coll} = 17\%$  ( $k_{coll} =$  collision rate). The rate constant of the consecutive reaction [Eq. (4)],  $k_4$  amounts to  $2.2 \times 10^{-10}$ (± 30%) cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>, which is in excellent agreement with the value obtained by experiments independently performed with isolated CHF<sub>2</sub><sup>+</sup> ions,  $k = 2.1 \times 10^{-10}$ (± 30%) cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>.

The formation of HO<sub>2</sub> has been directly probed by experiments, performed at high pressure, that allow detection of the intermediate of the reaction shown in Eq. (3). The stabilization and observation of these intermediates are very rare; in a few cases reported only the ionic products have been characterized.<sup>[3a,16]</sup> Here, the intermediate formally denoted as  $[C, H_2, F_2, O_2]^{+}$  has been obtained by ionization of  $O_2/$ CH<sub>2</sub>F<sub>2</sub> mixtures, using <sup>18</sup>O<sub>2</sub> to avoid mixing of the isobaric O<sub>2</sub> and CHF groups. The high-resolution collisionally activated dissociation (CAD) spectrum of isolated [C, H<sub>2</sub>, F<sub>2</sub>,  ${}^{18}O_2$ ]<sup>++</sup> ions shows two main peaks (Figure 2a):  $CHF_2^+$  and  ${}^{18}O_2^{+}$ . This pattern would indicate the presence of two complexes,  $[H^{18}O_2 - CHF_2^+]$  and  $[^{18}O_2 - CH_2F_2]$ . Alternatively, the latter complex could account for both the  $CHF_2^+$  and  ${}^{18}O_2^{+}$ fragments, considering the quite close ionization energies of  $O_2$  and  $CH_2F_2$  and the easy dissociation of  $CH_2F_2^{++}$  into  $CHF_2^+ + H^{\bullet} (\Delta H^{\circ} = 11.8 \text{ kcal mol}^{-1}).^{[9]}$  The latter is actually consistent with the very low  $CH_2F_2^{+}$  signal observed in the chemical ionization (CI) spectrum. The analysis of the neutral species coming from the dissociation of the reaction intermediate clarifies the issue, showing the nature of the radical formed in the reaction.

To this end, all neutral fragments have been separated from all charged species and reionized by the neutral fragments reionization (N<sub>f</sub>R) technique.<sup>[17]</sup> The spectrum of Figure 2b shows an intense peak at m/z 37 corresponding to the H<sup>18</sup>O<sub>2</sub><sup>+</sup> ion, that can only be formed by reionization of the H<sup>18</sup>O<sub>2</sub><sup>+</sup> radical. Notably, the failure to detect CH<sub>2</sub>F<sub>2</sub><sup>++</sup>, from reionization of the neutral counterpart of <sup>18</sup>O<sub>2</sub><sup>++</sup>, confirms its easy dissociation into CHF<sub>2</sub><sup>+</sup>. An intense peak at m/z 37 has also been obtained by negative reionization to H<sup>18</sup>O<sub>2</sub><sup>--</sup> (inset



*Figure 2.* High-resolution CAD spectrum (a) and  ${}^{+}N_{f}R^{+}$  spectrum (b) of [C, H<sub>2</sub>, F<sub>2</sub>,  ${}^{18}O_2$ ]<sup>++</sup> ions. The inset shows the  ${}^{+}N_{f}R^{-}$  spectrum with the H ${}^{18}O_2^{-}$  peak at *m/z* 37. Unresolved peaks at *m/z* 50 (CF<sub>2</sub><sup>++</sup>), 33 (CH<sub>2</sub>F<sup>+</sup>), 32 (CHF<sup>++</sup>), 31 (CF<sup>+</sup>), 19 (F<sup>+</sup>, H ${}^{18}O^{+})$ , and 18 ( ${}^{18}O^{++}$ ). The  ${}^{+}N_{f}R^{+}$  spectrum of unlabeled [C, H<sub>2</sub>, F<sub>2</sub>,  $O_2$ ]<sup>++</sup> ions display an intense signal corresponding to the HO<sub>2</sub><sup>++</sup> ion superimposed to CH<sub>2</sub>F<sup>+</sup> (*m/z* 33).

of Figure 2b). Accordingly, the positive detection of  $H^{18}O_2^+$ and  $H^{18}O_2^-$  unambiguously demonstrates that the ionic population contains a detectable amount of reaction intermediate that dissociates into  $CHF_2^+$  and  $H^{18}O_2^-$ .

Quantum chemical calculations performed at the CCSD(T)/6-311 + + G(3df,3pd)//B3LYP/6-311 + G(d) level of theory are in very good agreement with the experimental evidence. The reaction shown in Eq. (3) is computed to be exothermic by 25.5 kcalmol<sup>-1</sup>, a value quite close to that evaluated from experimental available data ( $\Delta H^{\circ} = -24$  kcal mol<sup>-1</sup>).<sup>[9]</sup> According to the energy profile picture in Figure 3, the reactants generate an encounter complex [O<sub>2</sub>-CH<sub>2</sub>F<sub>2</sub>]<sup>++</sup> (**A**) where O<sub>2</sub><sup>++</sup> interacts at a distance of 2.345 Å with the fluorine atom of CH<sub>2</sub>F<sub>2</sub>. The binding energy of [O<sub>2</sub>-CH<sub>2</sub>F<sub>2</sub>]<sup>++</sup> is 13.3 kcalmol<sup>-1</sup>.



*Figure 3.* Energy profile of the reaction between  $O_2^{\cdot+}$  and  $CH_2F_2$ .  $\Delta H^{\circ}$  values at 298 K in kcal mol<sup>-1</sup>, bond lengths in Å.

The encounter complex **A** easily isomerizes to  $[HO_2-CHF_2]^{+}$  (**B**), where the unpaired electron is entirely located on the newly formed HO<sub>2</sub> moiety interacting through the oxygen atom with the carbon atom of  $CHF_2^{+}$ . The structures of the two intermediates are consistent with the experimental

(4)

observations; in particular, the deep well of **B**, 59.1 kcal mol<sup>-1</sup> below the reactants, accounts for the experimental detection of the [HO<sub>2</sub>-CHF<sub>2</sub>]<sup>+</sup> complex in sufficient amounts to identify its dissociation radical product. Along the double-well path, the encounter complex **A** is expected to undergo back dissociation in competition with the isomerization, which is consistent with the moderate efficiency of the reaction at room temperature. The rate constant is likely to be sensitive to temperature, as observed for instance with the analogue reaction of O<sub>2</sub><sup>++</sup> with CH<sub>4</sub>.<sup>[18]</sup>

In conclusion, in this study we have given direct experimental evidence for an oxygen-centered radical formed by electrophilic C–H bond activation in the gas phase. This is the first experimental detection of both the ion and radical formed in this elementary step. The model reaction also shows that two-electron processes can generate species potentially relevant to functionalization or other chemical cycles. The ensuing chemistry strictly depends on the thermodynamic and kinetic constraints of the reaction pathways; nonetheless this result brings into sharp focus the intimate link existing between ion and neutral chemistry.

## **Experimental Section**

The N<sub>f</sub>R experiments were performed using a modified ZABSpec oa-TOF instrument (VG Micromass) of EBE-TOF configuration described elsewhere.<sup>[19]</sup> High-resolution CI spectra were recorded at 25000 full width at half maximum (fwhm) at the first detector, to rule out isobaric contaminations and assign the elemental composition [C, H<sub>2</sub>, F<sub>2</sub>, <sup>18</sup>O<sub>2</sub>]<sup>+</sup>. High-resolution CAD spectra of mass- and energyselected ions were recorded in a gas cell located in the TOF sector (0.8 keV, collision gas: Helium, transmittance T = 80%). The N<sub>f</sub>R spectra were recorded in a pair of cells located after the magnet: i) the mass-selected ions underwent high-energy (8 keV) CAD in the first cell by collision with He (T=80%); ii) the charged fragments were removed at the exit of the cell by a pair of high-voltage deflecting electrodes; iii) a beam containing only neutral fragments entered the second cell, where they were reionized by collision either with  $O_2$  $(^{+}N_{f}R^{+})$  or with Xe  $(^{+}N_{f}R^{-})$  (T = 80%). No signal was observed by switching the deflector on in the absence of the reionizing gas.

The kinetics of the ion-molecule reactions was studied by an EXTREL FTMS 2001 double-cell mass spectrometer. The O2++ ions were generated in the "source cell" by electron impact (EI) (50 ms, 30 eV) ( $P O_2 = 1.0 \times 10^{-7}$  Torr). After a cooling period of 4.1 seconds, they were isolated and transferred to the "analyzer cell" containing  $CH_2F_2$  (P  $CH_2F_2 = 2.3-9.1 \times 10^{-8}$  Torr). The pressure calibration was carried out using the known rate constant of the reference reaction  $CH_4$ .+  $CH_4$  $\rightarrow$   $CH_5^+$ + CH<sub>3</sub>  $(k=1.1\times10^{-9})$ + $(\pm$ 15%) cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>),<sup>[20]</sup> and the reading was further corrected for the response factor of  $CH_2F_2$ .<sup>[21]</sup> The  $O_2^{+}$ ,  $CHF_2^+$  and  $CH_2F^+$ intensities fit the equations derived by consecutive kinetics, where  $O_2^{+}$  fits a pseudo-first order kinectics.<sup>[22]</sup> In independent experiments, the  $CHF_2^+$  ions were generated in the analyzer cell by EI (50 ms, 18 eV) of  $CH_2F_2$  (*P*  $CH_2F_2 = 2.1-11 \times 10^{-8}$  Torr), isolated after a cooling time of 1.5 seconds and allowed to react. The bimolecular rate constants k (cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>) were obtained by the pseudo-unimolecular rate constants  $k_{(obs)}$  (relative square deviation 5%) and the neutral reactant density. The reaction efficiency  $k/k_{coll}$  ( $k_{coll}$  = collision rate constant) was calculated according to the ADO theory.[23]

Calculations were performed by locating the lowest stationary points at the B3LYP<sup>[24]</sup> level of theory with the 6-311 + G(d) basis set. Intrinsic reaction coordinate (IRC) calculations were used for the assignment of the saddle points.<sup>[25]</sup> The energy of all the stationary points was computed at the higher level of calculation CCSD(T)<sup>[26]</sup> using the 6-311 ++ G(3df,3pd) basis set and the CCSD(T) energies were corrected to 298.15 K by adding the zero point energy correction and the thermal correction computed using the scaled harmonic vibrational frequencies evaluated at B3LYP/6-311+G(d) level. All calculations were performed using Gaussian 03.<sup>[27]</sup>

Received: October 12, 2011 Published online: December 23, 2011

**Keywords:** C-H activation · density functional calculations · gas-phase reactions · mass spectrometry · radicals

- [1] a) H. Schwarz, Angew. Chem. 2011, 123, 10276; Angew. Chem. Int. Ed. 2011, 50, 10096; b) J. Roithová, D. Schröder, Chem. Rev. 2010, 110, 1170; c) G. E. Johnson, R. Mitrić, V. Bonačić-Koutecký, A. W. Castleman, Jr., Chem. Phys. Lett. 2009, 475, 1; d) R. H. Crabtree, Nat. Chem. 2009, 1, 348; e) D. K. Böhme, H. Schwarz, Angew. Chem. 2005, 117, 2388; Angew. Chem. Int. Ed. 2005, 44, 2336. For further reading about electrophilic activation of hydrocarbons, see: f) G. A. Olah, Acc. Chem. Res. 1987, 20, 422; g) J. J. Schneider, Angew. Chem. 1996, 108, 1132; Angew. Chem. Int. Ed. Engl. 1996, 35, 1068; h) A. E. Shilov, G. B. Shul'pin, Chem. Rev. 1997, 97, 2879; i) R. A. Periana, D. J. Taube, S. Gamble, H. Taube, T. Satoh, H. Fujii, Science 1998, 280, 560; j) S. S. Stahl, J. A. Labinger, J. E. Bercaw, Angew. Chem. 1998, 110, 2298; Angew. Chem. Int. Ed. 1998, 37, 2180; k) R. H. Crabtree, J. Chem. Soc. Dalton Trans. 2001, 2437; 1) A. A. Fokin, P. R. Schreiner, Chem. Rev. 2002, 102, 1551; m) J. A. Labinger, J. Mol. Catal. A 2004, 220, 27.
- [2] a) S. Feyel, J. Döbler, D. Schröder, J. Sauer, H. Schwarz, Angew. Chem. 2006, 118, 4797; Angew. Chem. Int. Ed. 2006, 45, 4681; b) D. Schröder, J. Roithovà, Angew. Chem. 2006, 118, 5835; Angew. Chem. Int. Ed. 2006, 45, 5705; c) S. Feyel, J. Döbler, R. Höckendorf, M. K. Beyer, J. Sauer, H. Schwarz, Angew. Chem. 2008, 120, 1972; Angew. Chem. Int. Ed. 2008, 47, 1946; d) G. E. Johnson, R. Mitrić, E. C. Tyo, V. Bonačić-Koutecký, A. W. Castleman, Jr., J. Am. Chem. Soc. 2008, 130, 13912; e) A. Božović, D. K. Böhme, Phys. Chem. Chem. Phys. 2009, 11, 5940; f) Z.-C. Wang, X.-N. Wu, Y.-X. Zhao, J.-B. Ma, X.-L. Ding, S.-G. He, Chem. Phys. Lett. 2010, 489, 25; g) X.-L. Ding, Y.-X. Zhao, X.-N. Wu, Z.-C. Wang, J.-B. Ma, S.-G. He, Chem. Eur. J. 2010, 16, 11463; h) Y.-X. Zhao, X.-N. Wu, Z.-C. Wang, S.-G. He, X.-L. Ding, Chem. Commun. 2010, 46, 1736; i) N. Dietl, R. F. Höckendorf, M. Schlangen, M. Lerch, M. K. Beyer, H. Schwarz, Angew. Chem. 2011, 123, 1466; Angew. Chem. Int. Ed. 2011, 50, 1430; j) N. Dietl, C. van der Linde, M. Schlangen, M. K. Beyer, H. Schwarz, Angew. Chem. 2011, 123, 5068; Angew. Chem. Int. Ed. 2011, 50, 4966; k) K. Chen, Z.-C. Wang, M. Schlangen, Y.-D. Wu, X. Zhang, H. Schwarz, Chem. Eur. J. 2011, 17, 9619.
- [3] a) G. de Petris, A. Troiani, M. Rosi, G. Angelini, O. Ursini, *Chem. Eur. J.* 2009, 15, 4248; b) N. Dietl, M. Engeser, H. Schwarz, Angew. Chem. 2009, 121, 4955; Angew. Chem. Int. Ed. 2009, 48, 4861; c) N. Dietl, M. Engeser, H. Schwarz, Chem. Eur. J. 2009, 15, 11100; d) N. Dietl, M. Engeser, H. Schwarz, Chem. Eur. J. 2010, 16, 4452; e) G. de Petris, A. Cartoni, A. Troiani, V. Barone, P. Cimino, G. Angelini, O. Ursini, Chem. Eur. J. 2010, 16, 6234.
- [4] One- and two-electron processes strictly correspond to H<sup>•</sup> and H<sup>-</sup> transfer, respectively; however they may not occur by pure homolytic and heterolytic rupture of the C–H bond when innersphere electron transfer is involved (see Ref. [11]).
- [5] A. A. Fokin, P. R. Schreiner, Adv. Synth. Catal. 2003, 345, 1035, and references therein.
- [6] F. Dong, S. Heinbuch, Y. Xie, J. J. Rocca, E. R. Bernstein, Z.-C. Wang, K. Deng, S.-G. He, J. Am. Chem. Soc. 2008, 130, 1932.

## Angewandte Communications

- [7] a) J. H. Lunsford, Langmuir 1989, 5, 12; b) C. E. Mooney, L. C. Anderson, J. H. Lunsford, J. Phys. Chem. 1991, 95, 6070; c) J. H. Lunsford, Angew. Chem. 1995, 107, 1059; Angew. Chem. Int. Ed. Engl. 1995, 34, 970; d) F. Cavani, F. Trifirò, Catal. Today 1999, 51, 561; e) G. B. Shul'pin, Y. N. Kozlov, G. V. Nizova, G. Süss-Fink, S. Stanislas, A. Kitaygorodskiy, V. S. Kulikova, J. Chem. Soc. Perkin Trans. 2 2001, 1351.
- [8] In the Pt<sup>+</sup>-catalyzed coupling of CH<sub>4</sub> and NH<sub>3</sub>, the suggested intermediate CH<sub>2</sub>NH<sup>•</sup> has been identified by threshold ionization mass spectrometry; a) M. Aschi, M. Brönstrup, M. Diefenbach, J. N. Harvey, D. Schröder, H. Schwarz, *Angew. Chem.* **1998**, *110*, 858; *Angew. Chem. Int. Ed.* **1998**, *37*, 829; b) R. Horn, G. Mestl, M. Thiede, F. C. Jentoft, P. M. Schmidt, M. Bewersdorf, R. Weber, R. Schlögl, *Phys. Chem. Chem. Phys.* **2004**, *6*, 4514.
- [9] a) NIST Chemistry WebBook, NIST Standard Reference Database Number 69, Gaithersburg MD, 20899, http://webbook.nist.gov (Eds.: P. J. Linstrom, W. G. Mallard), 2005; b) M. W. Chase, Jr., NIST-JANAF Themochemical Tables, Fourth Edition, J. Phys. Chem. Ref. Data, Monograph 9, 1998, 1–1951.
- [10] G. P. Brasseur, J. J. Orlando, G. S. Tyndall, Atmospheric Chemistry and Global Change, Oxford University Press, Oxford, 1999.
- [11] a) P. Blowers, K. Hollingshead, J. Phys. Chem. A 2009, 113, 5942;
   b) http://unfccc.int/resource/docs/convkp/kpeng.pdf.
- [12] a) A. D. N. J. De Grey, DNA Cell Biol. 2002, 21, 251; b) P. Wentworth, Jr., A. D. Wentworth, X. Zhu, I. A. Wilson, K. D. Janda, A. Eschenmoser, R. A. Lerner, Proc. Natl. Acad. Sci. USA 2003, 100, 1490.
- [13] M. Toru, Y. Ichiro, J. Phys. Chem. C 2011, 115, 5792.
- [14] a) A. B. Raksit, *Int. J. Mass Spectrom. Ion Processes* 1986, 69, 45;
   b) R. A. Morris, A. A. Viggiano, S. T. Arnold, J. F. Paulson, *J. Phys. Chem.* 1995, 99, 5992.

- [15] R. C. Bell, K. A. Zemski, A. W. Castleman, Jr., J. Phys. Chem. A 1999, 103, 2992.
- [16] G. de Petris, S. Garzoli, A. Troiani, Chem. Phys. Lett. 2007, 435, 219.
- [17] C. A. Schalley, G. Hornung, D. Schröder, H. Schwarz, *Chem. Soc. Rev.* **1998**, 27, 91.
- [18] At room temperature the reaction between  $O_2^{++}$  and  $CH_4$  is very slow ( $k \approx 5 \times 10^{-12}$  cm<sup>3</sup>s<sup>-1</sup>molecule<sup>-1</sup>) and gives the insertion product CH<sub>2</sub>OOH<sup>+</sup>. The rate constant increases up to  $5 \times 10^{-10}$  cm<sup>3</sup>s<sup>-1</sup> molecule<sup>-1</sup> below 300 K, and to  $7.4 \times 10^{-12}$  cm<sup>3</sup>s<sup>-1</sup> molecule<sup>-1</sup> above 300 K. See: a) E. R. Fisher, P. B. Armentrout, *J. Chem. Phys.* **1991**, *94*, 1150; b) S. Irle, K. Morokuma, *J. Chem. Phys.* **2001**, *114*, 6119, and references therein.
- [19] F. Bernardi, F. Cacace, G. de Petris, F. Pepi, I. Rossi, A. Troiani, *Chem. Eur. J.* 2000, 6, 537.
- [20] V. G. Anicich, J. Phys. Chem. Ref. Data 1993, 22, 1469.
- [21] J. E. Bartmess, R. M. Georgiadis, Vacuum 1983, 33, 149.
- [22] G. de Petris, A. Cartoni, A. Troiani, G. Angelini, O. Ursini, *Phys. Chem. Chem. Phys.* 2009, 11, 9976.
- [23] M. T. Bowers, T. Su, Interactions between Ions and Molecules, Plenum Press, New York, 1975.
- [24] a) A. D. Becke, J. Chem. Phys. 1993, 98, 5648; b) P. J. Stephens, F. J. Devlin, C. F. Chabalowski, M. J. Frisch, J. Phys. Chem. 1994, 98, 11623.
- [25] a) C. Gonzalez, H. B. Schlegel, J. Chem. Phys. 1989, 90, 2154;
   b) C. Gonzalez, H. B. Schlegel, J. Phys. Chem. 1990, 94, 5523.
- [26] a) R. J. Bartlett, Annu. Rev. Phys. Chem. 1981, 32, 359; b) K.
   Raghavachari, G. W. Trucks, J. A. Pople, M. Head-Gordon, Chem. Phys. Lett. 1989, 157, 479; c) J. Olsen, P. Jorgensen, H.
   Koch, A. Balkova, R. J. Bartlett, J. Chem. Phys. 1996, 104, 8007.
- [27] M. J. Frisch, et al. Gaussian 03, Revision D.01, Gaussian, Inc.; Wallingford, CT, 2004.