

Phorboxazole synthetic studies: the C3–C15 bis-oxane segment

Patrick B. Greer and William A. Donaldson *


Department of Chemistry, Marquette University, P.O. Box 1881, Milwaukee, Wisconsin 53201-1881, USA

Received 23 February 2000; revised 28 March 2000; accepted 30 March 2000

Abstract

The enantioselective synthesis of the C3–C15 bis-oxane segment of the phorboxazoles has been accomplished from 3-*t*-butyldiphenylsilyloxypropanal in 9 steps (>90% ee). © 2000 Elsevier Science Ltd. All rights reserved.


The phorboxazoles A and B (**1a** and **b**) are isomeric macrolides isolated from the marine sponge *Phorbas* sp.^{1a} In addition to exhibiting antifungal activity against *Candida albicans*, both phorboxazoles gave mean GI₅₀s < 8 × 10^{−10} M against most of the 60 tumor cell lines in the NCI panel. The complex structure of **1** consists of four oxane rings, two oxazole rings, 15 asymmetric centers, and a conjugated diene segment. The complete structure was assigned on the basis of NMR spectral studies, Mosher ester data, degradation, and model synthesis.¹ The outstanding biological activity of **1**, combined with its complex structural architecture, has led to significant activity by a number of research groups,² and a total synthesis by Forsyth's group.³ Recent reports by Hoffmann,^{2c} Smith^{2d} and Pattenden^{2e} have prompted us to report our own efforts on the synthesis of the C3–C15 bis-oxane segment **2**.



Lewis acid catalysed diene-aldehyde cyclocondensation⁴ of **3** with **4** in the presence of BF₃·Et₂O, followed by brief treatment with CF₃CO₂H gave dihydropyrone *rac*-**5** (Scheme 1). In our hands attempted asymmetric cyclocondensation by a number of literature procedures proved capricious.⁵ For this reason, establishment of optical activity was deferred to a later stage. Oxymercuration/reduction of **5** followed by reduction [LiAlH(OtBu)₃] and acylation gave **6**. Reaction of **6** with allyl

* Corresponding author.

trimethylsilane/TMSOTf/CH₃CN⁶ gave the α -C-glucoside **7**, which upon ozonolysis gave the aldehyde *rac*-**8**. The synthesis of a similar aldehyde was reported by Hoffmann's group^{2c} beginning from 8-oxabicyclo[3.2.1]octan-3-one (11 steps), and by Smith's group^{2d} beginning from 2,4-pentandione (12 steps); our route to **8** is significantly shorter (six steps).



Scheme 1. (P=TBDPS)

Chiral allylation of *rac*-**8** with *B*-allyldiisopinocampheylborane (prepared from (–)-(IPC)₂BOMe),⁷ followed by oxidative work up with NaBO₃, gave a separable mixture of the diastereomers (+)-**9** and (–)-**10** (41 and 42%, respectively). The absolute stereochemistry of each alcohol at C11 was assigned as (*S*) on the basis of the relative chemical shifts of their (*R*)- and (*S*)-Mosher's esters (>90% de each).

Ring closing metathesis⁸ has recently found utility in the preparation of unsaturated δ -lactones.⁹ To this end, esterification of (–)-**10** with acryloyl chloride gave (–)-**11**, which upon treatment with Grubbs' catalyst (0.3 equivalents) gave the unsaturated pyrone (–)-**2**.¹⁰

In summary, the C3–C15 bis-oxane segment of the phorboxazoles has been prepared in nine steps. Studies directed toward the preparation of other portions of the phorboxazoles continues in our laboratory.

Acknowledgements

The authors are grateful to the Marquette University Graduate School and the Marquette University Research Incentive Fund for financial support. P.B.G. is the recipient of a Johnson Wax research fellowship. Partial financial support was provided by the National Institutes of Health (GM-42641). The high resolution mass-spectral determinations were made at the Washington University Resource for Mass Spectrometry.

References

- (a) Searle, P. A.; Molinski, T. F. *J. Am. Chem. Soc.* **1995**, *117*, 8126–31; (b) Searle, P. A.; Molinski, T. F.; Brzezinski, L. J.; Leahy, J. W. *J. Am. Chem. Soc.* **1996**, *118*, 9422–23; (c) Molinski, T. F. *Tetrahedron Lett.* **1996**, *37*, 7897–80.
- C3–C15 bispyran segment: (a) Cink, R. D.; Forsyth, C. J. *J. Org. Chem.* **1997**, *62*, 5672–73; (b) Williams, D. R.; Clark, M. P.; Berliner, M. A. *Tetrahedron Lett.* **1999**, *40*, 2287–90; (c) Wolbers, P.; Hoffmann, H. M. R. *Tetrahedron* **1999**, *55*, 1905–14; (d) Smith III, A. B.; Verhoest, P. R.; Minbiole, K. P.; Lim, J. J. *Org. Lett.* **1999**, *1*, 909–12; (e) Pattenden, G.; Plowright, A. T. *Tetrahedron Lett.* **2000**, *41*, 983–86. For other references see Ref. 2e.

3. Forsyth, C. J.; Ahmed, F.; Cink, R. D.; Lee, C. S. *J. Am. Chem. Soc.* **1998**, *120*, 5597–98.
4. Danishefsky, S.; Bilodeau, M. T. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1380–1419.
5. On one occasion asymmetric LACDAC using the Keck procedure gave (–)-**5** (ca. 90% ee, 19% yield). Numerous attempts to repeat this result were unsuccessful. (a) Keck, G. E.; Li, X.-Y.; Krishnamurthy, D. *J. Org. Chem.* **1995**, *60*, 5998–99; (b) Bednarski, M.; Danishefsky, S. *J. Am. Chem. Soc.* **1986**, *108*, 7060–67; (c) Togni, A. *Organometallics* **1990**, *9*, 3106–13; (d) Ghosh, A. K.; Mathivanan, P.; Cappiello, J.; Krishnan, K. *Tetrahedron: Asymmetry* **1996**, *7*, 2165–68.
6. Hosomi, A.; Sakata, Y.; Sakurai, H. *Tetrahedron Lett.* **1984**, *25*, 2383–86.
7. Brown, H. C.; Jadhav, P. K. *J. Am. Chem. Soc.* **1983**, *105*, 2092–93.
8. Schuster, M.; Blechert, S. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2037–56; Grubbs, R. H.; Chang, S. *Tetrahedron* **1998**, *54*, 4413–50.
9. Nicolaou, K. C.; Rodriguez, R. M.; Mitchell, H. J.; van Delft, F. L. *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 1874–76; Ghosh, A. K.; Cappiello, J.; Shin, D. *Tetrahedron Lett.* **1998**, *39*, 4651–54; Cossey, J.; Bauer, D.; Bellosta, V. *Tetrahedron Lett.* **1999**, *40*, 4187–88; Ramachandran, P. V.; Reddy, M. V. R.; Brown, H. C. *Tetrahedron Lett.* **2000**, *41*, 583–86.
10. Compound (–)-**2**: $[\alpha]_D -38$ (c 1.0, CHCl₃); ¹H NMR (CDCl₃) δ 7.68–7.61 (m, 4H), 7.46–7.32 (m, 6H), 6.77 (ddd, $J=3.3, 5.3, 9.6$ Hz, 1H), 5.97 (br d, $J=10.1$ Hz, 1H), 5.06 (tt, $J=4.5, 9.0$ Hz, 1H), 4.52 (m, 1H), 4.24 (dq, $J=4.6, 9.1$ Hz, 1H), 3.97 (tt, $J=4.2$ Hz, 8.4 Hz, 1H), 3.81–3.63 (m, 2H), 2.36 (m, 2H), 2.28 (ddd, $J=5.5, 9.3, 15.3$ Hz, 1H), 2.03 (s, 3H), 1.97 (td, $J=3.1, 12.7$ Hz, 1H), 1.86–1.62 (m, 5H), 1.39 (td, $J=9.2, 12.9$ Hz, 1H), 1.02 (s, 9H); ¹³C NMR (CDCl₃) δ 170.3, 164.1, 144.9, 135.5, 133.8, 129.6, 127.7, 121.4, 75.0, 67.2, 66.4, 65.9, 60.2, 38.1, 36.8, 36.2, 34.4, 28.7, 26.8, 21.2, 19.2.